

AD _____

CONTRACT NO.: DAMD17-92-C-2066

TITLE: Biomonitoring and Hazard Assessment Evaluation of
Contaminated Groundwater at Aberdeen Proving Ground-
Edgewood Area Beach Point Peninsula

PRINCIPAL INVESTIGATOR: Dennis T. Burton, Ph.D.

CO-INVESTIGATORS: Randall S. Herriott, B.S. and Steven D.
Turley, M.S.

CONTRACTING ORGANIZATION: University of Maryland at College Park
Agricultural Experiment Station
Wye Research and Education Center
P.O. Box 169
Queenstown, MD 21658

REPORT DATE: August 30, 1994

TYPE OF REPORT: Annual Report (Report Period - July 31, 1993 to
July 30, 1994)

PREPARED FOR: U.S. Army Medical Research and
Materiel Command
Fort Detrick
Frederick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release
Distribution Unlimited

The views, opinions, and/or findings contained in this report are
those of the authors and should not be construed as an official
Department of the Army position, policy, or decision unless so
designated by other documents.

19941215 150

| REPORT DOCUMENTATION PAGE | | | Form Approved OMB No. 0704-0188 | |
|---|--|--|--|--|
| <small>Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204 Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.</small> | | | | |
| 1. AGENCY USE ONLY (Leave blank) | 2. REPORT DATE August 30, 1994 | 3. REPORT TYPE AND DATES COVERED Annual Report, 31 Jul 93 - 30 Jul 94 | | |
| 4. TITLE AND SUBTITLE Biomonitoring and Hazard Assessment Evaluation of Contaminated Groundwater at Aberdeen Proving Ground-Edgewood Area Beach Point Peninsula | | | 5. FUNDING NUMBERS DAMD17-92-C-2066 | |
| 6. AUTHOR(S) Dennis T. Burton, Randall S. Herriott, and Steven D. Turley | | | | |
| 7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) The University of Maryland System Agricultural Experiment Station Wye Research and Education Center P.O. Box 169 Queenstown, MD 21658 | | | 8. PERFORMING ORGANIZATION REPORT NUMBER WREC-94-B3 | |
| 9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick Frederick, Maryland 21702-5012 | | | 10. SPONSORING/MONITORING AGENCY REPORT NUMBER | |
| 11. SUPPLEMENTARY NOTES | | | | |
| 12a. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited. | | | 12b. DISTRIBUTION CODE | |
| 13. ABSTRACT (Maximum 200 words) Contaminated groundwater, which contained multiple heavy metals and chlorinated aliphatic hydrocarbons, from the surficial aquifer (well CC-33B) at Beach Point located in the Canal Creek Area of the U.S. Army Aberdeen Proving Ground-Edgewood Area, Aberdeen, Maryland, was evaluated for toxicity and environmental hazard. Toxicity was detected at various groundwater concentrations by 7 of 9 biomonitoring systems. When estimated maximum acceptable toxicant concentrations (MATC) were established, the data for algae, invertebrates and fish suggested that the groundwater would not be harmful at a concentration of 10% groundwater by volume. Likewise, no genotoxicity (Ames and SEC assays), developmental toxicity (FETAX), or chronic histopathology (9-month fish test) occurred at 10% groundwater by volume. The groundwater was considered to be a potentially excessive hazardous material to the benthic biota of the Bush River when a number of conservative assumptions (contaminant distribution and discharge rate of the aquifer) were used in the hazard assessment. However, the potential water quality impacts were judged to be minimal if a mixing zone were to be granted by the State of Maryland which allows for local exceedences of water quality standards. (Cont on reverse) | | | | |
| 14. SUBJECT TERMS Biomonitoring, bioassays, acute toxicity, chronic toxicity, Ames, SCE, FETAX, histopathology, groundwater, priority pollutants, heavy metals, chlorinated aliphatic hydrocarbons, hazard assessment, near-field dilution model, (Cont on reverse) | | | 15. NUMBER OF PAGES 652 | |
| | | | 16. PRICE CODE | |
| 17. SECURITY CLASSIFICATION OF REPORT Unclassified | 18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified | 19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified | 20. LIMITATION OF ABSTRACT Unclassified | |

13. ABSTRACT (Cont)

Near-field (ULINE model) and far-field (dye-tracer model) screening level dilution models were run to estimate the dilution of the groundwater discharge plume in the Bush River. The model analyses showed that a near-field dilution level of approximately 42:1 for the application of Maryland's acute aquatic life criteria and a near-field dilution level of 168:1 for the application of chronic criteria would occur. When the dilution factors were applied to groundwater quality at Beach Point, none of the heavy metals or chlorinated aliphatic compounds exceeded Maryland's current acute or chronic aquatic life criteria.

14. SUBJECT TERMS (Cont)

ULINE, far-field dilution model, dye tracer model, acute aquatic life criteria, and chronic aquatic life criteria.

FORWARD

Opinions, interpretations, conclusions, and recommendations are those of the authors and are not necessarily endorsed by the U.S. Army.

- () Where copyrighted material is quoted, permission has been obtained to use such material.
- () Where material from documents designated for limited distribution is quoted, permission has been obtained to use the material.
- (✓) Citations of commercial organizations and trade names in this report do not constitute an official Department of the Army endorsement or approval of the products or services of these organizations.
- (✓) In conducting research using animals, the investigator(s) adhered to the "Guide for the Care and Use of Laboratory Animals of the Institute of Laboratory Animal Resources, National Research Council (NIH Publication No. 86-23, Revised 1985).
- () In conducting research utilizing recombinant DNA technology, the investigator(s) adhered to current guidelines promulgated by the National Institutes of Health.

P. T. Burton
Principal Investigator's Signature

8-30-94
Date


| | |
|----------------------|--|
| Accession For | |
| NTIS CRA&I | <input checked="checked" type="checkbox"/> |
| DTIC TAB | <input type="checkbox"/> |
| Unannounced | <input type="checkbox"/> |
| Justification _____ | |
| By _____ | |
| Distribution / _____ | |
| Availability Codes | |
| Dist | Avail and/or Special |
| A-1 | |

Certification of Technical Data Conformity (May 1987)

The contractor, The University of Maryland System, hereby certifies that, to the best of its knowledge and belief, the technical data delivered herewith under Contract No. DAMD17-92-C-2066 is complete, accurate, and complies with all requirements of the contractor.

Date: August 22, 1994

Name and Title of Certifying Official:


Dennis T. Burton, Ph.D
Senior Research Scientist

EXECUTIVE SUMMARY

The primary objective of the study was to evaluate the use of several biomonitoring systems for assessing contaminated groundwater. Contaminated groundwater, which contained multiple heavy metals and chlorinated aliphatic hydrocarbons, was studied. The contaminated groundwater was withdrawn from the lower depths of the surficial aquifer at Beach Point which is located in the Canal Creek Area of the U.S. Army Aberdeen Proving Ground-Edgewood Area, Aberdeen, MD. Groundwater was withdrawn from well CC-33B, which is the most the highly contaminated well at Beach Point. A hazard assessment of the groundwater release into the Bush River was performed using the biomonitoring results.

Biomonitoring Evaluation

Several EPA priority pollutant heavy metals were found in the groundwater at Beach Point. The concentrations of cadmium, copper, nickel, and zinc exceeded one or more of EPA's numerical water quality criteria for the specific metal. A number of chlorinated aliphatic compounds were also found in the groundwater, several of which are EPA priority pollutants. The chlorinated aliphatic organics that occur at the lower depths of the aquifer may be present as a residual denser-than-water nonaqueous phase liquid (DNAPL). Of the organics present in the groundwater, none had octanol water partition coefficients ($\log k_{ow}$ or $\log P$) >3 ; thus, bioaccumulation was not a potential toxicological problem.

An array of nine biomonitoring systems integrated into a tiered hazard framework was evaluated during a 9-month study. The biomonitoring systems included a number of endpoints. The pH of the groundwater from well CC-33B was 4 (± 0.1); thus, many of the assays were conducted at both pH 4 and pH 7. The toxicity at pH 7 was studied so that the data could be used in the hazard assessment of the groundwater as it entered the Bush River which has pH values close to the neutral range.

Toxicity was detected at various groundwater concentrations by 7 of 9 biomonitoring systems. The Ames assay for mutagenicity was negative in all cases (1, 10, and 100% groundwater by volume). Japanese medaka (*Oryzias latipes*) growth was not affected by 9 months of exposure to 1 and 10% groundwater by volume in a chronic histopathology assay. No significant lesions were found in the Japanese medaka exposed to groundwater concentrations up to 10% groundwater by volume (highest concentration studied). A positive response was found for 100% groundwater in a sister chromosome exchange (SCE) assay for DNA damage when the groundwater sample was concentrated $\approx 50,000X$. The SCE assay was negative for unconcentrated 100% groundwater. The positive SCE response in the 50,000X concentrated sample was

judged not to be important to aquatic organisms in the receiving stream.

The lowest concentration of groundwater that caused no observable adverse effect (NOEC) at pH 4, in the test systems in which the NOEC value could be determined, was 10% groundwater by volume. A NOEC of 10% groundwater by volume occurred in 3 out of 4 tests in both a 7-d cladoceran (Ceriodaphnia dubia) and a 96-h frog (Xenopus laevis) embryo teratogenesis assay - Xenopus (FETAX); it occurred once in a 7-d fathead minnow (Pimephales promelas) test. The NOEC concentration was higher at pH 7 in both the fathead minnow and FETAX assays. The 10% groundwater by volume NOEC for the cladoceran at pH 4, however, did not change when the organism was exposed to buffered groundwater at pH 7.

When estimated maximum acceptable toxicant concentrations (MATC) were established, the data for alga (Selenastrum capricornutum), invertebrate (C. dubia), and fish (P. promelas) used in the biomonitoring study suggested that the groundwater would not be harmful at a concentration of 10% groundwater by volume. Likewise, no genotoxicity, developmental toxicity, or chronic histopathology occurred at 10% groundwater by volume. Thus, the biomonitoring data suggested that chronic toxicity may not occur in the Bush River at a dilution of 10:1. Chronic toxicity was predicted to occur in the Bush River if the groundwater entered the receiving stream at the concentrations which occur in well CC-33B.

Hazard Assessment Evaluation

A number of worst-case assumptions were made for the Beach Point hazard assessment when sufficient data were not available. The following assumptions were made for the calculation of the estimated discharge rate of the surficial aquifer into the Bush River. The surficial aquifer was considered to be homogeneous, characterized by isotropic flow conditions. The average saturated thickness of the aquifer was assumed to be 18.8 m (61.8 ft). The aquifer was assumed to discharge the entire length of the Beach Point Peninsula fronting the Bush River; a more realistic estimate of length of discharge from Beach Point is probably one third to one half, rather than the entire length.

With regard to the contaminant concentrations in the groundwater, the assumption was made for heavy metals that no retardation occurred via adsorption onto solid surfaces or trapping by clays through ion exchange. It was also assumed that no precipitation of the metals occurred when the pH shifted from 4 to neutrality when the groundwater entered the receiving stream. The assumption was made for the DNAPLs that no abiotic (chemical) or biotic (microbial) transformations occurred.

It was also assumed that the heavy metals and residual DNAPLs were homogeneously mixed throughout the complete aquifer at the highest concentration (not the average concentration) measured during the course of the study. It was assumed that the maximum concentrations of materials would not increase above those currently present in well CC-33B since the original sources of the contaminants were not longer present. The assumption was made that the highest concentrations of the heavy metals and chlorinated aliphatics in the aquifer all moved through the sediments into the Bush River without any biotransformations or other physical/chemical transformations occurring in the sediment or water column of the Bush River.

When the above assumptions were used in the hazard assessment, the groundwater may be considered to be a potentially hazardous material to the benthic biota of the Bush River. The hazard to water column aquatic biota would rapidly dissipate as the groundwater materials mix in the receiving stream. Because the potential water quality impacts were judged to be minimal, a mixing zone approach by the State of Maryland which allows for local exceedences of water quality standards may be pursued. The State of Maryland Code (COMAR) allows for a mixing zone on a case-by-case basis. Pollutant concentrations within a mixing zone may exceed the specified water quality standards within a localized vicinity of an outfall.

Near-field (ULINE model) and far-field (dye-tracer model with input data originally obtained for the Bush River) EPA-approved screening level dilution models were run to estimate the dilution of the groundwater discharge plume in the Bush River. The concentrations of the groundwater hazardous substances were estimated for the near-field and far-field in the Bush River using the mixing zone requirements of the State of Maryland.

The model analyses showed that a total near-field dilution (near-field dilution corrected for the influence of far-field dilution) of approximately 42:1 for the application of the State's acute aquatic life criteria and a near-field dilution level of 168:1 for the application of chronic criteria would occur. Thus, contaminants introduced via Beach Point groundwater into Bush River receiving waters at a concentration of 42 $\mu\text{g/L}$ would be diluted locally to a concentration of approximately 1 $\mu\text{g/L}$ or less. When the dilution factors were applied to groundwater quality at Beach Point, none of the heavy metals or chlorinated aliphatic compounds exceeded Maryland's current acute or chronic aquatic life criteria. The dilution study showed that detectable concentrations of some of the chlorinated organics would occur in the receiving stream when the conservative assumptions concerning the groundwater contaminants were used in the model.

Although an argument can be made for a mixing zone, it is highly unlikely that the contaminant loads assumed in the near-field dilution model will ever occur in the Bush River. A refinement of the conservative assumptions used in the hazard assessment would reduce the uncertainty regarding the volume of the discharge as well as the concentration of contaminants that may enter the receiving stream.

ACKNOWLEDGEMENTS

We thank Mr. Tommy R. Shedd (COR), Mr. Henry S. Gardner, and Dr. Robert A. Finch of the U.S. Army Medical Research and Development Command (USAMRDC), Fort Detrick, Frederick, Maryland, for their continual support and help throughout the study. Dr. Alan B. Rosencrance of USAMRDC is acknowledged for conducting the munitions analyses. We would also like to thank Mr. John G. Wrobel of the Environmental Preservation, Conservation and Restoration Division of the Directorate of Safety, Health and Environment, Aberdeen Proving Ground, MD for his help during the study. Mr. Scott Treherne of Jacobs Engineering Group provided the Beach Point surficial aquifer discharge rate calculation. Dr. Joseph L. DiLorenzo, Najarian Associates, Inc., performed the near- and far-field dilution models of the Beach Point groundwater discharge to the Bush River. We acknowledge the Research Methods Branch/Health Effects Research Division of USABRDL for supporting the study through U.S. Army Contract No. DAMD17-92-C-2066. This report is Contribution No. 8824 from the Maryland Agricultural Experiment Station.

TABLE OF CONTENTS

| | |
|---|----|
| FORWARD..... | 1 |
| CERTIFICATION OF TECHNICAL DATA CONFORMITY..... | 2 |
| EXECUTIVE SUMMARY..... | 3 |
| ACKNOWLEDGEMENTS..... | 7 |
| TABLE OF CONTENTS..... | 8 |
| LIST OF TABLES..... | 11 |
| LIST OF FIGURES..... | 12 |
| LIST OF APPENDICES..... | 13 |
| 1. INTRODUCTION..... | 18 |
| 2. OBJECTIVES OF STUDY..... | 19 |
| 3. BEACH POINT SITE DESCRIPTION..... | 20 |
| 3.1 Physiographic Setting..... | 20 |
| 3.2 Historical Use..... | 20 |
| 3.3 Groundwater Contamination..... | 20 |
| 3.4 Geology..... | 21 |
| 3.5 Groundwater Hydrology..... | 22 |
| 3.6 Surface Water Hydrology..... | 23 |
| 4. MATERIALS AND METHODS..... | 24 |
| 4.1 Biomonitoring Evaluation..... | 24 |
| 4.1.1 Acute Toxicity Tests..... | 28 |
| 4.1.1.1 Microtox®..... | 28 |
| 4.1.1.2 Rotifer..... | 29 |
| 4.1.1.3 Green Alga, Cladoceran, Fathead Minnow, and Japanese Medaka..... | 29 |
| 4.1.2 Short-term Chronic Toxicity Tests..... | 30 |
| 4.1.2.1 Green Alga..... | 30 |
| 4.1.2.2 Cladoceran..... | 31 |
| 4.1.2.3 Fathead Minnow..... | 31 |
| 4.1.3 Genotoxicity Tests..... | 32 |
| 4.1.3.1 Gene Mutation Assay..... | 32 |
| 4.1.3.2 Sister Chromatid Exchange Assay..... | 33 |
| 4.1.4 Developmental Toxicity Test..... | 34 |
| 4.1.5 Chronic Histopathology and Growth Test..... | 35 |
| 4.1.6 Chemical Analyses..... | 37 |
| 4.1.6.1 Comprehensive Chemical Analyses..... | 37 |
| 4.1.6.2 Routine Water Quality Analyses..... | 37 |

TABLE OF CONTENTS (CONTINUED)

| | | |
|---------|---|----|
| 4.1.7 | Test Endpoints and Data Analyses..... | 38 |
| 4.2 | Hazard Assessment Evaluation..... | 40 |
| 4.2.1 | Scope of the Hazard Assessment..... | 40 |
| 4.2.2 | Deviations from the Hazard Assessment Procedures..... | 41 |
| 4.2.3 | Estimated Discharge Rate of the Surficial Aquifer..... | 42 |
| 4.2.4 | Contaminant Distribution Assumptions in the Groundwater..... | 44 |
| 4.2.5 | Screening-level Assessment of the Near-Field Dilution of the Groundwater Plume in Bush River..... | 44 |
| 4.2.5.1 | Near-field Dilution Estimates..... | 45 |
| 4.2.5.2 | Far-field Dilution Estimates..... | 46 |
| 4.2.5.3 | Mixing Zone Consideration..... | 47 |
| 5. | RESULTS AND DISCUSSION..... | 48 |
| 5.1 | Biomonitoring Evaluation..... | 48 |
| 5.1.1 | Acute Toxicity Tests..... | 48 |
| 5.1.1.1 | Microtox®..... | 49 |
| 5.1.1.2 | Rotifer..... | 56 |
| 5.1.1.3 | Green Alga, Cladoceran, Fathead Minnow, and Japanese Medaka..... | 56 |
| 5.1.2 | Short-term Chronic Toxicity Tests..... | 60 |
| 5.1.2.1 | Green Alga..... | 60 |
| 5.1.2.2 | Cladoceran..... | 60 |
| 5.1.2.3 | Fathead Minnow..... | 61 |
| 5.1.3 | Genotoxicity Tests..... | 62 |
| 5.1.3.1 | Gene Mutation Assay..... | 62 |
| 5.1.3.2 | Sister Chromatid Exchange Assay..... | 62 |
| 5.1.4 | Developmental Toxicity Test..... | 63 |
| 5.1.5 | Chronic Histopathology and Growth Test..... | 64 |
| 5.1.6 | Chemical Analyses..... | 69 |
| 5.1.6.1 | Comprehensive Chemical Analyses..... | 69 |
| 5.1.6.2 | Routine Water Quality Analyses..... | 70 |
| 5.2 | Hazard Assessment Evaluation..... | 71 |
| 5.2.1 | Toxicity..... | 71 |
| 5.2.2 | Near-field Dilution in the Bush River..... | 73 |
| 5.2.3 | Mixing Zone Consideration..... | 75 |
| 5.2.3.1 | Mixing Zone Regulations..... | 75 |
| 5.2.3.2 | Site Applicability..... | 78 |
| 5.2.3.3 | Conformance with Aquatic Life Criteria..... | 79 |
| 5.2.3.4 | Re-evaluation of Hazard Assessment Assumptions as an Alternative to a Mixing Zone..... | 80 |

| | | |
|-----|-----------------------------------|----|
| 6. | CONCLUSIONS..... | 81 |
| 6.1 | Biomonitoring Evaluation..... | 81 |
| 6.2 | Hazard Assessment Evaluation..... | 82 |

TABLE OF CONTENTS (CONTINUED)

| | | |
|----|-----------------------|----|
| 7. | LITERATURE CITED..... | 83 |
|----|-----------------------|----|

LIST OF TABLES

| | | |
|----|--|----|
| 1. | SUMMARY OF THE BIOMONITORING TESTS CONDUCTED..... | 25 |
| 2. | SUMMARY OF THE TOXICITY ENDPOINTS/RESPONSES FOR BIOMONITORING TESTS CONDUCTED ON BEACH POINT GROUNDWATER (WELL CC-33B) FROM FEBRUARY 1993 TO DECEMBER 1993..... | 49 |
| 3. | SUMMARY OF THE FOUR BIMONTHLY CHEMICAL ANALYSES (RANGE OF CONCENTRATIONS) CONDUCTED ON RAW BEACH POINT GROUNDWATER (WELL CC-33B) FROM FEBRUARY 1993 TO DECEMBER 1993..... | 57 |
| 4. | LOG OCTANOL WATER PARTITION COEFFICIENTS OF THE CONTAMINANTS DETECTED IN WELL CC-33B..... | 71 |
| 5. | COMPUTED DILUTIONS OF THE GROUNDWATER METALS AND ORGANICS IN THE BUSH RIVER..... | 75 |
| 6. | APPLICABLE TOXIC SUBSTANCES CRITERIA FOR THE BUSH RIVER AND A COMPARISON OF PROJECTED NEAR-FIELD GROUNDWATER CONCENTRATIONS..... | 77 |

LIST OF FIGURES

| | | |
|----|--|----|
| 1. | Microtox® test results of raw (pH \approx 4) groundwater samples taken during the chronic histopathology study..... | 54 |
| 2. | Microtox® test results of buffered (pH \approx 7) groundwater samples taken during the chronic histopathology study..... | 55 |
| 3. | Cumulative mortality of Japanese medaka during the 9-month chronic histopathology exposure study..... | 67 |
| 4. | Percent cumulative mortality of Japanese medaka during the 9-month chronic histopathology exposure study..... | 68 |

LIST OF APPENDICES

1. MICROTOX® ASSAYS CONDUCTED ON RAW (pH ≈4) BEACH
POINT GROUNDWATER (WELL CC-33B), CHRONIC
HISTOPATHOLOGY EXPOSURE TANKS, AND DILUENT WATER..... A1-1
2. MICROTOX® ASSAYS CONDUCTED ON BEACH POINT BUFFERED
(pH ≈7) GROUNDWATER (WELL CC-33B) AND pH OF SAMPLES
USED IN ASSAYS..... A2-1
3. ROTIFER 24-H ACUTE TEST CONDUCTED ON BEACH POINT
RAW (pH ≈4) GROUNDWATER (WELL CC-33B) (TEST NO. 1)..... A3-1
4. ROTIFER 24-H ACUTE TEST CONDUCTED ON BEACH POINT
BUFFERED (pH ≈7) GROUNDWATER (WELL CC-33B)
(TEST NO. 1)..... A4-1
5. ROTIFER 24-H ACUTE TEST CONDUCTED ON BEACH POINT
RAW (pH ≈4) GROUNDWATER (WELL CC-33B) (TEST NO. 2)..... A5-1
6. ROTIFER 24-H ACUTE TEST CONDUCTED ON BEACH POINT
BUFFERED (pH ≈7) GROUNDWATER (WELL CC-33B)
(TEST NO. 2)..... A6-1
7. ROTIFER 24-H ACUTE TEST CONDUCTED ON BEACH POINT
RAW (pH ≈4) GROUNDWATER (WELL CC-33B) (TEST NO. 3)..... A7-1
8. ROTIFER 24-H ACUTE TEST CONDUCTED ON BEACH POINT
BUFFERED (pH ≈7) GROUNDWATER (WELL CC-33B)
(TEST NO. 3)..... A8-1
9. ROTIFER 24-H ACUTE TEST CONDUCTED ON BEACH POINT
RAW (pH ≈4) GROUNDWATER (WELL CC-33B) (TEST NO. 4)..... A9-1
10. ROTIFER 24-H ACUTE TEST CONDUCTED ON BEACH POINT
BUFFERED (pH ≈7) GROUNDWATER (WELL CC-33B)
(TEST NO. 4).....A10-1
11. GREEN ALGAL 96-H GROWTH TEST CONDUCTED ON BEACH
POINT RAW (pH ≈4) GROUNDWATER (WELL CC-33B)
(TEST NO. 1).....A11-1
12. GREEN ALGAL 96-H GROWTH TEST CONDUCTED ON BEACH
POINT BUFFERED (pH ≈7) GROUNDWATER (WELL CC-33B)
(TEST NO. 1).....A12-1
13. GREEN ALGAL 96-H GROWTH TEST CONDUCTED ON BEACH
POINT RAW (pH ≈4) GROUNDWATER (WELL CC-33B)
(TEST NO. 2).....A13-1

LIST OF APPENDICES (CONTINUED)

14. GREEN ALGAL 96-H GROWTH TEST CONDUCTED ON BEACH
POINT BUFFERED (pH \approx 7) GROUNDWATER (WELL CC-33B)
(TEST NO. 2).....A14-1
15. GREEN ALGAL 96-H GROWTH TEST CONDUCTED ON BEACH
POINT RAW (pH \approx 4) GROUNDWATER (WELL CC-33B)
(TEST NO. 3).....A15-1
16. GREEN ALGAL 96-H GROWTH TEST CONDUCTED ON BEACH
POINT BUFFERED (pH \approx 7) GROUNDWATER (WELL CC-33B)
(TEST NO. 3).....A16-1
17. GREEN ALGAL 96-H GROWTH TEST CONDUCTED ON BEACH
POINT RAW (pH \approx 4) GROUNDWATER (WELL CC-33B)
(TEST NO. 4).....A17-1
18. GREEN ALGAL 96-H GROWTH TEST CONDUCTED ON BEACH
POINT BUFFERED (pH \approx 7) GROUNDWATER (WELL CC-33B)
(TEST NO. 4).....A18-1
19. CLADOCERAN ACUTE AND 7-DAY SURVIVAL AND
REPRODUCTION TEST CONDUCTED ON BEACH POINT RAW
(pH \approx 4) GROUNDWATER (WELL CC-33B) (TEST NO. 1).....A19-1
20. CLADOCERAN ACUTE AND 7-DAY SURVIVAL AND
REPRODUCTION TEST CONDUCTED ON BEACH POINT BUFFERED
(pH \approx 7) GROUNDWATER (WELL CC-33B) (TEST NO. 1).....A20-1
21. CLADOCERAN ACUTE AND 7-DAY SURVIVAL AND
REPRODUCTION TEST CONDUCTED ON BEACH POINT RAW
(pH \approx 4) GROUNDWATER (WELL CC-33B) (TEST NO. 2).....A21-1
22. CLADOCERAN ACUTE AND 7-DAY SURVIVAL AND
REPRODUCTION TEST CONDUCTED ON BEACH POINT BUFFERED
(pH \approx 7) GROUNDWATER (WELL CC-33B) (TEST NO. 2).....A22-1
23. CLADOCERAN ACUTE AND 7-DAY SURVIVAL AND
REPRODUCTION TEST CONDUCTED ON BEACH POINT RAW
(pH \approx 4) GROUNDWATER (WELL CC-33B) (TEST NO. 3).....A23-1
24. CLADOCERAN ACUTE AND 7-DAY SURVIVAL AND
REPRODUCTION TEST CONDUCTED ON BEACH POINT BUFFERED
(pH \approx 7) GROUNDWATER (WELL CC-33B) (TEST NO. 3).....A24-1
25. CLADOCERAN ACUTE AND 7-DAY SURVIVAL AND
REPRODUCTION TEST CONDUCTED ON BEACH POINT RAW
(pH \approx 4) GROUNDWATER (WELL CC-33B) (TEST NO. 4).....A25-1

LIST OF APPENDICES (CONTINUED)

26. CLADOCERAN ACUTE AND 7-DAY SURVIVAL AND
REPRODUCTION TEST CONDUCTED ON BEACH POINT BUFFERED
(pH \approx 7) GROUNDWATER (WELL CC-33B) (TEST NO. 4).....A26-1
27. FATHEAD MINNOW ACUTE AND 7-DAY SURVIVAL AND GROWTH
TEST CONDUCTED ON BEACH POINT RAW (pH \approx 4)
GROUNDWATER (WELL CC-33B) (TEST NO. 1).....A27-1
28. FATHEAD MINNOW ACUTE AND 7-DAY SURVIVAL AND GROWTH
TEST CONDUCTED ON BEACH POINT BUFFERED (pH \approx 7)
GROUNDWATER (WELL CC-33B) (TEST NO. 1).....A28-1
29. FATHEAD MINNOW ACUTE AND 7-DAY SURVIVAL AND GROWTH
TEST CONDUCTED ON BEACH POINT RAW (pH \approx 4)
GROUNDWATER (WELL CC-33B) (TEST NO. 2).....A29-1
30. FATHEAD MINNOW ACUTE AND 7-DAY SURVIVAL AND GROWTH
TEST CONDUCTED ON BEACH POINT BUFFERED (pH \approx 7)
GROUNDWATER (WELL CC-33B) (TEST NO. 2).....A30-1
31. FATHEAD MINNOW ACUTE AND 7-DAY SURVIVAL AND GROWTH
TEST CONDUCTED ON BEACH POINT RAW (pH \approx 4)
GROUNDWATER (WELL CC-33B) (TEST NO. 3).....A31-1
32. FATHEAD MINNOW ACUTE AND 7-DAY SURVIVAL AND GROWTH
TEST CONDUCTED ON BEACH POINT BUFFERED (pH \approx 7)
GROUNDWATER (WELL CC-33B) (TEST NO. 3).....A32-1
33. FATHEAD MINNOW ACUTE AND 7-DAY SURVIVAL AND GROWTH
TEST CONDUCTED ON BEACH POINT RAW (pH \approx 4)
GROUNDWATER (WELL CC-33B) (TEST NO. 4).....A33-1
34. FATHEAD MINNOW ACUTE AND 7-DAY SURVIVAL AND GROWTH
TEST CONDUCTED ON BEACH POINT BUFFERED (pH \approx 7)
GROUNDWATER (WELL CC-33B) (TEST NO. 4).....A34-1
35. JAPANESE MEDAKA 96-H ACUTE TEST CONDUCTED ON BEACH
POINT RAW (pH \approx 4) GROUNDWATER (WELL CC-33B).....A35-1
36. LIST OF DATA REPORTS FOR THE AMES ASSAYS CONDUCTED
ON THE BEACH POINT HISTOPATHOLOGY EXPOSURE TANKS
AND APG-EA DILUENT WATER.....A36-1
37. LIST OF DATA REPORTS FOR THE SCE ASSAYS CONDUCTED
ON THE BEACH POINT HISTOPATHOLOGY EXPOSURE TANKS
AND APG-EA DILUENT WATER.....A37-1

LIST OF APPENDICES (CONTINUED)

38. FROG EMBRYO TERATOGENESIS ASSAY - Xenopus (FETAX)
CONDUCTED ON BEACH POINT RAW (pH \approx 4) GROUNDWATER
(WELL 33B) (TEST NO. 1).....A38-1
39. FROG EMBRYO TERATOGENESIS ASSAY - Xenopus (FETAX)
CONDUCTED ON BEACH POINT BUFFERED (pH \approx 7) GROUNDWATER
(WELL 33B) (TEST NO. 1).....A39-1
40. FROG EMBRYO TERATOGENESIS ASSAY - Xenopus (FETAX)
CONDUCTED ON BEACH POINT RAW (pH \approx 4) GROUNDWATER
(WELL 33B) (TEST NO. 2).....A40-1
41. FROG EMBRYO TERATOGENESIS ASSAY - Xenopus (FETAX)
CONDUCTED ON BEACH POINT BUFFERED (pH \approx 7) GROUNDWATER
(WELL 33B) (TEST NO. 2).....A41-1
42. FROG EMBRYO TERATOGENESIS ASSAY - Xenopus (FETAX)
CONDUCTED ON BEACH POINT RAW (pH \approx 4) GROUNDWATER
(WELL 33B) (TEST NO. 3).....A42-1
43. FROG EMBRYO TERATOGENESIS ASSAY - Xenopus (FETAX)
CONDUCTED ON BEACH POINT BUFFERED (pH \approx 7) GROUNDWATER
(WELL 33B) (TEST NO. 3).....A43-1
44. FROG EMBRYO TERATOGENESIS ASSAY - Xenopus (FETAX)
CONDUCTED ON BEACH POINT RAW (pH \approx 4) GROUNDWATER
(WELL 33B) (TEST NO. 4).....A44-1
45. FROG EMBRYO TERATOGENESIS ASSAY - Xenopus (FETAX)
CONDUCTED ON BEACH POINT BUFFERED (pH \approx 7) GROUNDWATER
(WELL 33B) (TEST NO. 4).....A45-1
46. SIX-MONTH INTERIM AND NINE-MONTH FINAL JAPANESE
MEDAKA CHRONIC HISTOPATHOLOGY GROWTH DATAA46-1
47. SUMMARY OF THE SIX-MONTH INTERIM AND NINE-MONTH
FINAL JAPANESE MEDAKA CHRONIC HISTOPATHOLOGY RESULTS...A47-1
48. COMPREHENSIVE CHEMICAL ANALYSES CONDUCTED ON RAW
(pH \approx 4) BEACH POINT GROUNDWATER (WELL CC-33B) AND
CHRONIC HISTOPATHOLOGY EXPOSURE TANKSA48-1
49. ROUTINE WATER QUALITY OF THE CHRONIC HISTOPATHOLOGY
EXPOSURE TANKS.....A49-1
50. ROUTINE WATER QUALITY CHEMICAL METHODS FOR ALL GRAB
SAMPLES TAKEN IN THE BIOMONITORING TRAILER AND ALL
SAMPLES ANALYZED AT THE UMD/WREC LABORATORY.....A50-1

LIST OF APPENDICES (CONTINUED)

51. SCREENING-LEVEL ASSESSMENT OF THE NEAR- AND FAR-
FIELD DILUTION OF THE SURFICIAL GROUNDWATER
DISCHARGE FROM BEACH POINT INTO THE BUSH RIVER.....51-1

SECTION 1

INTRODUCTION

The primary goal of ecotoxicological testing is to predict the effects of chemicals (single elements, compounds, and mixtures) and other stressors (e.g., heated wastewater, suspended solids, etc.) on the long-term health of individual organisms, populations, communities, and ecosystems. There is no perfect short-term (acute) or long-term (chronic) toxicity test which allows one to predict with certainty the effects of a single toxicant (not to mention a complex mixture of contaminants) on ecosystems. Although perfect prediction is not attainable (Barnhouse et al., 1986), a well selected array or battery of biomonitoring assays integrated into a tiered hazard assessment approach can be used to maximize predictability of adverse pollutant effects to aquatic ecosystems (for ex., see Dutka and Kwan, 1988; National Research Council, 1981; Schaeffer and Janardam, 1987).

The selection of an appropriate tier of hazardous assessment biomonitoring tests is based on the problem to be solved (Cairns, 1990; Dickson et al., 1979; Mackay et al., 1989). Single species assays are the most widely used hazard assessment systems (for ex., see Cairns, 1986; Herricks and Schaffer, 1985; Rand and Petrocelli, 1985). Hazard assessment schemes may also incorporate multispecies microcosm tests and ecosystem level simulations in their protocols. Single species toxicity tests, multispecies microcosm tests, and ecosystem level simulations all have advantages and limitations (Neuhold, 1986). The U.S. Army Biomedical Research and Development Laboratory's (USABRDL) Research Methods Branch uses a hazard assessment biomonitoring testing tier which is composed primarily of an array of single species assays.

The University of Maryland in collaboration with USABRDL is evaluating the use of several biomonitoring systems for assessment of potentially contaminated groundwater and sediment-associated contaminants at U.S. Army installations. The current report summarizes an evaluation of contaminated groundwater which contained multiple heavy metals and chlorinated aliphatic hydrocarbons. The contaminated groundwater was withdrawn from the surficial aquifer at Beach Point which is located in the Canal Creek Area of the U.S. Army Aberdeen Proving Ground-Edgewood Area, Aberdeen, MD. The definitive experimental phase of the study was conducted over a 9-month period from February 25, 1993 to December 6, 1993. In addition to the primary goal of the study to evaluate the performance of the biomonitoring systems, the biomonitoring results were also used in a hazard assessment of the groundwater release in the Bush River.

SECTION 2

OBJECTIVES OF STUDY

The primary objectives of the study were to:

- 1) Evaluate the acute toxicity of the groundwater using the 5- and 15-min Microtox® procedure (Photobacterium phosphoreum bioluminescent activity) and the 24-h LC50 Rotifer Toxkit™ (Brachionus rubens) screening test. In addition, the following acute bioassays were also performed: algal (Selenastrum capricornutum) 96-h EC50 growth test, cladoceran (Ceriodaphnia dubia) 48-h LC50, fathead minnow (Pimephales promelas) 96-h LC50, and Japanese medaka Oryzias latipes) 96-h LC50.
- 2) Evaluate chronic toxicity using the 96-h EC50 algal (S. capricornutum) growth test, 7-d cladoceran (C. dubia) survival and reproduction test, and 7-d fathead minnow (P. promelas) survival and growth test.
- 3) Determine the genotoxicity potential of unconcentrated and concentrated samples of the groundwater using the Ames assay and Sister Chromatid Exchange assay.
- 4) Determine the developmental toxicity potential of the groundwater using the frog (X. laevis) embryo teratogenesis assay - Xenopus (FETAX).
- 5) Determine the chronic histopathological potential of the groundwater using a 9-month Japanese medaka (O. latipes) growth and chronic histopathology test.
- 6) Quantify the major chemicals present in the groundwater and monitor the general water quality of the groundwater.

A secondary objective of the study was to use the biomonitoring data obtained in the biomonitoring system evaluations to assess the potential hazard of the groundwater as it moved into an aquatic ecosystem (Bush River).

SECTION 3

BEACH POINT SITE DESCRIPTION

3.1 Physiographic Setting

Beach Point is a peninsula located in the Canal Creek Area of the U.S. Army Aberdeen Proving Ground-Edgewood Area (APG-EA), Aberdeen, MD. The peninsula is bounded by the Bush River on the southeast and Kings Creek on the northwest. The long axis of the peninsula, which is oriented approximately 45° east of north, is ≈ 448 m ($\approx 1,470$ ft) long x ≈ 107 m (≈ 350 ft) wide. The maximum altitude of the land surface is ≈ 4.6 m (≈ 15 ft) above mean sea level (Nemath, 1989); the average elevation is ≈ 3 m (≈ 10 ft) above mean sea level (McGinnis et al., 1994).

3.2 Historical Use

Beach Point has been used for a number of activities which may have contributed to the contamination of the soils and groundwater as well as Kings Creek and Bush River (Nemath, 1989). The major operations include 1) mobile and fixed-based clothing impregnating plant activities during World War II; 2) liquid rocket fuel testing from the early 1960s through the 1970s; and 3) pyrotechnic testing with grenades and pots filled with obscurant smoke from the post-World War II period to about 1970.

As discussed in Nemath (1989), the clothing impregnating process involved the use of several solvents which included 1,1,2,2-tetrachlorethane, N,N'-dichloro-bis(2,4,6-trichlorophenyl)urea, chlorobenzene, and other solvents. Losses of the hazardous materials may have occurred through volatilization, spillage, or leakage, as well as direct discharge to Kings Creek and/or the Bush River. Historical photographs indicate that the materials may have also been discharged to small pits located near the clothing impregnating plants. Various fuels, oxidizers, and fire suppressants were used in the liquid rocket fuel testing programs. Wastewater from the tests was either discharged directly to the Bush River and/or Kings Creek, or may have been allowed to run off onto the ground surface. The primary materials used in the pyrotechnic testing were white obscurant smokes and fog oil. Materials associated with pyrotechnic testing often include aluminum, magnesium, zinc, lead, and titanium, as well as petroleum compounds, hexachloroethane, and other organic compounds (JEG, 1993).

3.3 Groundwater Contamination

Because of possible groundwater contamination from historical activities at Beach Point, USGS began placing observation wells at Beach Point in 1986 as part of a larger

study of the Canal Creek area (Lorah and Vroblesky, 1989). Five wells, located in three clusters, were originally drilled in the surficial aquifer (see Section 3.4) at Beach Point during 1986 in the proximity of the former clothing impregnating plants. A sixth well (CC-33B.1) was drilled in 1988. The three clusters contain the following wells. Cluster 32 consists of wells CC-32A and CC-32B which have screened depths below land surface of 3.2-4.7 and 6.4-7.9 m (10.5-15.5 and 21-26 ft), respectively. Wells 33A, 33B, and 33B.1 in cluster 33 have screened depths of 3.4-4.9, 18.9-20.4, and 12.5-14.0 m (11-16, 62-67, and 41-46 ft), respectively. Well 34A, with a screen interval of 4.3-5.8 m (14-19 ft), is the only well in cluster 34. Of the six monitoring wells at Beach Point in the surficial aquifer, four are shallow with screen depths that range from 3.2-6.9 m (10.5-26 ft), one well (CC-33B.1) is intermediate with a screen depth of 12.5-14.0 m (41-46 ft), and one well (CC-33) is deep with a screen depth of 18.9-20.4 m (62-67 ft).

Chemical monitoring (inorganic and organic constituents) by USGS during 1988 confirmed that hazardous chemicals from prior activities were present in the groundwater (Lorah and Vroblesky, 1989 and Lorah and Clark, 1992). Several inorganic constituents exceeded Federal drinking water standards; the highest concentrations were found in the deep well (CC-33B). Several heavy metals were also found in the wells. The highest concentrations were generally found in wells CC-33B and CC-33B.1; the concentrations of heavy metals in wells CC-33B and CC-33B.1 were similar. A number of volatile chlorinated aliphatic compounds were found in the groundwater at Beach Point. With the exception of chloroform in well CC-32A and methylene chloride in wells CC-33A and CC-33B.1, the highest concentrations of the volatile organics were found in the deep well CC-33B. Wells CC-33A and CC-33B only were studied at Beach Point during 1989. The concentrations of heavy metals and volatile organics were all higher in the deep well (CC-33B) than in the shallow well (CC-33A).

The chlorinated aliphatic organics that occur at the lower depths of the aquifer may be present as a residual denser-than-water nonaqueous phase liquid (DNAPL) (Cohen and Mercer, 1993; Knox et al., 1993). McGinnis et al. (1994) recently found that the surficial aquifer at Beach Point is a relatively low-resistivity lens pierced by high resistivity "plumes" that may indicate the presence of DNAPLs. Additional chemical data are needed from the high resistivity plumes in areas other than well CC-33B to confirm the presence of DNAPLs.

3.4 Geology

The geology of the Canal Creek area has been described by USGS (Oliveros and Vroblesky, 1989). JEG (1993) has recently summarized the geology of the Beach Point study site. Briefly,

the Aberdeen Proving Ground-Edgewood Area is underlain by coastal plain sediments consisting of unconsolidated clay, silt, and sand layers with occasional gravel lenses. At Beach Point the surficial sediments consist of a clayed soil to a depth of ≈ 1.2 m (4 ft), underlain by ≈ 18.2 m (≈ 60 ft) of fine- to medium-grained, well-sorted sand, interfingering with thin lenses of clay and silty sand and sometimes containing coarse sand and gravel layers (Oliveros and Vroblesky, 1989). The surficial unit, which appears to be part of the Potomac Group, is underlain by a clay layer called the upper confining unit. According to JEG (1993), the upper confining unit, which has a thickness of ≈ 27 m (88 ft), does not appear to have been penetrated at Beach Point. Thus, it is unlikely that the Canal Creek Aquifer below the upper confining unit has been contaminated by activities at Beach Point. The Beach Point surficial sediments and clay upper confining unit appear to dip gently (≈ 15 m/mile; 50 ft/mile) to the southeast.

3.5 Groundwater Hydrology

Groundwater at Beach Point is encountered at shallow depths from less than 4 m (13 ft) to about 4.9 m (16 ft) below ground surface in most areas under unconfined conditions (JEG, 1993). The surficial aquifer at Beach Point appears to be an isolated part of the surficial aquifer identified throughout the Canal Creek area by Oliveros and Vroblesky (1989). The surficial aquifer is hydraulically interconnected to Kings Creek, Bush River, and associated wetlands. Limited studies by USGS in 1988 and 1989 showed that the surficial aquifer was tidally influenced (Oliveros and Vroblesky, 1989). Quarterly groundwater elevation monitoring by USGS in wells installed at Beach Point showed that relatively constant water levels (within tidal variations) occurred over the year, with little seasonal variation (Oliveros and Vroblesky, 1989). According to JEG (1993), this indicates that the surficial aquifer at Beach Point is more strongly influenced by tidal conditions than by recharge from infiltrating precipitation.

A recent study of the six monitoring wells described in Section 3.3 above for Beach Point confirmed that the groundwater is tidally influenced (K-V Associates, Inc., 1994). The study showed that all wells exhibited tidal influence. There were varying amounts of tidal influence including both horizontal and vertical components in the groundwater. Wells with shallower screens (CC-32A, CC-33A, and CC-34A) exhibited smaller fluctuations relative to deeper wells with a more permeable stratum (CC-32B, CC-33B, and CC-33B.1). Groundwater flow directions and rates were also substantially influenced by tidal stage. The predominant direction of horizontal flow was to the northeast.

3.6 Surface Water Hydrology

Kings Creek, which drains ≈ 324 ha (≈ 800 acres), is essentially a tidal estuary associated with the Bush River. As discussed by JEG (1993), flow from the creek appears to occur mainly as a result of tidal flushing. Net advective flow resulting from the stream gradient appears to be minimal. Drainage into the main body of the creek is through numerous subsidiary streams and wetlands. The tidal range for the creek is typically <0.3 m (1 ft). Salinity generally varies from approximately 1 to 3 parts per thousand (ppt). The depth of the creek is <3 m (10 ft).

The Bush River, a tributary estuary of the Chesapeake Bay, has a tidal range of 0.15 to 0.46 m (0.5 to 1.5 ft) at Beach Point (JEG, 1993). The river is approximately 1.6 km (0.87 nautical mile) wide at Beach Point and is generally <1.8 m (6 ft) deep except in the shipping channel, where the depth is about 6.1 m (20 ft) (JEG, 1993). The salinity of the river in the vicinity of Beach Point can vary considerably on a seasonal basis; the average is ≈ 3 ppt (Pritchard, 1976).

As discussed by Carter (1976), the Bush River is ≈ 15.7 km (8.5 nautical miles) long measured along its axis from mouth to head. The depth increases generally from head to mouth with the maximum depth of 9.8 m (32 ft) near Briery Point; the mean depth is ≈ 1.8 m (6 ft). At its widest point, the river is ≈ 3.7 km (2 nautical miles) wide. The tide in the Bush River is semidiurnal with a mean range of 0.4 m (1.4 ft) reported for Pond Point. This tidal range results in an intertidal volume of $\approx 1.2 \times 10^7$ m³ (42.9×10^7 ft³). This is about 20% of the volume below mean tide level. At the time of maximum tidal velocity at the entrance, the discharge through the mouth of the river is ≈ 850 m³ sec⁻¹ (30×10^3 ft³ sec⁻¹). The major tributaries to the river include Otter Point Creek, Lauderick Creek, and Kings Creek. The drainage of the Bush River is $\approx 29.5 \times 10^3$ ha (113.9 miles²), resulting in a mean annual runoff of the order of 3.68 m³ sec⁻¹ (130 ft³ sec⁻¹) (Carter, 1976). Thus, waters in the Bush are derived mainly from the adjacent Chesapeake Bay.

SECTION 4

MATERIALS AND METHODS

4.1 Biomonitoring Evaluation

Groundwater was withdrawn from well CC-33B (Hartford County Permit No. HA-81-4049) which is the most highly contaminated well at Beach Point (Section 3.3). The well is 21.3 m deep (70 feet) and has a screened interval of 18.0-20.4 m (62-67 ft). The well pump intake was located at 19.8 m (65 feet). Groundwater was pumped continuously from the well at a rate of ≈ 7.5 L/min (2 gal/min).

Several components of the biomonitoring study were conducted on-site (see below) in an aquatic biomonitoring trailer with bioassay capabilities similar to a USABRDL trailer described by Herriott and Burton (1992). Groundwater and dilution water (APG-EA potable water) were supplied to the trailer via polyethylene pipe. The dilution water was charcoal filtered and aerated before use. Excess groundwater and diluent water from the trailer were collected, treated via charcoal, and pumped to the APG-EA Wastewater Treatment Plant for further treatment.

An array of biomonitoring systems were evaluated during the 9-month study. The biomonitoring systems included a number of endpoints. A summary of the biomonitoring tests conducted is given in Table 1. The pH of the groundwater from well CC-33B was 4 (± 0.1); thus, many of the assays were conducted at both pH 4 and pH 7. The toxicity at pH 7 was studied so that the data could ultimately be used in a hazard assessment of the groundwater as it enters the Bush River which generally has a pH in the low neutral range.

The experimental procedures and frequency of each assay are described in detail below. The following is a brief description of the tier of biomonitoring systems employed in the evaluation. Acute toxicity of the groundwater was evaluated three times each week using the 5- and 15-min Microtox® assay which uses microbial (Photobacterium phosphoreum) bioluminescent activity. In addition to providing rapid toxicity data, the test was also conducted to monitor the toxicity of the groundwater over time. A rapid 24-h rotifer (Branchious rubens) test was conducted to determine acute toxicity. The rotifer bioassay was conducted four times on a bimonthly basis during the course of the study at a pH of 4 and 7. Acute toxicity data were also obtained for the algal, invertebrate, and fish used in the short-term toxicity tests described in Section 4.1.2 below. An acute toxicity test was also conducted with the Japanese medaka (Oryzias latipes).

TABLE 1. SUMMARY OF THE BIOMONITORING TESTS CONDUCTED

| Test and/or Species | Type of Test | Test No. | Test Periods | Comments |
|--------------------------|---|------------------|--|---|
| Microtox® (bacterium) | 5- and 15-min EC50 | N/A | 03/24/93 - 12/06/93 | Grab and/or composite sample ^a |
| Rotifer | 24-h LC50 | 1 2 3 4 | 04/10/93 - 04/12/93 06/02/93 - 06/04/93 08/24/93 - 08/26/93 12/01/93 - 12/03/93 | Grab and composite sample ^b |
| Green alga | 96-h EC50; 96-h Growth | 1 2 3 4 | 04/10/93 - 04/14/93 06/03/93 - 06/07/93 08/24/93 - 08/28/93 12/01/93 - 12/05/93 | Grab and composite sample ^b |
| Cladoceran | 48-h LC50; 7-d Survival and reproduction | 1 2 3 4 | 04/07/93 - 04/14/93 06/01/93 - 06/08/93 08/24/93 - 08/31/93 11/30/93 - 12/07/93 | Grab and composite sample ^c |
| Fathead minnow | 96-h LC50; 7-d Survival and growth | 1 2 3 4 | 04/07/93 - 04/14/93 06/02/93 - 06/09/93 08/24/93 - 08/31/93 11/30/93 - 12/07/93 | Grab and composite sample ^c |
| Japanese medaka | 96-h LC50 | N/A | 11/30/93 - 12/04/93 | Grab and composite sample ^c |
| Genotoxicity (bacterium) | Ames assay | 1 2 3 4 | 04/14/93 06/02/93 08/23/93 11/30/93 | Grab and/or composite sample ^{d,e} |

TABLE 1. (CONTINUED)

| Test and/or Species | Type of Test | Test No. | Test Periods | Comments |
|---|----------------|--------------------------|--|--|
| Genotoxicity (Chinese hamster ovary cells) | SCE assay | N/A N/A N/A N/A | 06/02/93 07/19/93 08/23/93 11/30/93 | Grab and/or composite sample ^f |
| Developmental toxicity (African clawed frog) | 4-d FETAX | 1 2 3 4 | 04/10/93 - 04/14/93 06/03/93 - 06/07/93 08/26/93 - 08/30/93 12/03/93 - 12/07/93 | Grab and composite sample ^e |
| Chronic histopathology and growth (Japanese medaka) | 6 and 9 Months | N/A | 02/25/93 - 11/29/93 | Flow-through exposure conducted in biomonitoring lab |
| Comprehensive chemical analyses | N/A | 1 2 3 4 | 04/14/93 06/02/93 08/23/93 11/30/93 | Grab and composite sample ^g |
| Munitions analyses | N/A | 1 2 3 4 | 04/14/93 06/02/93 09/30/93 11/29/93 | Grab sample of 100% groundwater |
| Routine water quality analyses | N/A | N/A | Daily/weekly | Grab sample ^h |

^a Grab sample of 100% groundwater and composite grab samples of 10 and 1% groundwater by volume taken 3 times a week; composite grab sample of diluent water taken once a week.

TABLE 1. (CONTINUED)

| Test and/or Species | Type of Test | Test No. | Test Periods | Comments |
|---------------------|---|----------|--------------|----------|
| ^b | Grab sample of 100% groundwater; composite grab sample of diluent water. | | | |
| ^c | Grab sample of 100% groundwater and composite grab sample of diluent water renewed every 24 h. | | | |
| ^d | Composite grab samples of 10 and 1% groundwater by volume and diluent water taken for analysis on 04/14/93 and 06/02/93. | | | |
| ^e | Grab sample of 100% groundwater and composite grab samples of 10% groundwater by volume and diluent water taken for analysis on 08/23/93 and 11/30/93. | | | |
| ^f | Samples taken for analyses were as follows: composite grab sample of 10% groundwater by volume on 06/02/93, grab sample of 100% groundwater on 07/19/93 and 11/30/93, and composite grab sample of diluent water on 08/23/93. | | | |
| ^g | Grab sample of 100% groundwater and composite grab samples of 10 and 1% groundwater by volume and diluent water taken for analysis during each test period listed. | | | |
| ^h | Grab samples taken from chronic histopathology tanks as described in Section 4.1.6.2. | | | |

The following 4- to 7-d short-term toxicity tests, which were used to estimate chronic toxicity, were performed on a bimonthly basis: 96-h algal (Selenastrum capricornutum) growth test; 7-d cladoceran (Ceriodaphnia dubia) survival and growth test; and 7-d fathead minnow (Pimephales promelas) survival and growth test. Four bimonthly tests were conducted with each species at pH 4 and pH 7. In addition to the short-term methods used to estimate chronic toxicity, growth data at 6 and 9 months from the chronic Japanese medaka histopathology test described below were also used as chronic toxicity endpoints.

Gene mutation potential was determined using the Ames Salmonella/mammalian-microsome reverse assay. Primary DNA damage was evaluated by the sister chromatid exchange (SCE) assay. Developmental toxicity was determined by the 96-h frog embryo teratogenesis assay-Xenopus (FETAX) using the African clawed frog, Xenopus laevis. Genotoxicity and developmental toxicity assays were conducted at bimonthly intervals during the same periods as the above acute and short-term chronic tests were conducted. Chronic histopathological changes were evaluated using the Japanese medaka as the experimental model. Both unexposed and fry exposed to diethylnitrosamine (DEN) were exposed continuously under flow-through test conditions for 6- and 9-month exposure periods.

Comprehensive chemical analyses of the raw groundwater, test dilutions in the chronic histopathology assay, and diluent water were performed four times at bimonthly intervals. The chemical analyses were conducted during the same periods that the above bimonthly assays were conducted. Routine water quality analyses were also conducted at various frequencies on a weekly basis.

4.1.1 Acute Toxicity Tests

4.1.1.1 Microtox®

The Microtox® test (Microbics Corp., Carlsbad, CA) is a rapid acute toxicity test that may be completed in less than one hour. The test is based on the reduction in bioluminescence of the marine bacterium P. phosphoreum when exposed to a sample of unknown toxicity. The degree of light reduction, an indication of metabolic inhibition in the test preparation, indicates the degree of toxicity of the sample. The Microtox® test procedures followed were those outlined in Herriott and Burton (1992) which were derived from Microtox®'s operating manual (Microtox®, 1988). A Microtox® Model 500 Analyzer with PC version 6.3 software was used for both a 5-min and 15-min test on all samples.

Microtox® assays were initiated on April 2, 1993. The assays were conducted on-site three times a week until the biomonitoring study was completed on December 6, 1993. Both 5- and 15-min assays were conducted on 100, 10, and 1% groundwater

by volume samples as well as APG-EA diluent water. The 10 and 1% groundwater by volume samples were taken from the chronic histopathology study tanks as described below in Section 4.1.5. A 5- and 15-min assay was conducted once each week on 100% groundwater adjusted to pH 7 with 10 N NaOH.

4.1.1.2 Rotifer

The 24-h freshwater rotifer Toxkit™ screening test (Bio-Response Systems, Halifax, Nova Scotia) was used to determine the potential toxicity of the groundwater at pH 4 and 7 (groundwater buffered with 10 N NaOH). pH-adjusted aliquots of the same groundwater used for the short-term chronic toxicity tests were used for the rotifer assays (see Section 4.1.2). The test utilized newly hatched rotifers (*B. rubens*) <4 h old. The rotifers used in the tests were hatched from cysts supplied in the Rotifer ToxKit™. Rotifer ToxKit™ synthetic medium was used to hatch the cysts, rear the organisms before testing, and as diluent media for the tests. The static tests were conducted in glass Petri dishes containing 10 mL of test solution.

The groundwater, which was used within 6 h from the time of collection for the rotifer bioassays, was held in glass containers at 4°C until used in the tests. A geometric series of five groundwater concentrations (plus controls) was used. Three replicates of 10 organisms each were used at each test concentration. All tests were conducted at $25 \pm 0.5^\circ\text{C}$. Routine water quality (alkalinity, conductivity, DO, hardness, pH, and temperature) was measured at the beginning and end of each test. The methods used for the chemical analyses are given in Section 4.1.6.2. All tests were conducted under a 16-h light:8-h dark photoperiod (fluorescent lights at 60-85 foot candles).

4.1.1.3 Green Alga, Cladoceran, Fathead Minnow, and Japanese Medaka

Acute toxicity values were calculated where possible at pH 4 and 7 for the green alga, cladoceran, and fathead minnow from the data obtained during the short-term chronic tests described in Section 4.1.2. With regard to the green alga, EPA's Office of Research and Development considers the 96-h algal test for growth to be a short-term chronic test for determining the toxicity of effluents (Horning and Weber, 1985; Weber et al., 1989) as do other investigators for evaluating single chemicals (for ex., see Hughes et al., 1988 and Suter, 1993). EPA's Toxic Substance Control Act office considers the 96-h test to be an acute test (U.S. EPA, 1985a and 1986a). Because we used the short-term chronic method (Section 4.1.2.1), we analyzed the data as chronic data; however, we also analyzed and reported the results as 96-h acute data so that acute:chronic ratios could be calculated for use in the hazard assessment (Section 4.2).

Forty-eight-h LC50s and 96-h LC50s were determined where possible for the cladoceran and fathead minnow, respectively. One 96-h acute toxicity test was conducted with the Japanese medaka using the procedure of Weber (1991). The age of the Japanese medaka used in the test was 22 d old at the start of the exposure; the test procedure recommends fish between the ages of 1-14 d old be used.

4.1.2 Short-term Chronic Toxicity Tests

The specific test methods for the short-term chronic tests are given below. Deviations from the test methods are discussed where appropriate. A geometric series of five groundwater concentrations (plus controls) was used in most tests. The groundwater samples used in all tests were obtained daily and used within 6 h at each 24-h renewal (see below). All groundwater samples were transported in glass containers on ice and held at 4°C until used for the tests. Each groundwater sample was split into two aliquots. One aliquot was maintained at pH 4 and the second buffered to pH 7 (10 N NaOH). All bimonthly short-term chronic bioassays were conducted at pH 4 and 7. The same pH-adjusted aliquots were also used for the rotifer bioassays (Section 4.1.1.2) and FETAX assays (4.1.4). The bioassays were conducted at the University of Maryland Wye Research and Education Center (UMD/WREC) Aquatic Toxicology Laboratory.

4.1.2.1 Green Alga

The short-term chronic toxicity of the groundwater to the green alga (S. capricornutum) at pH 4 and 7 (10 N NaOH) was determined four times by the EPA procedures given in Weber et al. (1989). A starter culture of S. capricornutum was obtained from the culture collection at the University of North Texas, Denton, TX. Stock algal cultures were reared in 2.5 L Pyrex culture flasks containing 1 L of sterilized double strength "AAP" algal assay medium, with sufficient P added to achieve a 20:1 N:P ratio as described in Miller et al. (1978). Cultures were maintained in a constant temperature incubator under constant cool-white fluorescent lights (≈ 300 foot candles) at a temperature of $25 \pm 0.2^\circ\text{C}$ on a shaker table oscillating at 100 rpm ($\pm 10\%$). Log growth cells were used to start all tests.

Algal test solutions were prepared by dilution of the groundwater with filtered sterilized assay media. Test solutions (100 mL total volume) were dispensed into 250 mL Delong flasks and inoculated with S. capricornutum cells in log growth to achieve a density of $\approx 1 \times 10^4$ cells/mL. Triplicates were prepared for each treatment. The flasks were placed on a shaker table in an incubator set at the culturing conditions described above. Growth measurements (cell density) were made from all replicates in each treatment at 0, 24, 48, 72, and 96 h. Algal

cell density was determined from a 1 mL sample with a Model ZBI Coulter Counter (Coulter Electronics, Inc., Hialeah, FL). The instrument was calibrated with each use via hemocytometer counts. Test solutions were not renewed during the 96-h studies.

4.1.2.2 Cladoceran

The chronic toxicity of the groundwater at pH 4 and 7 to C. dubia was determined four times by the EPA static renewal method (solutions renewed daily) given in Weber et al. (1989). The cladoceran was cultured at $25 \pm 1^\circ\text{C}$ in 600 mL glass beakers filled with 400 mL of 20% Perrier:80% reverse osmosis water amended with selenium ($2 \mu\text{g Se/L}$ as Na_2SeO_3) as recommended by Winner (1989). The diet consisted of a mixture of Cerophyl® (Cerophyl Laboratories, Inc., Kansas City, MO) and the green alga, S. capricornutum, added to the cladoceran culture to achieve final concentrations of $120 \mu\text{g Cerophyl}^\circ/\text{mL}$ and 6.7×10^5 S. capricornutum cells/mL. Starter cultures of C. dubia were obtained from the Center for Lake Superior Environmental Studies, University of Wisconsin - Superior.

All neonates used in the 7-d survival and reproduction tests were produced by cladocerans in culture that had released at least three broods. The initial age of the neonates in each test was <12 h old. The tests were conducted in 50 mL glass beakers containing 30 mL of test solution. All tests were conducted in an environmental chamber at $25 \pm 1^\circ\text{C}$ under a 16-h light:8-h dark photoperiod (fluorescent lights; 60-85 foot candles at the surface of the culture vessels). All test organisms were fed daily as described above at each 24-h renewal. Routine water quality was taken at the beginning and end of each 24-h renewal. The methods used for the chemical analyses are discussed in Section 4.1.6.2.

4.1.2.3 Fathead Minnow

The toxicity of the groundwater at pH 4 and 7 to fathead minnows (P. promelas) was determined four times by the EPA static renewal method (solutions renewed daily) given in Weber et al. (1989). All larvae used in the 7-d survival and growth tests were <24 h old at the start of the test. The tests were conducted in 600 mL glass beakers containing 400 mL of test solution. The dilution water was a 20% Perrier:80% reverse osmosis water. All test organisms were fed brine shrimp (Artemia sp.) nauplii <24 h old daily at each 24-h renewal. All tests were conducted at $25 \pm 1^\circ\text{C}$ under a 16-h light:8-h dark photoperiod (fluorescent lights; 60-85 foot candles). Routine water chemistry was performed at the beginning and end of each renewal. Dry weight was determined by drying at 100°C for a minimum of 12 h.

Fathead minnow larvae were obtained from the UMD/WREC culture maintained at $25 \pm 1^\circ\text{C}$ in UMD/WREC non-chlorinated well water (mean dissolved oxygen = 8.2; pH = 7.8; conductivity = 161 $\mu\text{S}/\text{cm}$; alkalinity = 53 mg/L as CaCO_3 ; hardness = 52 mg/L as CaCO_3). The UMD/WREC culture procedures were similar to those recommended by Peltier and Weber (1985). The UMD/WREC culture was initiated with mature fathead minnows obtained from the U.S. EPA Environmental Monitoring and Support Laboratory - Cincinnati, Ohio.

Spawning fish were cultured in fiberglass tanks (2.4 x 0.8 x 0.5 m) containing 0.2 m UMD/WREC well water held at $25 \pm 1^\circ\text{C}$. The spawning adults were fed a diet of frozen brine shrimp (*Artemia* sp.; Argent Chem. Lab., Redmond, WA) and TetraMin® Staple Food (Ramfab Aquarium Products Co., Oak Ridge, TN) twice daily. Excess food was removed daily. Four sets of spawning fathead minnows were maintained in the culture tanks at a ratio of 1 male:4 females. Replacement spawners were rotated at approximately 3-month intervals. Fathead minnow embryos were collected on spawning substrates (10 cm I.D. x 20 cm long PVC pipe sections cut longitudinally in equal portions) and transferred to 19 L aquaria at $25 \pm 1^\circ\text{C}$ in UMD/WREC well water for hatching. All stages of the fish were reared under a 16-h light:8-h dark photoperiod (fluorescent lights; 60-85 foot candles).

4.1.3 Genotoxicity Tests

4.1.3.1 Gene Mutation Assay

The Ames assay was used to predict chemical mutagenic activity which in turn may serve as a carcinogen prescreen test (Ames et al., 1973). *Salmonella typhimurium*/mammalian-microsome reverse mutation assays were conducted four times on the groundwater and APG diluent water samples described below. The assays were conducted on both unconcentrated and concentrated (10X via XAD-2 resin extracts) samples of the groundwater and diluent water. The Ames mutagenicity assays were conducted by Hazleton Washington, Inc., Vienna, VA.

Grab samples of groundwater were collected in polypropylene containers as follows. Raw groundwater was taken directly from the well fed line to the biomonitoring laboratory. Groundwater samples of 10 and 1% groundwater by volume were taken as composite samples from the chronic histopathology fish tanks described in Section 4.1.5. Grab samples of APG-EA diluent water were taken from a large polypropylene tank with a 99% particle replacement time of ≈ 12 h. Thirty-one liters (1 L for an unconcentrated sample and 30 L for a 10X sample) of each material were siphoned into Nalgene polycarbonate carboys, packed in ice, and transported to Hazleton Washington, Inc. in insulated containers. The unconcentrated samples were analyzed by Hazleton

Washington, Inc. Protocol No. HLA Protocol 401W, Edition 17. The concentrated (10X) samples were analyzed by Protocol No. HLA Protocol 401X, Edition 18.

Detailed experimental procedures for the unconcentrated and 10X tests are given in the protocols listed above. Briefly, the mutagenicity assays evaluated the groundwater and diluent water samples for their ability to induce reverse mutations at the histidine locus in the genome of specific S. typhimurium tester strains both in the presence and absence of an exogenous metabolic activation system of mammalian microsomal enzymes derived from Aroclor 1254-induced rat liver. The tester strains used in the assays were TA98 and TA100. The assays were conducted using two plates per dose level in the presence of microsomal enzymes. Six dose levels of the groundwater and diluent water samples were tested in both the presence and absence of S9 along with appropriate vehicle controls (three plates per dose), negative controls, and positive controls. Resin controls were also run for the 10X samples. The doses tested in the 10X assays varied based on the amount of extractable organics recovered from the test material.

4.1.3.2 Sister Chromatid Exchange Assay

The sister chromatid exchange (SCE) assay was used as a test for possible genetic damage induced by chemical mutagens (Perry and Evans, 1975). We are aware that many mutagens are active in SCE assays, yet, it is still incompletely understood how DNA damage or perturbations of DNA synthesis give rise to SCE (Hoffman, 1991). The assay was performed once on concentrated samples ($\approx 50,000X$) of 100% raw groundwater, 10% groundwater by volume, and APG-EA diluent water. One assay was also conducted on unconcentrated 100% groundwater. The SCE assays were conducted by Hazleton Washington, Inc., Vienna, VA.

Grab samples of groundwater were collected in polypropylene containers as follows. The 100% groundwater samples were taken directly from the well fed line to the biomonitoring laboratory. The 10% groundwater by volume sample was taken as a composite sample from the chronic histopathology fish tanks described below. A grab sample of APG-EA diluent water was taken from a large polypropylene tank with a 99% particle replacement time of ≈ 12 h. Thirty liters were siphoned into Nalgene polycarbonate carboys, packed in ice, and transported to Hazleton Washington, Inc. in insulated containers. The samples were assayed by Hazleton Washington, Inc. using their Genetics Protocol No. 438, Edition 15.

The experimental procedures for the SCE assay are given in the protocol listed above. Briefly, 30 L of each sample were concentrated in DMSO to a final volume of 6 mL ($\approx 50,000X$) by rotoevaporation. The unconcentrated 100% groundwater sample was

not concentrated by rotoevaporation. The extract was evaluated for its ability to induce sister chromatid exchanges (determined at metaphase) in Chinese hamster ovary (CHO) cells in the presence and absence of an exogenous metabolic activation system of mammalian microsomal enzymes derived from Aroclor 1254-induced rat liver. The SCE assays were conducted by exposing cultured cells to the test extract, growing the cells with the thymidine analog of 5-bromo-2'-deoxyuridine for about two cell cycles, and making chromosome preparations that were Giemsa-stained for SCE. The doses tested were selected based on the amount of extractable organics recovered from the test article. Single cultures of CHO cells were incubated with five concentrations of the extract with and without metabolic activation. Sister chromatid exchange frequencies were analyzed in the cultures treated with the two highest doses with second generation cells and from solvent and positive control cultures.

4.1.4 Developmental Toxicity Test

Four bimonthly developmental toxicity tests were conducted at pH 4 and 7 using the frog embryo teratogenesis assay - Xenopus (FETAX). The assay is a 96-h quantitative developmental assay used to screen for developmental toxicants in aquatic media. The assays were conducted using the static renewal (solutions renewed every 24 h) test method Designation E 1439-91 of the American Society for Testing and Materials (ASTM, 1992a). Embryo lethality and malformations were determined; growth retardation was not evaluated. The identification and interpretation of malformations in the embryos at 96 h were made via the atlas of Bantle et al. (1991). Aliquots of the same groundwater used for the rotifer, algal, and short-term chronic toxicity biomonitoring tests were used for the FETAX assays (Section 4.1.2).

Embryos between normal stage 8 blastulae and normal stage 11 gastrulae were obtained from X. laevis breeding colonies at the UMD/WREC as described below. The embryos were de-jellied in a 2% L-cysteine solution (2 g of L-cysteine per 98 mL of FETAX solution). Once de-jellied, the embryos were rinsed and re-suspended in FETAX solution (ASTM, 1992a). The embryos were tested in glass petri dishes containing 10 mL of solution. Two replicates of 25 embryos/replicate were used for each test treatment. As required by the study protocol, four replicates of 25 embryos/replicate were used for the control treatment. The tests were conducted at $24 \pm 0.2^\circ\text{C}$ under a 16-h light: 8-h dark photoperiod (fluorescent lights; ≈ 75 foot candles at the surface of the test medium) in a constant temperature environmental chamber.

The UMD/WREC X. laevis adult colony was maintained in flow-through (≈ 4 replacement volumes per day) circular polyethylene aquaria (0.91 m I.D. x 0.36 m high) with a water depth of 10 cm. Each aquarium contained a maximum of 10 adults. UMD/WREC non-

chlorinated deep well water (water quality given in Section 4.1.2.3) held at 23.5 ± 0.5 °C served as the culture medium. All frogs were fed every 5-6 d with commercial beef liver supplemented with liquid vitamins (PolyViSol; Mead-Johnson Nutritionals, Evansville, IN). The colony was held under a photoperiod of 16h light:8 h dark. Mating pairs were bred in the dark in 23.5 ± 0.5 °C UMD/WREC non-chlorinated water at ≈ 70 d intervals by injecting 400 and 800 I.U. of human chorionic gonadotropin (HCG) in the dorsal lymph sac of the males and females, respectively. Amplexus occurred 4-6 h after injecting HCG; egg deposition occurred 9-12 hours following HCG injection. The original breeding stock was obtained from Xenopus I (Ann Arbor, MI).

4.1.5 Chronic Histopathology and Growth Test

Chronic histopathologic changes were evaluated using the Japanese medaka (*O. latipes*) as the experimental model. The Japanese medaka is a sensitive laboratory model for screening environmental pollutants which may induce histopathological changes and neoplasms (for ex., see Hawkins et al., 1988; Klaunig et al., 1984; Metcalfe, 1989). Both unexposed and fry exposed to diethylnitrosamine (DEN) were exposed continually under flow-through test conditions for a 9-month period. A subset of organisms was taken after 6 months of exposure for morphometric measurements and histopathological evaluation. The USABRDL test designation was Protocol No. 401-001R.

The fish were exposed to one of three treatments: 10% groundwater by volume, 1% groundwater by volume, or APG-EA diluent water (control). A 100% groundwater treatment could not be used because the pH of the groundwater was 4 (± 0.1) which would have caused excessive mortality over the 9-month exposure period. Consideration was given to buffering the 100% groundwater to pH 7 and conducting the 100% treatment at pH 7. However, a preliminary evaluation of the buffered groundwater at pH 7 showed that excessive precipitation of metals occurred. Thus, 100% groundwater was not buffered to pH 7 and used as an experimental treatment.

The flow-through test solutions were delivered by a solenoid-activated proportional dilutor system which was constructed primarily of glass and stainless steel; some silicon tubing was also used. The test concentrations were delivered to twelve 19 L (5 gal) glass aquaria (4 aquaria at 10% groundwater by volume; 4 at 1% groundwater by volume; and 4 control aquaria); each aquarium contained a volume of ≈ 16 L (4.25 gal). All aquaria were held at 25 ± 1 °C in a constant temperature water bath. The dilutor was calibrated to complete one full cycle every 2.5 to 3.5 min. During a cycle, tanks 1-4 each received 300 ± 15 mL of APG-EA diluent water, tanks 5-8 each received 300 ± 15 mL of 10% groundwater by volume, and tanks 9-12 each

received 300 ± 15 mL of 1% groundwater by volume.

Both unexposed fry and fry exposed to DEN, were reared off-site at USABRDL until 17 d old. The DEN-initiated fish were exposed to 10 mg/L DEN for 48 h when the organisms were 12 to 14-d old. Prior to the start of the exposure to groundwater, the 17-d old fish were randomized into 6 groups of 60 fish/group for both the unexposed and DEN-initiated groups. The fish were suspended in 12 1-L mesh-bottom glass beakers in the appropriate flow-through test aquaria in the biomonitoring laboratory. The fish were held in the beakers for one week after which they were released into the aquaria.

Japanese medaka, 17-22 d old, were fed microworms two feedings per day and live brine shrimp (*Artemia* sp.) (<24 h old) two feedings per day (30 brine shrimp/fish). Pre-adult fish, 23-30 d old, were fed Tetramin® flake food two feedings per day and live brine shrimp <24 h old (one feeding per day; 40 brine shrimp per fish). Adult fish, >30 d old, were fed Tetramin® flake food (three feedings per day on Tuesday and Thursday and two feedings per day on the remaining days) and live brine shrimp (one feeding per day on Monday, Wednesday, and Friday). The ration was adjusted as the size of the fish increased. Tanks were cleaned on an as needed basis (usually 1-2 times a week) by scrubbing algae from the sides of the tanks, allowing the debris to settle, and then siphoning. Tetramin® was fed ad libitum for 15-30 min during each feeding.

The number of test organisms alive in each tank were monitored and recorded daily. Moribund fish were euthanized and fixed in Bouin's solution for subsequent histological observation. The dilutor cycle time was calculated and recorded daily. The volume of groundwater and diluent water delivered to the aquaria were checked weekly. When necessary, cycle time and/or volume distributions were adjusted. The dilutor was occasionally shutdown (for no more than one hour) and cleaned on an as needed basis. Daily water quality (DO, pH, and temperature) was determined in all aquaria. Additional water quality tests (alkalinity, hardness, conductivity, total residual chlorine, free available chlorine, and total ammonia-nitrogen) were performed once a week in all aquaria (Section 4.1.6.2). A 16-h light:8-h dark photoperiod (fluorescent lights at 70-100 foot candles) was maintained throughout the study. Unionized ammonia-nitrogen was determined by the method of Thurston et al. (1979). Comprehensive chemical analyses were performed four times at bimonthly intervals as discussed in Section 4.1.6.1 on 100, 10, and 1% groundwater by volume and APG-EA diluent water during the test periods shown in Table 1.

On day 180, 20 Japanese medaka in each tank were removed and taken back to USABRDL for fixation (Bouin's solution) and subsequent histological observation. Wet weight and standard

length measurements were taken on all fish. The morphometric data were taken to assess the effects of a chronic 6-month exposure to the contaminated groundwater as well as the general health of the fish. On day 278, when the exposure was completed, the remaining Japanese medaka were also taken back to USABRDL for morphometric measurements and subsequent histological analysis. The histological analyses for the 6- and 9-month exposures were performed by Experimental Pathology Laboratories, Inc., Herndon, VA.

4.1.6 Chemical Analyses

4.1.6.1 Comprehensive Chemical Analyses

Comprehensive chemical analyses were performed four times at bimonthly intervals on 100, 10, and 1% groundwater by volume and APG-EA diluent water during the test periods shown in Table 1. The comprehensive chemical analyses included general water quality, metals, volatile organics, acid compounds, base/neutral compounds, pesticides, herbicides, organophosphorus pesticides, and munitions. The elements and/or compounds analyzed in each group are presented in the data tables discussed in Section 5.1.6.1. The 100% groundwater samples were grab samples taken directly from the well fed to the biomonitoring trailer. The 10 and 1% groundwater by volume samples were composite samples taken from the four 10 and 1% groundwater by volume replicate treatment tanks in the chronic histopathology study (Section 4.1.5). Grab samples of APG-EA diluent water were taken from a large polypropylene tank with a 99% particle replacement time of ≈ 12 h.

The groundwater and APG-EA diluent water samples were placed in appropriate containers for various analyses. The containers were placed on ice and delivered to a vendor for the analyses. The first comprehensive analysis was performed by Biospherics Inc., Laurel, MD, for all materials with the exception of the munitions. The remainder of the analyses were performed by Gascoyne Inc., Baltimore, MD, for all materials with the exception of the munitions. The methods used for the analyses of all materials are given in the data tables discussed in Section 5.1.6.1. The four munitions samples were analyzed by USABRDL via in-house procedures (U.S. Biomedical Research and Development Laboratory, 1993).

4.1.6.2 Routine Water Quality Analyses

Routine water quality was measured in all histopathology treatment tanks. Dissolved oxygen, pH, and temperature were measured daily. Alkalinity, hardness, conductivity, total residual chlorine, free available chlorine, and total ammonia-nitrogen were measured once a week (all tests were performed on the same days). Unionized ammonia-nitrogen was determined by the method of Thurston et al. (1979). The methods used for the

analyses are presented in Section 5.1.6.2. The same methods used in the biomonitoring trailer were also used for the toxicity studies conducted at the University of Maryland Wye Research and Education Center.

In addition to the temperature measurements made in the aquaria during the chronic histopathology test, temperature was monitored continuously in Tank No. 3 via a strip chart recorder (Cole-Parmer Thermistor Recorder Model No. 08354-15, Cole-Palmer Instrument Co., Chicago, IL).

4.1.7 Test Endpoints and Data Analyses

The test endpoint for the Microtox® 5- and 15-min EC50s was a reduction in bioluminescence. The EC50s and their 95% fiducial limits were determined by probit analysis using the software program supplied by Microtox® (Microtox®, 1988). The test endpoint for the acute effects of groundwater to the green alga was growth, measured as density (cells/mL). The 96-h EC50s for growth were estimated by using the "inhibition proportion" technique recommended by Horning and Weber (1985). The technique uses quantal analyses (e.g., probit or moving average angle methods) to estimate EC50s and their 95% fiducial or confidence limits. Since the assumptions of the quantal analysis are not met in the classical sense because of the very nature of the growth data, the count data at each treatment were averaged and subsequently converted to "inhibition proportions" using the formula below before a moving average angle analysis was performed (Stephan, 1978).

$$I = C - T / C * 100$$

where: C = the mean growth of the controls
T = the mean growth at a given treatment

The 96-h EC50s and their 95% confidence limits for embryo malformations in the FETAX assays were determined by the moving average angle method using an EPA statistical program (Stephan, 1978). The test endpoint for all 24-h LC50 toxicity tests with rotifers, 48-h and 7-d LC50 tests with cladocerans, and 96-h and 7-d LC50 tests with fathead minnows was mortality. The LC50s and their 95% confidence limits were determined by the moving average angle method (Stephan, 1978).

The test endpoint for the chronic toxicity of groundwater at pH 4 and 7 to the green alga was growth measured as density (cells/mL). The no-observed-effect concentrations (NOEC) and lowest-observed-effect concentrations (LOEC) were determined by Dunnett's test. Dunnett's test consists of an analysis of variance (ANOVA) to determine the error term, which is then used in a multiple comparison test for comparing each of the treatment means with the control mean. The assumptions upon which the use

of Dunnett's test are contingent are that the observations within treatments are independent and normally distributed, with homogeneity of variance. The chi-square test for normality and Bartlett's test for homogeneity of variances were performed before the Dunnett's test was used. The above statistical tests were performed using Toxstat (Gulley et al., 1989) at a minimum probability level of 0.05.

The endpoints for the 7-d survival and reproduction tests with Ceriodaphnia were survival and young production. The endpoints for the fathead minnow 7-d survival and growth tests were survival and growth. The endpoints for the 96-h FETAX assay were survival and number of malformations. The statistics used for the LC50 data and FETAX EC50 (malformations) data are given above. NOECs and LOECs were determined as follows. The adult raw cladoceran survival data were analyzed by Fisher's Exact test. Arc-sine square root transformations were made on the FETAX percent embryo survival and percent embryo malformation data as well as the fathead minnow percent survival raw data before further data analyses were performed. With the exception of the cladoceran survival data, all data were then subjected to a chi-square test of normality and Bartlett's test for homogeneity of variance.

When the data sets met the assumptions of normality and homogeneity of variance, a parametric statistic was used. Dunnett's test was used when the number of replicates was constant among treatments. A t-test with Bonferroni adjustment of error rate was performed when the number of replicates was not constant among treatments. When a data set failed to meet the assumptions of normality or homogeneity of variance, a nonparametric statistic was used. Steel's Many-One Rank test was performed when equal number of replicates were used. The statistical tests were performed using Toxstat (Gulley et al., 1989). A minimum probability level of 0.05 was used for all tests.

The morphometric endpoints for the Japanese medaka chronic histopathology study after 6 and 9 months of exposure were wet weight and standard length. The raw data were first checked for normality and homogeneity of variance by the Kolmogorov-Smirnov test and Levene median test, respectively. The wet weight data after 6 and 9 months of exposure passed the tests for normality and homogeneity of variance and were subsequently analyzed by a general linear model type III test (replicates within treatment of the error term). The standard length data after 6 and 9 months of exposure were not normally distributed; thus, the nonparametric Kruskal Wallis one way ANOVA on ranks statistic was used. A statistically significant difference ($\alpha = 0.05$) was found for length after 9 months of exposure using the nonparametric Kruskal Wallis test. Thus, Dunn's multiple comparison test was used to enumerate the difference between the

control and each experimental treatment. All statistical tests with the exception of the general linear model were performed via Sigma Stat (1992). The general linear model was run with SAS (1987). A minimum probability level of 0.05 was used for all tests. The histopathological data enumerated by Experimental Pathology Laboratory, Inc. at 6 and 9 months were not treated statistically.

4.2 Hazard Assessment Evaluation

The U.S. Army Aberdeen Proving Ground Directorate of Safety, Health, and Environment (DSHE) is conducting a risk assessment of the Canal Creek area as part of their Installation/Restoration Program. The potential hazard of the contaminated surficial aquifer at Beach Point, which appears to be an isolated part of the surficial aquifer identified throughout the Canal Creek area (Section 3.5), is being addressed by DSHE as an operable unit in a focus feasibility study of Beach Point. A hazard assessment of the contaminated surficial aquifer to aquatic organisms was performed in this study using the toxicity data obtained in the biomonitoring evaluation study. The ASTM Designation E 1023-84 (Reapproved 1988) standard guide for assessing the hazard of a material to aquatic organisms was used as the basis for the hazard assessment evaluation (ASTM, 1992b).

4.2.1 Scope of the Hazard Assessment

The hazard assessment process is complex and requires decisions at a number of points (ASTM, 1992b). The validity of a hazard assessment depends on the soundness of the decisions and the accuracy of information used. All decisions should be based on reasonable worst-case analyses so that an appropriate assessment can be completed for the least cost that is consistent with scientific validity. A number of worst-case assumptions were made for the Beach Point assessment when sufficient data were not available in order to err on the conservative side during the evaluation. The use of worst case assumptions obviously introduces a significant bias in the hazard assessment.

As discussed in the ASTM guide (ASTM, 1992b), the hazard assessment procedure is an iterative process for assessing the hazard of a material to aquatic organisms. The process basically considers the relationship between a material's measured or estimated environmental concentrations and the adverse effects likely to occur to aquatic organisms. Normally the iterative process proceeds as follows. Unavailable necessary information concerning environmental concentrations and adverse effects is obtained through a stepwise process that starts with inexpensive information gathering and progresses to more comprehensive data analyses and experimental studies if necessary. At the end of each iteration (usually a maximum of three) the estimated or measured environmental concentrations are compared with

information on possible adverse effects to determine the adequacy of the available data for assessing hazard. If it is not possible to conclude that hazard is either minimal or potentially excessive after an iteration, further iterations are conducted until the hazard is adequately characterized.

4.2.2 Deviations from the Hazard Assessment Procedures

The hazard evaluation process for Beach Point deviated from some procedures recommended in the ASTM guide (ASTM, 1992b). Most of the deviations, all of which are discussed below, were dictated by the study site and the biomonitoring evaluation study design. The following deviations occurred. The ASTM guide was written for the hazard evaluation of specific chemicals or a group of chemicals that have similar biological, chemical, physical, and toxicological properties. The contaminated groundwater at Beach Point is a multiple mixture of heavy metals and chlorinated aliphatic compounds which have very different properties. Because the groundwater contained a mixture of contaminants, the complete mixture was tested rather than individual components of the mixture.

Aquatic toxicological models currently do not exist which allow for good predictions of toxicity for contaminant mixtures that contain chemicals with different modes of toxic action. Since the toxic interactions of the contaminants could not be readily determined, the concentrations of the groundwater chemicals were expressed as percent groundwater by volume rather than individual chemical mass. The mass of each contaminant in the groundwater, however, can be calculated from the chemical analysis data obtained during the biomonitoring phase of the study if desired.

The ASTM guide recommends a phased hazard iteration process (ASTM, 1992b). Several toxicological phases, however, were conducted simultaneously during the biomonitoring evaluation. That is, low-cost, medium-cost, and high-cost toxicological data collection/iteration phases were all conducted simultaneously during the biomonitoring evaluation studies.

Phase II (medium-cost information) and Phase III (high-cost information) of the ASTM hazard assessment recommend that acute and chronic toxicity tests, respectively, be conducted with estuarine organisms when a material enters estuarine waters. The Bush River is a low salinity estuary at Beach Point (Carter, 1976); however, all of the biomonitoring studies were conducted with freshwater organisms. Tests with estuarine organisms were initially considered in the design phase of the biomonitoring study. However, the pH of the groundwater at Beach Point was 4 ± 0.1 . Preliminary pH adjustment studies in freshwater showed that some heavy metals were "salted out" of solution when the pH was raised from 4 to 7. The "salting out" effect observed in

freshwater would increase at a salinity of 20 ppt which is the test salinity normally required for the EPA acute and short-term chronic test methods for estuarine organisms (Weber, 1991; Weber et al., 1989). Several heavy metals in the groundwater occurred at concentrations which could be toxic (U.S. EPA, 1986b); thus, a decision was made not to conduct estuarine tests because the tests would have underestimated the toxicity of the complex chemical mixture of the contaminants. As will be shown in the Results and Discussion Section, less toxicity occurred in several of the freshwater biomonitoring tests conducted at pH 7 relative to those conducted at pH 4. Finally, the State of Maryland Code (COMAR) considers the Bush River Area (Sub-Basin 26.08.02.03-1B) as "freshwater" for the purposes of applying numerical toxic substance criteria (see Section 5.2.3.1).

One can argue that differences exist between the toxicological sensitivities of freshwater and saltwater organisms. Indeed, there is no question that examples could be given to support either side of the argument with individual materials present in the groundwater at Beach Point. EPA has developed both freshwater and saltwater numerical water quality criteria for most of the priority pollutant heavy metals present in the groundwater at Beach Point (U.S. EPA, 1986b). As one would predict from the general toxicological literature, differences exist between EPA's freshwater and saltwater acute and chronic criteria for the various metals. Little research has been conducted on the comparative sensitivities of freshwater versus saltwater organisms exposed to multiple mixtures of heavy metals and aliphatic organics. Since toxicological differences exist between various freshwater and saltwater organisms and few data are available to predict differences in multiple mixtures, we assumed that the freshwater biomonitoring organisms would serve as reasonable surrogates for estuarine organisms in the Bush River.

It is known that the groundwater enters the Bush River through the bottom sediments (Section 3.5). One could argue that benthic as well as water column organisms should also be included in the suite of test organisms. No benthic organisms were tested in the biomonitoring evaluation of the groundwater. A recent analysis of numerical water quality criteria chemicals by EPA shows that freshwater and saltwater benthic organisms, in general, have toxicological sensitivities similar to those of water column organisms (U.S. EPA, 1993a). Thus, toxicity data for water column organisms were used to predict toxicity for benthic organisms.

4.2.3 Estimated Discharge Rate of the Surficial Aquifer

The discharge rate of the Beach Point surficial aquifer to the Bush River had to be estimated in order to estimate the environmental concentrations of the groundwater in the Bush

River. An estimated discharge rate of the surficial aquifer at Beach Point was made by Jacobs Engineering Group (JEG, 1994). The horizontal transport value, V_{hor} , for the discharge calculation was taken from a 25-h study conducted in late March 1994 by K-V Associates, Inc. (1994). V_{hor} , which was 0.27 m/d (0.89 ft/d) in a direction of 61.8°NE magnetic north, approximately 24° riverward from the main peninsula axis, was taken from well CC-33B.1 which is an intermediate depth well at Beach Point (Section 3.3). The assumption was made that V_{hor} was continuous throughout the surficial aquifer at Beach Point (JEG, 1994). V_{hor} orientation (flow direction) was assumed to be consistent with actual aquifer conditions over one complete tidal cycle (discounting short-term fluctuations due to tidal changes).

The flowing assumptions were also made for the calculation of the estimated discharge rate of the surficial aquifer at Beach Point (JEG, 1994). The surficial aquifer was considered to be homogeneous, characterized by isotropic flow conditions. The average saturated thickness of the aquifer was assumed to be 18.8 m (61.8 ft). The Bush River frontage along the peninsula is ≈448 m (1,470 ft). Approximately 366 m (1,200 ft) of frontage is directed in a NE direction; approximately 82 m (270 ft) of the northern peninsula terminus is directed in a NW direction. The aquifer was assumed to discharge the entire length of the Beach Point Peninsula fronting the Bush River. Thus, the cross-sectional area of saturated surficial aquifer fronting Bush River was assumed to be ≈8,443 m² (90,883 ft²). The length and area of the aquifer fronting the Bush River are believed to be quite conservative. A more realistic estimate of length of discharge from Beach Point is probably one third to one half, rather than the entire length (Lorah, 1994). The directional component of V_{hor} orthogonal to the Bush River frontage was used rather than assuming V_{hor} is orientated perpendicular (orthogonal) to the Bush River frontage in the surficial aquifer. That is, the measured horizontal velocity (0.27 m/d) was multiplied by a sin (24°) factor along the 366 m (1,200 ft) frontage.

The calculation of the estimated discharge rate of the surficial aquifer to the Bush River made by Jacobs Engineering Group (JEG, 1994) is:

$$\begin{aligned} \text{Discharge} &= (0.27 \text{ m/d velocity})(\sin 24^\circ)(18.8 \text{ m depth})(366 \text{ m frontage}) + (0.27 \text{ m/d velocity})(18.8 \text{ m depth})(82 \text{ m frontage}) \\ &= 1,175.5 \text{ m}^3/\text{d} \quad (41,508.7 \text{ ft}^3/\text{d}) \end{aligned}$$

4.2.4 Contaminant Distribution Assumptions in the Groundwater

As discussed in Section 3.3, chemical monitoring of the wells at Beach Point showed that the highest concentrations of heavy metals in 1988 occurred in the mid-depth (CC-33B.1) and deep (CC-33B) wells. The highest concentrations of chlorinated aliphatic organics occurred primarily in the deep well as a residual DNAPL plume. Only one shallow well (CC-33A) and one deep well (CC-33B) were studied at Beach Point during 1989. The concentrations of heavy metals and volatile organics were all higher in the deep well (CC-33B) than in the shallow well (CC-33A).

The following assumptions were made for the hazard assessment. The assumption was made for heavy metals that no retardation occurred via adsorption onto solid surfaces or trapping by clays through ion exchange. It was also assumed that no precipitation of the metals occurred when the pH shifted from 4 to neutrality when the groundwater entered the receiving stream. With regard to the DNAPLs, the assumption was made that no abiotic (chemical) or biotic (microbial) transformations occurred. We assumed that the heavy metals and residual DNAPLs were homogeneously mixed throughout the complete aquifer. In addition, the highest concentration (not the average concentration) of each contaminant measured during the four bimonthly chemical analyses (Section 4.1.6.1) was assumed to be homogeneously distributed throughout the aquifer. It was also assumed that the maximum concentrations of materials would not increase above those currently present in well CC-33B since the original sources of the contaminants were not longer present (Section 3.2). The assumption was made that the highest concentrations of the heavy metals and chlorinated aliphatics in the aquifer all moved through the sediments into the Bush River. Finally, the assumption was made that 1,176 m³/d of contaminated groundwater moved into the Bush River. We are aware that some of the groundwater flow, which is influenced by tidal stage, may be in the direction of Kings Creek (Section 3.5).

4.2.5 Screening-level Assessment of the Near-Field Dilution of the Groundwater Plume in Bush River

A screening level assessment of the near-and far-field dilution of the groundwater discharge from Beach Point was conducted by Najarian Associates, Inc. (1994) under contract to USABRDL. The primary objective of the study was to estimate, using EPA-approved screening-level techniques, the initial ("near-field") dilution and flushing rate ("far-field" dilution) of the groundwater discharge plume into the Bush River. The assessment, when combined with the discharge concentration data, provides first-order estimates of near-field concentrations. That is, it gives receiving water concentrations that would be

expected over relatively short distances from the discharge site (e.g., ones, tens, and hundreds of meters) and over short time periods (e.g., seconds and minutes).

The second objective was to compute concentrations for selected toxic constituents and to compare the results to available baseline data. As a result, potential water quality impacts can be quantified. The third objective was to assess the applicability of a Maryland "regulatory mixing zone", i.e., a localized discharge zone in which local water quality standards may be exceeded (U.S. EPA, 1991a). Such zones are routinely allowed for wastewater treatment plant discharges. If applicable to the Beach Point groundwater discharge plume, a mixing zone may allow for local exceedences of water quality standards.

4.2.5.1 Near-field Dilution Estimates

The near-field model approach was to evaluate the volume of groundwater emanating from Beach Point as a function of the potential dilution available within the Bush River. Within the near-field, it was postulated that the groundwater plume intersects the channel of the Bush River and seeps into the River as a submerged buoyant plume. Based on the available geologic information about the site, it was assumed the discharge from the surficial aquifer was limited to a relatively narrow band that parallels the interface between the Bush River and the Beach Point Peninsular. Since the groundwater plume is less dense (less saline) than the adjacent estuarine water, it would rise and subsequently mix with the ambient receiving waters. Thus, it would be diluted through mechanisms of buoyancy-based entrainment and mixing associated with ambient currents.

As discussed in Najarian Associates, Inc. (1994), the same physical mechanisms are known to dilute effluent plumes discharged from outfall diffusers of wastewater treatment plants (WWTP) and electrical generation facilities. Thus, the dilution of groundwater emanating from Beach Point can be conceptualized as the dilution of a discharge from a line diffuser. This approach allows the near-field impact of the groundwater discharge to be simulated using established plume dilution models that have been developed primarily for such outfall diffusers.

The limitations of the approach are twofold. First, most plume dilution models also simulate momentum-based entrainment, a phenomenon that has a negligible effect upon the groundwater plume. Therefore, the port configuration on the conceptualized line diffuser must be set to minimize initial momentum. Secondly, the conceptualization of the groundwater discharge surface as a line diffuser will concentrate the plume's impact within the near-field condition. As this effect reduces the dimensions of the near-field plume and calculated levels of dilution, the model result will provide a conservative estimate

of field conditions.

A series of EPA-approved, plume dilution models for surface waters are given in Muellenhoff et al. (1985). The EPA models are usually applied to discharges from line sources such as outfall diffuser manifolds. In the present case, however, the groundwater discharge is emanating from the river bed. While all EPA-approved plume models could be run, ULINE (uniform linear density flume model) was selected for the dilution calculations because it is based directly on laboratory experiments and has proved it's usefulness in recent outfall studies conducted throughout the United States (Najarian Associates, Inc., 1994). The ULINE model also neglects initial discharge momentum which is consistent with the groundwater plume movement into the Bush River. The ULINE model is designed to give dilution ratios of a discharge in a receiving stream; not distance isopleths from the source of a discharge.

Outfall dilution studies are typically conducted under "critical" minimum dilution conditions (Najarian Associates, Inc., 1994). These conditions are defined as the lowest 10th percentile ambient currents, and highest 10th percentile ambient density stratification conditions. However, Maryland regulations require that the acute aquatic life toxicity criteria be applied under conditions of mean low water and "minimum daily averaged 1-h tidal velocity" (near slack tide) (COMAR 26.08.02.05C); chronic aquatic life criteria are applied under conditions of mean water level and average tidal velocity (COMAR 26.08.02.05D). Thus, the input data were selected for the ULINE near-field model to reflect these required conditions.

To adapt ULINE to the study area, an input of site-specific data was required. The model input data fell into three categories which included 1) groundwater discharge parameters; 2) receiving water parameters; and 3) outfall diffuser parameters. The estimation of the groundwater input parameters were those given in Section 4.2.3. The ambient receiving water input parameters (e.g., ambient current velocity, salinity, temperature, and depth of discharge in the Bush River) are given in Najarian Associates, Inc. (1994). The conceptualized outfall discharge parameters for an assumed 448-m line source of buoyancy are also given in Najarian Associates, Inc. (1994).

4.2.5.2 Far-field Dilution Estimates

The Bush River receiving waters can be divided into two zones based on physical mechanisms of mixing. In the near-field, rapid mixing is caused primarily by the discharge plumes's buoyancy and initial discharge momentum, in combination with ambient currents. In the far-field, the plume is mixed passively by ambient processes such as turbulence and tidal dispersion. Far-field length scales are typically kilometer distances from

the discharge site; far-field time scales are hours, days, or longer.

Three EPA methods were initially considered for the far-field dilution estimates (i.e., the flushing rate) in the Bush River (U.S. EPA, 1985b). Far-field dilutions were calculated and reported by Najarian Associates, Inc. (1994) for the Bush River via the three methods, which included the tidal prism method, fraction of freshwater method, and dye-tracer method. The dye-tracer method was selected for further computations because dye-tracer data collected by Carter (1976) for the Bush River provided a more conservative estimate of the far-field dilution than the first two models which were based on segmented (complete mixing) approaches.

Total near-field dilutions, which correct for the influence of far-field dilutions (Fisher et al., 1979), were calculated for Maryland's acute and chronic aquatic life toxicity criteria under Spring and Fall conditions. The smaller of the Spring or Fall calculated dilution factors for both the acute and chronic toxicity conditions were then used to estimate the near-field acute and chronic receiving water concentrations of all the contaminants found in the Beach Point groundwater. The estimate for each contaminant was determined by using the highest concentration of the material found in the groundwater (see Section 4.2.4) divided by the dilution factor.

4.2.5.3 Mixing Zone Consideration

Mixing zones are areas where an effluent discharge undergoes initial dilution; they may also be extended to cover secondary mixing in the ambient waterbody. A mixing zone is an allocated impact zone where acute and chronic water quality criteria can be exceeded as long as a number of protections are maintained as required by EPA and the State (U.S. EPA, 1991a). The State of Maryland allows a mixing zone as a policy issue, but requires spatial dimensions to limit the areal extent of the mixing zone. Mixing zones may be allowed on a case-by-case basis. The concept of mixing zones is normally applied in the case of wastewater treatment facilities and power plants that discharge to waterways. Arguments will be presented in Section 5.2.3 that the mixing zone concept could be considered for the groundwater discharge in the Bush River.

SECTION 5

RESULTS AND DISCUSSION

5.1 Biomonitoring Evaluation

The Results and Discussion Section is organized as follows. The results and discussion for all of the biomonitoring systems are presented in separate sections for each test system. The endpoints/responses for each biomonitoring toxicity test are summarized in Table 2. The table is organized as Tests Nos. 1, 2, 3, and 4 which reflects the bimonthly test design. The Microtox®, Japanese medaka LC50, Ames, SCE, and Japanese medaka chronic histopathology and morphometric results are presented under the Test No. 1 column for space convenience purposes only; the tests were not quarterly tests as the heading implies. The raw data, water quality data, and statistical analyses for the biomonitoring test systems as well as the chemical analyses results are given in separate Appendices as referred to in the appropriate sections for each test system.

5.1.1 Acute Toxicity Tests

5.1.1.1 Microtox®

A summary of the Microtox® 5- and 15-min EC50 (reduction of bioluminescence) results is given in Table 2. The test data for the 100, 10, and 1% groundwater by volume samples, as well as the APG-EA diluent water are given in Appendix 1, Tables A1-1 to A1-4, respectively. The data for the Microtox® assays conducted on the groundwater buffered to a pH of ≈ 7 and the pH of the samples used in the assays are given in Appendix 2, Tables A2-1 and A2-2.

The Microtox® 5- and 15-min EC50s for 100% groundwater at pH 4 ranged from 10.6-29.1 and 17.0-45.2% groundwater by volume, respectively (Table 2). The 5- and 15-min EC50s for the 100% groundwater adjusted to pH 7 ranged from 16.9-59.1 and 18.4-95.6% groundwater by volume, respectively. The raw groundwater was less toxic at pH 7 than at pH 4. Although the 5- and 15-min EC50s were variable over the course of the study, the raw and buffered groundwater both appeared to become less toxic as the study progressed (Fig. 1 and 2). As will be shown below, the same trend was not evident in the other acute and short-term chronic test systems. Less toxicity was observed in the 15-min EC50s relative to the 5-min EC50s.

Toxicity was detected in 10% groundwater by volume one time during the study. The 5-min EC50 was 98.6% groundwater by volume; no 15-min EC50 was obtained for the same sample. The EC50 is most likely spurious because the 95% fiducial limits ranged from 88 to 300% groundwater by volume. No toxicity was

TABLE 2. SUMMARY OF THE TOXICITY ENDPOINTS/RESPONSES FOR BIOMONITORING TESTS CONDUCTED ON BEACH POINT GROUNDWATER (WELL CC-33B) FROM FEBRUARY 1993 TO DECEMBER 1993

| Bioassay | Endpoint | Value ^a | | | |
|--------------------------|--------------------------|--------------------|------------|------------|------------|
| | | Test No. 1 | Test No. 2 | Test No. 3 | Test No. 4 |
| Microtox®: | | | | | |
| 100% (pH 4) | 5-min EC50 ^b | 10.6-29.1 | N/A | N/A | N/A |
| 100% (pH 7) | 5-min EC50 ^b | 16.9-59.1 | N/A | N/A | N/A |
| 100% (pH 4) | 15-min EC50 ^b | 17.0-45.2 | N/A | N/A | N/A |
| 100% (pH 7) | 15-min EC50 ^b | 18.4-95.6 | N/A | N/A | N/A |
| 10% | 5-min EC50 ^b | 98.6 ^c | N/A | N/A | N/A |
| 10% | 15-min EC50 ^b | Not toxic | N/A | N/A | N/A |
| 1% | 5-min EC50 ^b | Not toxic | N/A | N/A | N/A |
| 1% | 15-min EC50 ^b | Not toxic | N/A | N/A | N/A |
| Diluent H ₂ O | 5-min EC50 ^b | Not toxic | N/A | N/A | N/A |
| Diluent H ₂ O | 15-min EC50 ^b | Not toxic | N/A | N/A | N/A |
| Rotifer: | | | | | |
| pH 4 | 24-h LC50 | 72 | 64 | 70 | 95 |
| pH 7 | 24-h LC50 | Not toxic | No LC50 | No LC50 | Not toxic |
| Green alga: | | | | | |
| pH 4 | 96-h EC50 ^d | 59 | 51 | 56 | 55 |
| pH 4 | NOEC ^d | 56 | 18 | 18 | 18 |
| pH 4 | LOEC ^d | 100 | 32 | 32 | 32 |
| pH 7 | 96-h EC50 ^d | No EC50 | 58 | 55 | 50 |
| pH 7 | NOEC ^d | 56 | 32 | 18 | 18 |
| pH 7 | LOEC ^d | 100 | 56 | 32 | 32 |

TABLE 2. (CONTINUED)

| Bioassay | Endpoint | Value ^a | | | |
|------------------|-------------------|--------------------|------------|------------|------------|
| | | Test No. 1 | Test No. 2 | Test No. 3 | Test No. 4 |
| Cladoceran: | | | | | |
| pH 4 | 48-h LC50 | 65 | 65 | 65 | 63 |
| pH 4 | 7-d LC50 | 65 | 65 | 65 | 42 |
| pH 4 | NOEC ^e | 18 | 10 | 10 | 10 |
| pH 4 | LOEC ^e | 32 | 18 | 18 | 18 |
| pH 7 | 48-h LC50 | Not toxic | Not toxic | No LC50 | No LC50 |
| pH 7 | 7-d LC50 | 43 | 72 | No LC50 | 40 |
| pH 7 | NOEC ^e | f | 9 | 10 | 10 |
| pH 7 | LOEC ^e | f | 9 | 18 | 18 |
| Fathead minnow: | | | | | |
| pH 4 | 96-h LC50 | 64 | 22 | 45 | 62 |
| pH 4 | 7-d LC50 | 62 | 22 | 39 | 51 |
| pH 4 | NOEC ^h | 32 | 10 | 32 | 32 |
| pH 4 | LOEC ^h | 56 | 18 | 56 | 56 |
| pH 7 | 96-h LC50 | Not toxic | Not toxic | Not toxic | Not toxic |
| pH 7 | 7-d LC50 | Not toxic | No LC50 | Not toxic | Not toxic |
| pH 7 | NOEC ^h | 56 | 56 | Not toxic | Not toxic |
| pH 7 | LOEC ^h | 100 | 100 | Not toxic | Not toxic |
| Japanese medaka: | | | | | |
| pH 4 | 96-h LC50 | No LC50 | N/A | N/A | N/A |

TABLE 2. (CONTINUED)

| Bioassay | Endpoint | Value ^a | | | |
|---------------------------------|-----------------------|--------------------|------------|------------|------------|
| | | Test No. 1 | Test No. 2 | Test No. 3 | Test No. 4 |
| Ames: | | | | | |
| 100% | Mutagenicity | i | i | Negative | Negative |
| 100% (10X) | Mutagenicity | i | i | Negative | Negative |
| 10% | Mutagenicity | Negative | Negative | Negative | Negative |
| 10% (10X) | Mutagenicity | Negative | Negative | Negative | Negative |
| 1% | Mutagenicity | Negative | Negative | Negative | Negative |
| 1% (10X) | Mutagenicity | Negative | Negative | i | i |
| Diluent H ₂ O | Mutagenicity | Negative | Negative | Negative | Negative |
| Diluent H ₂ O (10X) | Mutagenicity | Negative | Negative | Negative | Negative |
| SCE: | | | | | |
| 100% (Conc) | DNA damage | Positive | N/A | N/A | N/A |
| 100% | DNA damage | Negative | N/A | N/A | N/A |
| 10% (Conc) | DNA damage | Negative | N/A | N/A | N/A |
| Diluent H ₂ O (Conc) | DNA damage | Negative | N/A | N/A | N/A |
| FETAX: | | | | | |
| pH 4 | 4-d LC50 | Not toxic | Not toxic | Not toxic | No LC50 |
| pH 4 | 4-d EC50 ^j | No EC50 | 89 | No EC50 | No EC50 |
| pH 4 | NOEC ^k | 10 | 10 | 10 | 32 |
| pH 4 | LOEC ^k | 18 | 18 | 18 | 56 |
| pH 7 | 4-d LC50 | Not toxic | No LC50 | Not toxic | No LC50 |
| pH 7 | 4-d EC50 ^j | No EC50 | No EC50 | No EC50 | No EC50 |
| pH 7 | NOEC ^k | 18 | 18 | 18 | 18 |
| pH 7 | LOEC ^k | 32 | 32 | 32 | 32 |

TABLE 2. (CONTINUED)

| Bioassay | Endpoint | Value ^a | | | |
|------------------------------------|------------------|-----------------------|------------|------------|------------|
| | | Test No. 1 | Test No. 2 | Test No. 3 | Test No. 4 |
| Chronic histopathology and growth: | | | | | |
| 10% | 6/9-m Wet weight | No effect | N/A | N/A | N/A |
| 1% | 6/9-m Wet weight | No effect | N/A | N/A | N/A |
| Diluent H ₂ O | 6/9-m Wet weight | N/A | N/A | N/A | N/A |
| 10% | 6/9-m Std. lgth. | No effect | N/A | N/A | N/A |
| 1% | 6/9-m Std. lgth. | No effect | N/A | N/A | N/A |
| Diluent H ₂ O | 6/9-m Std. lgth. | N/A | N/A | N/A | N/A |
| 10% | 6/9-m Lesions | Negative ¹ | N/A | N/A | N/A |
| 1% | 6/9-m Lesions | Negative ¹ | N/A | N/A | N/A |
| Diluent H ₂ O | 6/9-m Lesions | Negative ¹ | N/A | N/A | N/A |

52

^a All endpoints are given as percent groundwater by volume.

^b Range of all EC50s for reduction in bioluminescent activity conducted from April-December.

^c Only one 5-min EC50 for reduction in bioluminescent activity occurred during the quarter.

^d Test endpoint- reduction in growth (cell density).

^e Test endpoint- reduction in neonate production.

^f The NOEC and LOEC could not be obtained because significant ($\alpha = 0.05$) reductions in neonate production relative to the controls occurred at all concentrations down to 32% buffered groundwater by volume, which was the lowest concentration studied.

^g The NOEC and LOEC could not be obtained because significant ($\alpha = 0.05$) reductions in neonate production relative to the controls occurred at all concentrations down to 18% buffered groundwater by volume, which was the lowest concentration studied.

^h Test endpoint- reduction in growth (dry weight) for Test Nos. 1 and 2 at pH 4 and Test No. 1 at pH 7; the test endpoint for Test Nos. 3 and 4 at pH 4 and Test No. 2 at pH 7 was an increase in mortality rather than a reduction in growth.

TABLE 2. (CONTINUED)

| Bioassay | Endpoint | Value ^a | | | |
|----------|---|--------------------|------------|------------|------------|
| | | Test No. 1 | Test No. 2 | Test No. 3 | Test No. 4 |
| i | Assay not conducted. | | | | |
| j | 96-h EC50 for malformations. | | | | |
| k | Test endpoint- increased number of malformations. | | | | |
| l | See Section 5.1.5 for additional information. | | | | |




Figure 1. Microtox® test results of raw (pH \approx 4) groundwater samples taken during the chronic histopathology study.




Figure 2. Microtox® test results of buffered (pH ≈ 7) groundwater samples taken during the chronic histopathology study.

detected during the study in the 1% groundwater by volume samples or the APG-EA diluent water.

The toxicity of the groundwater as shown by the Microtox® assay is not surprising when one considers the complex mixture of the contaminants in the groundwater (Section 5.1.6.1; Table 3). For example, the 5-min EC50s for both copper and zinc are less than the groundwater concentrations shown in Table 3 (Qureshi et al., 1982). Similarly, several of the volatile organics at concentrations below the concentrations in the raw groundwater have been shown to be toxic via the 5- and/or 15-min Microtox® analysis (Kaiser and Ribo, 1988). Chloroform, chlorobenzene, trichloroethane, and tetrachloroethane all have 5- and/or 15-min EC50s below the concentrations found in the groundwater.

5.1.1.2 Rotifer

The results of the 24-h LC50 tests at pH 4 and 7 with the rotifer are summarized in Table 2. The test data for Test Nos. 1, 2, 3, and 4 at pH 4 and 7 are given in Appendices 3 and 4, 5 and 6, 7 and 8, and 9 and 10, respectively. The 24-h LC50 values for the four acute rotifer tests conducted at pH 4 ranged from 64-95% groundwater by volume. No toxicity was found at pH 7 in the first and fourth bimonthly tests. Some mortality occurred in the 100% groundwater treatment during the second and third tests; however, no LC50s could be calculated.

5.1.1.3 Green Alga, Cladoceran, Fathead Minnow, and Japanese Medaka

The 96-h EC50s (reduction in growth) for the green alga exposed to raw groundwater at pH 4 and buffered groundwater at pH 7 are given in Table 2. The raw data at pH 4 and 7 for Test Nos. 1-4 are given in Appendices 11-18. The 96-h EC50s for reduction in growth at pH 4 ranged from 51-59% groundwater by volume (Table 2). With the exception of Test No. 1 in which no EC50 could be calculated because <50% reduction in growth occurred in the study, the 96-h EC50s were essentially the same in the groundwater buffered to pH 7. The EC50s at pH 7 in Test Nos. 2-4 ranged from 50-58% groundwater by volume.

The 48-h acute LC50 toxicity data for the cladoceran at pH 4 and 7 are summarized in Table 2. The raw data, including test water quality, for Test Nos. 1-4 at pH 4 and 7 are given in Appendices 19-26. Groundwater at pH 4 was acutely toxic to the cladoceran (Table 2). The 48-h LC50s ranged from 63-65% groundwater by volume for the four tests. Buffered groundwater was not toxic in Test Nos. 1 and 2. Some toxicity occurred in Tests Nos. 3 and 4; however, no 48-h LC50 could be calculated because <50% mortality occurred in 48 h.

TABLE 3. SUMMARY OF THE FOUR BIMONTHLY CHEMICAL ANALYSES
(RANGE OF CONCENTRATIONS) CONDUCTED ON RAW BEACH
POINT GROUNDWATER (WELL CC-33B) FROM FEBRUARY
1993 TO DECEMBER 1993

| Parameter | Concentration | Unit |
|------------------------------|---------------|---------------------------|
| <u>General Water Quality</u> | | |
| Alkalinity | <1 - 21 | mg/L as CaCO ₃ |
| Hardness | 170 - 230 | mg/L as CaCO ₃ |
| Conductivity | 1,600 - 2,020 | μs/cm |
| Total Suspended Solids | <5 | mg/L |
| Total Organic Carbon | 2 - 4 | mg/L |
| Calcium | 16.5 - 21 | mg/L as Ca |
| Magnesium | 30 - 43 | mg/L as Mg |
| Manganese | 1.7 - 2.1 | mg/L as Mn |
| Potassium | 2.5 - 4.1 | mg/L as K |
| Sodium | 257 - 350 | mg/L as Na |
| Sulfate | 62 - 91 | mg/L as SO ₄ |
| Sulfide | <1 | mg/L as S |
| Chloride | 490 - 580 | mg/L as CL |
| Ammonia-Nitrogen | <0.1 | mg/L as N |
| Nitrate-Nitrogen | 0.1 - 2.5 | mg/L as N |
| Nitrite-Nitrogen | <0.01 | mg/L as N |
| Phosphorus | 0.12 - 2.64 | mg/L as P |
| Fluoride | 0.2 - 1.2 | mg/L as F |
| Iron | 4.6 - 7.7 | mg/L as Fe |
| Cyanide | <0.01 - 0.06 | mg/L as CN |
| Bromide | 0.4 - 1.5 | mg/L as Br |
| pH | 3.9 - 4.1 | Std. Unit |

TABLE 3. (CONTINUED)

| Parameter | Concentration | Unit |
|---------------------------------|----------------|------------|
| <u>Heavy Metals^a</u> | | |
| Aluminum | 810 - 1,500 | µg/L as Al |
| Antimony | <5 | µg/L as Sb |
| Arsenic | <5-7 | µg/L as As |
| Beryllium | <5 | µg/L as Be |
| Boron | 50 - 70 | µg/L as B |
| Cadmium | 0.8 - 1.4 | µg/L as Cd |
| Chromium | <2 | µg/L as Cr |
| Cobalt | 80 - 110 | µg/L as Co |
| Copper | 5.4 - 8 | µg/L as Cu |
| Lead | <5 | µg/L as Pb |
| Mercury | <0.5 | µg/L as Hg |
| Molybdenum | <10 | µg/L as Mo |
| Nickel | 120 - 180 | µg/L as Ni |
| Selenium | <5 | µg/L as Se |
| Silver | <1 | µg/L as Ag |
| Thallium | <5 | µg/L as Ti |
| Tin | <50 | µg/L as Sn |
| Zinc | 210 - 310 | µg/L as Zn |
| <u>Volatile Organics</u> | | |
| Chlorobenzene | 2 | µg/L |
| Chloroform | 10 - 12 | µg/L |
| 1,1,2-Trichloroethane | 37 - 110 | µg/L |
| 1,1,1,2-Tetrachloroethane | 3 | µg/L |
| 1,1,2,2-Tetrachloroethane | 9,000 - 17,000 | µg/L |

TABLE 3. (CONTINUED)

| Parameter | Concentration | Unit |
|--------------------------------|---------------|------|
| <u>Volatile Organics Con't</u> | | |
| 1,1-Dichloroethene | 1 | µg/L |
| cis-1,2-Dichloroethene | 110 - 640 | µg/L |
| Trichloroethene | 450 - 1,600 | µg/L |
| Tetrachloroethene | 30 - 90 | µg/L |
| Vinyl Chloride | 2 | µg/L |

^a The metal concentrations are total metal; not dissolved metal.

The 96-h LC50 data for the fathead minnow at pH 4 and 7 are summarized in Table 2. The raw data, including test water quality, for Test Nos. 1-4 at pH 4 and 7 are given in Appendices 27-34. Groundwater at pH 4 was acutely toxic to the fathead minnow (Table 2). The 96-h LC50s ranged from 22-64% groundwater by volume for the four tests. Buffered groundwater at pH 7 was not toxic in any test after a 96-h exposure.

One 96-h LC50 toxicity test was conducted with Japanese medaka at pH 4 (Table 2; Appendix 35). The groundwater was toxic at pH 4 after a 96-h exposure; however, a 96-h LC50 could not be calculated because <50% mortality occurred in 96 h. An acute toxicity test was not conducted at pH 7 since the groundwater was not acutely toxic to Japanese medaka at pH 4.

The acute toxicity of the groundwater to the rotifer, cladoceran, and fathead minnow at pH 4 may be attributable to the heavy metals in the groundwater (Table 3). Several EPA priority pollutant heavy metals (cadmium, copper, nickel, and zinc) were found in the groundwater. The concentration of zinc in the groundwater (when adjusted for hardness) exceeded the EPA numerical water quality criterion of 120 µg/L for freshwater invertebrates and fish (U.S. EPA, 1987). Furthermore, metals such as copper and zinc exist primarily as divalent cations at a pH of 4 which is the most toxic form of the metal (Lee, 1973; Sprague, 1985). It is well established that the toxicity of metals in chemical mixtures is additive for many aquatic animals (Marking, 1985). It is likely that the toxicity observed in the study may have been additive or greater than additive (de March,

1988). The reduction and/or elimination of toxicity at pH 7 for the same test organisms is most likely related to the reduction in toxicity of heavy metals as they shift from a divalent cation at pH 4 to less toxic species at pH 7 (Lee, 1973). In contrast to the acute toxicity of the groundwater at pH 4 to the rotifer, cladoceran, and fathead minnow, a 96-h LC50 could not be calculated for the Japanese medaka because <50% mortality occurred.

No single heavy metal at the concentrations present in the groundwater can account for the toxicity observed in the green alga (U.S. EPA, 1986b). Few data are available on the possible joint toxicity of heavy metals or organics to green algae (Faust et al., 1994). The toxicity at pH 4 and 7 was essentially the same (Table 2) which suggests that heavy metals may not be important toxicologically to the alga. No single priority pollutant organic for which there are toxicity data can account for the toxicity observed in the alga (U.S. EPA, 1986b). One can only speculate that the toxicity of the groundwater to the alga may be related to 1) the joint toxicity of the metals, 2) joint toxicity of the organics and/or 3) interaction of the metals and organics.

5.1.2 Short-term Chronic Toxicity Tests

5.1.2.1 Green Alga

The NOECs and LOECs (reduction in cell density) for the green alga exposed to groundwater at pH 4 and 7 are summarized in Table 2. The test data and statistical analyses for Test Nos. 1-4 are given in Appendices 11-18. At pH 4, the NOECs for the four tests ranged from 18-56% groundwater by volume. The LOECs ranged from 32-100% groundwater by volume. At pH 7, the NOECs and LOECs, respectively, ranged from 18-56 and 32-100% groundwater by volume. With the exception of Test No. 2, the NOECs and LOECs were exactly the same at pH 4 and 7 for the same test period. The NOECs and LOECs were less at pH 7 than at pH 4 in Test No. 2 only. Thus, with the exception of one test, no difference in algal toxicity was found for groundwater at pH 4 and pH 7. Some variation in toxicity was found between the four sets of tests. The lowest NOEC found in any test was 18% groundwater by volume. As stated above in Section 5.1.1.3, one can only speculate that the toxicity of the groundwater to the alga may be related to the 1) joint toxicity of the metals, 2) joint toxicity of the organics and/or 3) interaction of the metals and organics.

5.1.2.2 Cladoceran

The 7-d LC50s, NOECs, and LOECs for the cladoceran exposed to groundwater at pH 4 and 7 are summarized in Table 2. The test data and statistical analyses for Test Nos. 1-4 are given in Appendices 19-26. The groundwater at pH 4 was toxic in all

tests. The short-term chronic 7-d LC50s at pH 4 ranged from 42-65% groundwater by volume. The NOECs (reduction in neonate production) at pH 4 ranged from 10-32% groundwater by volume while the LOECs ranged from 18-32% groundwater by volume in the four tests.

The groundwater at pH 7 was essentially as toxic to the cladoceran as it was at pH 4. The 7-d LC50s ranged from 40-72% groundwater by volume in three of the four tests. Toxicity was observed in Test No. 3; however, the 7-d LC50 could not be calculated because <50% mortality occurred after 7 d of exposure. The NOECs and LOECs in Test Nos. 1 and 2 could not be determined because significant reductions ($\alpha = 0.05$) in neonate production occurred down to 32% and 18% groundwater by volume, respectively, which were the lowest concentrations used in the study (Appendix 20, Table A20-3 and Appendix 22, Table A22-3). The NOECs and LOECs in Test Nos. 3 and 4 were both 10% and 18%, respectively. The NOECs and LOECs were the same in Test Nos. 3 and 4 in both raw and buffered groundwater.

5.1.2.3 Fathead Minnow

The 7-d LC50s, NOECs, and LOECs for the fathead minnow exposed to groundwater at pH 4 and 7 are summarized in Table 2. The test data and statistical analyses for Test Nos. 1-4 are given in Appendices 27-34. The groundwater at pH 4 was toxic in all tests. The short-term chronic 7-d LC50s at pH 4 ranged from 22-62% groundwater by volume. The NOECs at pH 4 ranged from 10-32% groundwater by volume while the LOECs ranged from 18-56% groundwater by volume in the four tests. The NOEC and LOEC endpoints at pH 4 were a reduction in growth for Test Nos. 1 and 2 and an increase in mortality in Test Nos. 3 and 4.

The groundwater was less toxic when buffered to pH 7 (Table 2). With regard to the 7-d LC50s at pH 7, no statistically significant ($\alpha = 0.05$) mortality occurred in three of the four tests. Some mortality was observed in Test No. 2; however, a 7-d LC50 could not be calculated because <50% mortality occurred in 7 d of exposure. A significant reduction ($\alpha = 0.05$) in growth occurred in Test No. 1 while a significant increase in mortality occurred in Test No. 2. The NOECs for Test Nos. 1 and 2 were 56% groundwater by volume, while the LOECs were 100% groundwater by volume. No NOECs or LOECs were obtained in the third and fourth tests because statistically significant ($\alpha = 0.05$) increases in mortality or reductions in growth did not occur.

The chronic toxicity observed for the cladoceran and fathead minnow may be related to several heavy metals and at least one organic present in the groundwater (Table 3). As discussed below in Section 5.1.6.1, cadmium, nickel, and zinc concentrations in the groundwater are equal to or exceed the EPA freshwater chronic numerical water quality criteria of 1.1, 160, and 110 $\mu\text{g/L}$

(hardness dependent criteria; 100 mg/L as CaCO₃ used), respectively (U.S. EPA, 1980a; 1984a; 1987). None of the priority pollutant organics found in the groundwater currently have numerical water quality criteria values because insufficient data exist to develop criteria (Potts, 1994). However, 1,1,2,2-tetrachlorethane concentrations found in the groundwater exceed the EPA freshwater chronic LOEC (U.S. EPA, 1980b). Thus, it is likely that the chronic toxicity observed in the cladoceran and fathead minnow tests is a combination of metals and organics.

In contrast to the general reduction in acute toxicity when the rotifer and fathead minnow were tested in buffered groundwater, the chronic toxicity NOECs and LOECs values for the cladoceran were the same at pH 4 and 7 (Table 2). Thus, the suggestion above that toxicity attributable to heavy metals is reduced at the higher pH does not appear to be valid for the cladoceran in the chronic tests. The reason for this observation is not clear.

5.1.3 Genotoxicity Tests

5.1.3.1 Gene Mutation Assay

The results of the Ames mutagenicity assays are summarized in Table 2. Both unconcentrated and concentrated (10X) groundwater Ames assays were conducted on 10% groundwater by volume, 1% groundwater by volume, and control water in Test Nos. 1 and 2; 100% groundwater, 10% groundwater by volume, and control water were assayed in Test Nos. 3 and 4. The reference for each study data report is given in Appendix 36 so that a copy can be obtained if further information is desired; the actual data reports were not included in the Appendix because of the excessive length of each report.

All of the unconcentrated and concentrated (10X) 100% groundwater, 10% groundwater by volume, 1% groundwater by volume, and diluent water assays were found to be non-mutagenic (negative) with tester strains TA98 and TA100 in both the presence and absence of an exogenous metabolic activation system of mammalian microsomal enzymes derived from Aroclor-induced rat liver (S9 mix). One deviation from the above results was found in the unconcentrated APG-EA diluent water in Test No. 1. With tester strain TA98 in the absence of S9 mix, the response was evaluated as marginal in two trials due to the presence of 1 dose with an MR \geq 1.5 which was not accompanied by a dose response.

5.1.3.2 Sister Chromatid Exchange Assay

The sister chromatid exchange (SCE) assay was performed once on concentrated samples (\approx 50,000X) of 100% groundwater, 10% groundwater by volume, APG-EA diluent water, and once on an unconcentrated 100% groundwater sample (Table 2). The actual

data reports were not included in the Appendix of this report because of their excessive length. The reference for each study report is given in Appendix 37.

The concentrated extract of 100% groundwater was considered positive for inducing sister chromatid exchange in Chinese hamster ovary cells under the nonactivation conditions of the study and weakly positive under the activation (exogenous metabolic activation system of mammalian microsomal enzymes derived from Aroclor-induced rat liver S9 mix) conditions of the study (Table 2). The unconcentrated sample of 100% groundwater was considered negative for inducing sister chromatid exchange in Chinese hamster ovary cells under the activation and nonactivation conditions of the assay. The concentrated extracts of the 10% groundwater by volume and APG-EA diluent water were also considered negative for inducing sister chromatid exchange in Chinese hamster ovary cells under the activation and nonactivation conditions of the assay.

A positive response was found for 100% groundwater in the SCE assay when the groundwater sample was concentrated $\approx 50,000X$. The SCE assay was negative for unconcentrated 100% groundwater. It is not clear that the SCE positive response in the 50,000X concentrated sample is important to aquatic organisms. As discussed below in Section 5.2, the potential for the heavy metals and aliphatic organics present in the groundwater to bioaccumulate is <100 -fold because the $\log K_{ow}$ s of the materials are ≤ 3 . It is clear that no chemical or physical process in the Bush River could cause a 50,000X increase in the concentration of the materials in the near-field of the groundwater discharge. Thus, organisms in the immediate vicinity of the groundwater discharge would never be exposed to a 50,000X increase in the contaminants.

5.1.4 Developmental Toxicity Test

The 4-d LC50, 4-d EC50 (malformations), NOEC, and LOEC results for the FETAX assays conducted in groundwater at pH 4 and 7 are summarized in Table 2. The test data, statistical analyses, and types and numbers of malformed embryos that occurred after 96 h of exposure for Test Nos. 1-4 are given in Appendices 38-45. Little embryo lethality occurred in the groundwater at pH 4 or 7 (Table 2). The raw groundwater was not toxic to the embryos at pH 4 in the first three tests. Some toxicity occurred in Test No. 4; however, a LC50 could not be calculated because $<50\%$ mortality occurred. The buffered groundwater was not toxic in two of the four test. Some mortality occurred in Test Nos. 2 and 4; however, a LC50 could not be calculated.

Significant ($\alpha = 0.05$) embryo malformations occurred in the raw and buffered groundwater in all assays. A 96-h EC50

(malformations) of 89% groundwater by volume was obtained in the second test at pH 4; 96-h EC50s could not be calculated for any of the other tests. The NOECs and LOECs for the four groundwater tests at pH 4 were 10 and 18% groundwater by volume, respectively, in the first three tests and 32 and 56% groundwater by volume in Test No. 4. The NOECs and LOECs for the four groundwater tests at pH 7 were all 18 and 32% groundwater by volume, respectively. With the exception of the fourth test, the buffered groundwater was less toxic than the raw groundwater. For some unexplainable reason, the raw groundwater was less toxic than the buffered groundwater in Test No. 4.

The types of malformed embryos (as described by Bantle et al., 1991) after 96 h of exposure in raw groundwater were primarily coiled guts, multiple edema, and abdominal edema (see last table in each of Appendices 38-45). Severe, notochordal and facial malformations were also observed in <10% of the total malformations. Fewer malformations were observed in the buffered groundwater tests; however, the same types of malformations that occurred in the raw groundwater were observed in the buffered groundwater assays (primarily coiled guts, multiple and abdominal edema with <10% severe, notochord, and facial). The incidences of malformations were greater at the higher test concentrations in both the raw and buffered groundwater.

The developmental toxicity found in the FETAX assays is most likely related to the heavy metals present in the groundwater. Several heavy metals, including copper, cadmium, and zinc, have been shown to cause developmental problems in lower vertebrate aquatic organisms (Weis and Weis, 1989). Dawson et al. (1985) found that mixtures of heavy metals (copper, cadmium, lead, and zinc) from acidic mine sources caused teratogenic effects and mortality when evaluated by FETAX. When the pH was adjusted from lows which ranged from 3.2 to 5.9 to pH 7, toxicity and teratogenicity decreased. The same response occurred in the present study with the exception of Test No. 4 in which the raw groundwater was less toxic than the buffered groundwater. The possible role of the organics in the groundwater is not clear since FETAX data do not exist for the individual materials (Bantle, 1994).

5.1.5 Chronic Histopathology and Growth Test

The Japanese medaka growth and histopathology results at 6 and 9 months of exposure to 10% groundwater by volume, 1% groundwater by volume, and diluent water are briefly summarized in Table 2. The wet weight and standard length raw data, means, and standard deviations of the means for each replicate tank in each treatment at 6 and 9 months are given in Appendix 46, Tables A46-1 and A46-3, respectively. The statistical analyses of the growth data at 6 and 9 months are given in Appendix 46, Tables A46-2 and A46-4.

No significant difference ($\alpha = 0.05$) in wet weight or standard length was found between treatments after 6 and 9 months of exposure. With regard to the 9-month length data only, a statistically significant difference ($\alpha = 0.003$) in standard length was found using the nonparametric Kruskal Wallis test (Appendix 46, Table A46-4). However, when Dunn's multiple comparison test was used to enumerate the difference between the control and each experimental treatment, no statistically significant difference ($\alpha = 0.05$) was found. Standard length was judged not to be affected by a 9-month exposure to the groundwater because 1) Dunn's test could not discern any statistical difference ($\alpha = 0.05$) between groups and 2) the numerical differences between mean lengths in the study groups were quite small. For example, the mean standard lengths of the four replicates combined at each treatment were 32.0, 31.8, and 32.5 mm respectively, for the control, 1% groundwater by volume, and 10% groundwater by volume groups.

The cumulative mortality of Japanese medaka during the 9-month study is shown graphically in Figure 3. Percent cumulative mortality is given in Figure 4. Cumulative percent mortality at 6 months for the controls, fish exposed to 1% groundwater by volume, and 10% groundwater by volume was 4.5, 2.5, and 2.1%, respectively. Cumulative percent mortality at 9 months for the controls, fish exposed to 1% groundwater by volume, and 10% groundwater by volume was 22.1, 11.7, and 7.2%, respectively. To the authors knowledge, there are no test mortality acceptability criteria for a 9-month test. If one uses the mortality acceptability criteria for early life stage (ELS) toxicity tests which run for 1-2 months after hatch or fry swim-up, the mortality observed in this study falls within ELS acceptability criteria (Goodman, 1986). For example, the ELS test acceptability criteria for all eight freshwater species listed in the draft ASTM standard guide (Japanese medaka are not include in the guide) run from 60-75% (Goodman, 1986). That is, for a test to be acceptable, 60-75% of the control organisms must be alive at the end of the study depending on the species being studied. Excluding fish sacrificed at 6 months, 77.9% of the control Japanese medaka were living at the end of the 9-month period.

The histopathological findings for the 6- and 9-month exposures are described in detail in the pathology report by Experimental Pathology Laboratories, Inc. (1994). A summary of the findings is given in Appendix 47. Briefly, the major conclusions of the histopathology study are as follows. After 6 months of exposure to control water, 1% groundwater by volume, and 10% groundwater by volume, liver neoplasia (hepatocellular adenoma and carcinoma) occurred only among Japanese medaka that had been previously initiated with DEN. Seven of 40 medaka were affected in the fish exposed to 10% groundwater by volume as compared to 2 of 40 among the controls and 1 of 40 Japanese medaka exposed to 1% groundwater by volume.

After 9 months of exposure, hepatocellular adenomas or carcinomas, single or multiple, occurred in DEN-initiated Japanese medaka that had been exposed to control water, 1% groundwater by volume, and 10% groundwater by volume. There were as many neoplasms among the controls (8 of 122) as among the 10% exposure group (8 of 144). There were two liver neoplasms (1 hepatocellular adenoma and 1 carcinoma) in Japanese medaka exposed to 10% groundwater by volume (no DEN exposure). Two additional liver neoplasms not related to hepatocytes occurred in the DEN 10% groundwater by volume group. One was a cholangioma and the other was a carcinoma, cell of origin unknown. In the DEN 1% groundwater by volume exposure groups there were five (5 of 144) hepatocellular neoplasms, all adenomas. At 9 months there was no dose-related response for liver neoplasia in Japanese medaka exposed to groundwater.

Other neoplasms that occurred sporadically among the various exposure groups at 6 and 9 months included lymphosarcoma, seminoma, thyroid follicular cell adenoma, thyroid follicular cell carcinoma, hemangioma of the gills, and a gallbladder papilloma.

Granulomatous inflammation of the subcutaneous tissue of the lower and/or upper jaw occurred in small numbers of male fish at 6 months. The same lesion occurred among Japanese medaka of all exposure groups at 9 months. The incidence was substantially greater among males than among females. For example, 144 males overall versus 17 females were diagnosed with granulomatous inflammation of the lower jaw. Granulomatous inflammation or granulomas occurred in a variety of additional tissues among all exposure groups. The reason for the higher incidence of these lesions in males versus females is not known. Trauma may be an initiating factor in the granulomatous lesions of the jaws, but the reason why the jaws of these fish should be traumatized is not known.

Granulomas in the heart, hematopoietic tissue, liver, and spleen occurred in the highest incidence in the controls and in the lowest incidence in Japanese medaka exposed to 10% groundwater by volume. The reasons for this lower incidence among fish exposed to 10% groundwater by volume as compared to the controls is not known.

The DEN-induced hepatic neoplasia found in this study have been observed by several investigators using the Japanese medaka as a model to evaluate carcinogenic potential of selected chemicals in aquatic organisms (for ex., Bunton, 1990; Lauren et al., 1990; Hinton et al., 1988). Protocols for studying DEN hepatocarcinogenicity vary but generally involve exposure to 10-100 mg/L DEN in ambient water for several days or weeks followed by several weeks for tumor development (Hinton et al., 1985). The DEN exposure period in this study was 48 h, which is




Figure 3. Cumulative mortality of Japanese medaka during the 9-month chronic histopathology exposure study.




Figure 4. Percent cumulative mortality of Japanese medaka during the 9-month chronic histopathology exposure study.

substantially shorter than the exposure periods normally used by other investigators. Ichikawa and Takayama (1979) have shown that the length of exposure and length of grow-out can enhance hepatic neoplasia induced by DEN.

Little, if any, data are available on the potential carcinogenicity of heavy metals to the Japanese medaka (Hawkins, 1994). Similarly, short-chain halogenated hydrocarbons have received little study relative to other organic groups, e.g., nitroso compounds, polynuclear aromatic hydrocarbons, and aromatic amines (Walker et al., 1985). 1,1,2,2-Tetrachloroethane was not found to be carcinogenic to Japanese medaka exposed to concentrations up to 14 mg/L in a recent study by Hawkins (1991).

The four bimonthly comprehensive chemical analyses for the 9-month study are given in Appendix 48 (see Section 5.1.6.1). The general water quality data, means, and standard deviations of the means for each replicate tank in each treatment for the 9-month study are given in Appendix 49 (see Section 5.1.6.2).

5.1.6 Chemical Analyses

5.1.6.1 Comprehensive Chemical Analyses

A summary of the raw groundwater general water quality, heavy metals, and volatile organics measured in the samples of the four bimonthly comprehensive chemical analyses is given in Table 3. The range of the lowest and highest concentrations of the four analyses is presented. The comprehensive results of each bimonthly chemical analysis, including the values for 100% groundwater, 10% groundwater by volume, 1% groundwater by volume, and diluent water are given in Appendix 48, Tables A48-1, A48-2, A48-3, and A48-4 for Test Nos. 1, 2, 3, and 4, respectively. The tables in Appendix 48 include the test method and detection limit for each chemical. In contrast to Table 3, which contains only the range of materials actually measured in the samples, all materials measured and quantified as well as materials not detected during analysis are included in Appendix 48.

No compounds in the following groups were detected at EPA's quantitation limits for groundwater: 1) acid or base/neutral compounds; 2) pesticides; 3) herbicides; or 4) organo-phosphorus pesticides. The following munitions were not detected at a quantitation limit of 50 $\mu\text{g/L}$: 1) octahydro-1,3,5,7-tetranitro-1,3,5,7-tetrazocine (HMX); 2) hexahydro-1,3,5-trinitro-1,3,5-triazine (RDX); 3) 1,3,5-trinitrobenzene (TNB); 4) N,2,4,6-tetranitro-N-methylaniline (tetryl); 5) trinitrotoluene (TNT); 6) 2,4-dinitrotoluene (2,4-DNT); or 7) 2,6-dinitro-toluene (2,6-DNT).

The general water chemistry parameters of the groundwater summarized in Table 3 show that the groundwater is a moderately

hard water (hardness = 170-230 mg/L as CaCO_3). The pH of 3.9-4.1 is low relative to that which occurs in most surface waters. Some surface waters high in tannic acid or those waters impacted by acid rain may also have pH values in the same range (Baker et al., 1990). Ammonia-nitrogen was <0.1 mg/L in all samples; thus, nonionized ammonia would not be expected to play a role in toxicity (Thurston et al., 1979).

Several EPA priority pollutant heavy metals were found in the groundwater (Table 3). The concentrations of cadmium, copper, nickel, and zinc exceeded one or more of EPA's numerical water quality criteria for the specific metal. Cadmium, nickel, and zinc concentrations were equal to or exceeded the EPA freshwater chronic numerical water quality criteria of 1.1, 160, and 110 $\mu\text{g/L}$ (hardness dependent criteria; 100 mg/L as CaCO_3 used), respectively (U.S. EPA, 1980a; 1984a; 1987). Zinc also exceeded the freshwater acute criteria of 120 $\mu\text{g/L}$ (hardness dependent criteria; 100 mg/L as CaCO_3 used). Copper exceeded EPA's saltwater acute criterion of 2.9 $\mu\text{g/L}$ (U.S. EPA, 1984b); a chronic saltwater copper criterion does not exist (Potts, 1994). Aluminum was also present at high concentrations in the groundwater; however, EPA has not established numerical water quality criteria for the metal in freshwater or saltwater under acute or chronic exposure conditions (Potts, 1994).

A number of chlorinated aliphatic compounds were found in the groundwater (Table 3). Several of the organics were EPA priority pollutants. None of the priority pollutant organics found in the groundwater currently have numerical water quality criteria values because insufficient data exist to develop criteria (Potts, 1994). EPA does give the LOEC for several of the compounds where criteria are not available. 1,1,2,2-Tetrachlorethane concentrations found in the groundwater exceed the EPA freshwater chronic LOEC of 2,400 $\mu\text{g/L}$; no freshwater acute LOEC is given by EPA (U.S. EPA, 1980b). None of the 10 volatile organics found in the groundwater have octanol water partition coefficients ($\log k_{ow}$ or $\log P$) >3 (Table 4). Thus, bioaccumulation is not a potential toxicological problem (see Section 5.2.1).

5.1.6.2 Routine Water Quality Analyses

The raw data, mean, and standard deviation of the mean for each exposure tank in the chronic histopathology study are given in Appendix 49. The methods for determining the routine water quality are given in Appendix 50. Although some of the water quality parameters varied as a function of the three treatments, water quality within a given treatment was quite consistent over the 9-month study. The continuous temperature measurement recorded from Tank No. 3 during the 9-month study verified the constant temperature conditions which were measured daily in each exposure tank.

TABLE 4. LOG OCTANOL WATER PARTITION COEFFICIENTS OF THE ORGANIC CONTAMINANTS DETECTED IN WELL CC-33B

| Contaminant | Log K_{ow} |
|---------------------------|------------------|
| Chlorobenzene | 2.8 ^a |
| Chloroform | 1.9 ^b |
| 1,1,2-Trichloroethane | 2.2 ^b |
| 1,1,1,2-Tetrachloroethane | 3.0 ^c |
| 1,1,2,2-Tetrachloroethane | 2.4 ^b |
| 1,1-Dichlorethene | 2.1 ^a |
| cis-1,2-Dichloroethene | 1.8 ^d |
| Trichloroethene | 2.4 ^b |
| Tetrachloroethene | 2.9 ^b |
| Vinyl chloride | 1.4 ^c |

^a Value taken from Howard (1989).

^b Value taken from U.S. EPA (1991b).

^c Value taken from Mackay et al. (1993).

^d Value taken from Howard (1990).

5.2 Hazard Assessment Evaluation

5.2.1 Toxicity

As discussed in Section 4.2, the ASTM guide recommends a phased hazard iteration process (ASTM, 1992b). Phase I (low-cost), Phase II (medium-cost), and Phase III (high-cost) toxicological data collection/iterations were all conducted simultaneously during the biomonitoring evaluation studies. Thus, the decision that normally would be made at the end of each phase concerning whether or not the potential hazard should be assessed as either 1) minimal, 2) potentially excessive, or 3) uncertain was not made until the end of the biomonitoring study when all of the toxicological studies were complete.

One important decision regarding bioaccumulation in aquatic organisms was made at the end of Phase I. The hazardous heavy metals present in the groundwater (Table 3) do not bioaccumulate to any degree in their metallic form in tissues (Williamson et al., 1993). Some heavy metals will bioaccumulate in the shells of benthic organisms; however, bioaccumulation in the shells is not believed to interfere with the normal physiological function of the organisms. With regard to the halogenated organics present in the groundwater, the log K_{ow} s (log P) of all the organics are ≤ 3 (Table 4). Bioaccumulation of a material up to 100-fold above background (bioconcentration factor or BCF = 100) can occur when the log K_{ow} s = 3 (U.S. EPA, 1991c). Thus,

bioaccumulation was judged not to be an issue in the hazard assessment.

Toxicity was detected at various groundwater concentrations by all of the biomonitoring systems at pH 4 with the exception of two assays (Section 5.1). The Ames test for mutagenicity was negative in all cases including 100% groundwater as was the 9-month Japanese medaka chronic growth and histopathological test up to 10% groundwater by volume. A positive response was found in 100% groundwater in the SCE assay when the groundwater sample was concentrated $\approx 50,000\times$; the SCE assay was negative in 100% groundwater which was not concentrated. As discussed in Section 5.1.3.2, the SCE findings for concentrated 100% groundwater were not considered to be important to aquatic organisms in the near-field of the groundwater discharge.

The lowest concentration of groundwater that caused no observable adverse effect at pH 4, in the test systems in which the NOEC value could be determined, was 10% groundwater by volume. A NOEC of 10% groundwater by volume occurred in 3 out of 4 tests in both the cladoceran and FETAX assays; it occurred once in the fathead minnow test. The NOEC concentration was higher at pH 7 in both the fathead minnow and FETAX assays. The 10% groundwater by volume NOEC for cladoceran at pH 4, however, did not change when the organism was exposed to buffered groundwater at pH 7.

Maximum acceptable toxicant concentrations (MATC) are normally required for chronic toxicity estimates in a hazard assessment (ASTM, 1992b). The MATC is the highest concentration of a material that would have no statistically significant observed effect on the survival, growth, or reproduction of a test species during continuous exposure throughout a life-cycle or partial life cycle test. The short-term chronic tests with the green alga and cladoceran can be used to estimate the MATC for a plant and invertebrate (Weber et al., 1989). The green algal test is a multi-generation assay. The cladoceran test is a partial life cycle test. The MATC for the alga and cladoceran would be 18 and 10% groundwater by volume, respectively.

One can estimate chronic toxicity by calculating the acute-to-chronic ratio (ACR) which is simply the ratio of the LC50 to the NOEC (U.S. EPA, 1991a). The ACRs for the fathead minnow at pH 4 ranged from 1.4 to 2.0 in the four bimonthly tests. Thus, chronic toxicity of the groundwater to the fathead minnow would be estimated to be ≈ 2 -fold greater than the acute toxicity. The estimated MATC of the groundwater to the fathead minnow would be $\approx 11\%$ groundwater by volume. Although the 9-month Japanese medaka growth test is not a partial life cycle test by definition, the groundwater had no effect on Japanese medaka growth or chronic histopathology at 10% groundwater by volume after a 9-month exposure.

The estimated MATCs for the alga, invertebrate, and fish in the biomonitoring study suggest that the groundwater would not be harmful at a concentration of 10% groundwater by volume. Likewise, no genotoxicity, developmental toxicity, or chronic histopathology occurred at 10% groundwater by volume. Thus, the biomonitoring data suggest that chronic toxicity may not occur in the receiving stream at a dilution of 10:1. Chronic toxicity, however, would be predicted to occur in the Bush River if the groundwater entered the receiving stream at the concentrations which occur in well CC-33B.

In conclusion, if all of the conservative assumptions presented in Section 4.2 concerning the groundwater are assumed to occur (e.g., homogeneously mixed heavy metals and DNAPLs, no chemical or biotic transformations of the materials, large release volumes through the sediments to the Bush River, etc.), the groundwater would be considered a potentially excessive hazardous material to the benthic biota of the Bush River (ASTM, 1992b). The hazard to water column aquatic organisms would rapidly dissipate as the materials mixed in the receiving stream. We argue below that even if all of the conservative assumptions in Section 4.2 for the groundwater were used, that potential water quality impacts would be minimal if a mixing zone was granted by the State of Maryland which allows for local exceedences of water quality standards. The State of Maryland allows on a case-by-case basis a mixing zone where acute and chronic water quality criteria may be exceeded (see Section 5.2.4). Standards for a mixing zone for acute and chronic sediment quality do not currently exist.

5.2.2 Near-field Dilution in the Bush River

The results of the ULINE near-field model study are given in Najarian Associates, Inc. (1994). A copy of the Najarian report is included in this report as Appendix 51. The ULINE near-field model results indicate that a near-field dilution ratio of 159:1 is appropriate for application of the Maryland Department of the Environment's (MDE) acute aquatic life toxicity criteria. The model shows that a far-field dilution factor of 1,006:1 is appropriate for application of chronic toxicity criteria. A dilution factor of 159:1 means that when 1 part of Beach Point groundwater is diluted in 159 parts of Bush River water, MDE's acute criteria are met at the edge of the acute mixing zone. MDE's acute and chronic aquatic life criteria are the same as those derived by EPA. Note that far-field flushing parameters are not included in the ULINE near-field model. When near-field dilution is corrected for the influence of far-field flushing, the resulting total near-field dilution ratios will decrease (see below).

A model sensitivity analysis was performed to evaluate the effects of various parameter variations on the near-field ULINE

model output. The results of the model sensitivity analysis are given in Figures 2-6 of the Najarian report in Appendix 51. The sensitivity analysis showed that variations in groundwater discharge will change the dilution ratios. A groundwater discharge rate of 1,176 m³/d (Section 4.2.3) gives a dilution ratio of 159. A higher discharge rate of 2,160 m³/d, for example, will decrease the near-field ratio to 88. The length of the groundwater discharge in the Bush River affects the dilution ratio. The length of the discharge used in the model was assumed to be 448 m (entire length of Beach Point Peninsular). However, a more conservative estimate of approximately half the length of the peninsular (i.e., 224 m) gives a near-field dilution ratio of 81; a length of approximately one third gives a factor of 54. The sensitivity analysis also showed that dilution ratios are somewhat dependent upon the assumed discharge depth, are sensitive to ambient current speed, and relatively insensitive to salinity.

The total near-field dilutions (near-field dilution corrected for the influence of far-field dilution) for MDE's acute toxicity criteria under Spring and Fall conditions were 42 and 43, respectively. The total near-field dilutions for chronic toxicity criteria were 168 and 185, respectively, under Spring and Fall conditions (Appendix 51). (As discussed in Section 5.2.1 above, no acute or chronic toxicity occurred in any of the biomonitoring systems at a dilution of 10). Using the above total near-field dilution factors, near-field concentrations can be estimated for toxicants of interest. The estimate is obtained by dividing the groundwater contaminant concentration by the dilution factor. For example, using the Spring dilution factor of 42, a pollutant concentration of 42 µg/L would be diluted to 1 µg/L within the near-field mixing zone limit of the Beach Point discharge in the Bush River. Using this relationship, the near-field concentrations for all the groundwater contaminants can be calculated.

The computed acute and chronic dilutions for the highest concentration of each heavy metal and aliphatic organic measured in well CC-33B (Table 3) are given in Table 5. The computed dilutions for both the acute and chronic values were determined using the Spring acute and chronic dilution factors (i.e., 42 and 168). An examination of the results in Table 5 show that the expected near-field impacts would be near or below the analytical method detection limit for most of the contaminants. For the heavy metals and organics that would be discharged in the Beach Point groundwater under the worst-case assumptions of this hazard assessment, the analysis suggests that on-site contamination will not produce any discernable adverse effect on the water quality of the Bush River.

TABLE 5. COMPUTED DILUTIONS OF THE GROUNDWATER METALS AND ORGANICS IN THE BUSH RIVER^a

| Contaminant | Maximum Groundwater Concentration | <u>Bush River Concentration</u> | |
|---------------------------|---|---------------------------------|----------------------|
| | | Acute ^b | Chronic ^c |
| <u>Heavy Metals</u> | | | |
| Aluminum | 1,500 | 35.71 | 8.93 |
| Arsenic | 7 | 0.17 | 0.04 |
| Boron | 70 | 1.67 | 0.42 |
| Cadmium | 1.4 | 0.03 | 0.01 |
| Cobalt | 110 | 0.19 | 0.05 |
| Copper | 8 | 0.19 | 0.05 |
| Nickel | 180 | 4.29 | 1.07 |
| Zinc | 310 | 7.38 | 1.85 |
| <u>Volatile Organics</u> | | | |
| Chlorobenzene | 2 | 0.05 | 0.01 |
| Chloroform | 12 | 0.29 | 0.01 |
| 1,1,2-Trichloroethane | 110 | 2.62 | 0.65 |
| 1,1,1,2-Tetrachloroethane | 3 | 0.07 | 0.02 |
| 1,1,2,2-Tetrachloroethane | 17,000 | 404.76 | 101.19 |
| 1,1-Dichloroethene | 1 | 0.02 | 0.006 |
| cis-1,2-Dichloroethene | 640 | 15.24 | 3.81 |
| Trichloroethene | 1,600 | 38.10 | 9.52 |
| Tetrachloroethene | 90 | 2.14 | 0.54 |
| Vinyl chloride | 2 | 0.05 | 0.01 |

^a All heavy metal and volatile organic concentrations are given as $\mu\text{g/L}$.

^b Based on the maximum concentration measured during the four bimonthly chemical analyses and dilution ratio of 42:1 (see Section 5.2.2).

^c Based on the maximum concentration measured during the four bimonthly chemical analyses and dilution ratio of 168:1 (see Section 5.2.2).

5.2.3 Mixing Zone Consideration

5.2.3.1 Mixing Zone Regulations

The State of Maryland allows for a mixing zone on a case-by-case basis. The State regulations regarding mixing zones are discussed in detail in the report by Najarian Associates, Inc. (1994). Briefly, as discussed in the Najarian report, a mixing

zone is defined by the State of Maryland (COMAR 26.08.01.01B) as "an area contiguous to a discharge where surface water quality or groundwater quality does not have to meet: (a) all water quality criteria, or (b) all requirements otherwise applicable to the natural water." That is, pollutant concentrations within the mixing zone may exceed the specified water quality standards within a localized vicinity of the outfall. The dimensions of the zone are normally comparable to those associated with the initial dilution process (e.g., 10-100 m). Thus, aquatic organisms would only be exposed to concentrations exceeding the specified criteria for a brief period during a transit of the mixing zone. COMAR 26.08.02.05A further states that mixing zones may be allowed only if they meet a number of requirements which ensure that no interference to aquatic ecosystem uses occurs outside the mixing zone (the list of requirements are given in Najarian Associates, Inc., 1994).

Four types of numerical toxic substance criteria are promulgated by the MDE: 1) human health criteria through ingestion of public water supplies; 2) the wholesomeness of fish for human consumption; 3) fresh, estuarine, and salt water aquatic life criteria from acute toxicity impacts; and 4) fresh, estuarine, and salt water aquatic life criteria from chronic toxicity impacts. For the purpose of applying numerical toxic substance criteria, the Bush River Area (Sub-Basin 02-13-07) is classified by MDE as "freshwater" (COMAR 26.08.02.03-1B) and as "Use II" (shellfish harvesting waters). The toxic substances criteria for both ambient surface waters ("freshwater") and human health ("fish consumption") for the pollutants in well CC-33B at Beach Point are given in Table 6. Toxic criteria related to drinking water do not apply to shellfish harvesting waters.

The following regulations (COMAR 26.08.02.05D) pertain to the application of toxic substance chronic criteria for the protection of aquatic life. First, in terms of size, the Regulatory Mixing Zone (i.e., the mixing zone in which the chronic criteria are applied) may not exceed 10 percent of the cross-sectional area of the receiving waters. Also, the chronic criteria are applied under flow conditions determined from site-specific data for the mean tide level, average tidal velocity, and, when appropriate, the design stream flow. Based on MDE's discretion, a plume dilution study may also be required for the application of these criteria. In accordance with COMAR 26.08.02.06(a), conditions pertinent to the application of toxic substance chronic criteria are also relevant to the application of toxic substance human health criteria.

The following regulations (COMAR 26.08.02.05C) pertain to the application of toxic substance acute criteria for the protection of aquatic life. Such acute criteria (for low velocity dischargers) must be met within a short distance from the outfall using the most restrictive of the following:

TABLE 6. APPLICABLE TOXIC SUBSTANCES CRITERIA FOR THE BUSH RIVER AND A COMPARISON OF PROJECTED NEAR-FIELD GROUNDWATER CONCENTRATIONS^a

| Constituent | Acute Conditions | | Chronic Conditions | |
|---------------------------|-----------------------|------------------|-----------------------|--|
| | Aquatic Life Criteria | Near-Field Conc. | Aquatic Life Criteria | Near-Field Conc. Human Health Fish Consumption |
| <u>Heavy Metals</u> | | | | |
| Aluminum | b | 35.71 | b | 8.93 c |
| Arsenic (III) | 360 ^d | 0.17 (tot) | 190 ^d | 0.04 (tot) c |
| Cadmium | 3.9 ^d | 0.03 | 1.1 ^d | 0.01 c |
| Copper | 18 ^d | 0.19 | 12 ^d | 0.05 c |
| Nickel | 1,400 ^d | 4.29 | 160 ^d | 1.07 100 |
| Zinc | 120 ^d | 7.38 | 110 ^d | 1.85 c |
| <u>Volatile Organics</u> | | | | |
| Chlorobenzene | e | 0.05 | e | 0.01 c |
| Chloroform | 28,000 ^f | 0.29 | 1,240 ^f | 0.01 c |
| 1,1,2-Trichloroethane | e | 2.62 | 9,400 ^f | 0.65 c |
| 1,1,1,2-Tetrachloroethane | e | 0.07 | e | 0.02 c |
| 1,1,2,2-Tetrachloroethane | e | 404.76 | 2,400 ^f | 101.19 c |
| 1,1-Dichloroethene | e | 0.02 | e | 0.006 c |
| cis-1,2-Dichloroethene | e | 15.24 | e | 3.81 c |
| Trichloroethene | e | 38.10 | e | 9.52 807 |
| Tetrachloroethene | e | 2.14 | e | 0.54 c |
| Vinyl chloride | e | 0.05 | e | 0.01 c |

^a All heavy metal and volatile organic concentrations are given as $\mu\text{g/L}$.

^a EPA acute and chronic aquatic life criteria are pH dependent; State of Maryland has no standard for aluminum.

^b No State of Maryland criteria.

^c Acute and chronic aquatic life criteria are hardness dependent (100 mg/L CaCO_3 used).

^d No State of Maryland or EPA criteria.

^e No State of Maryland or EPA criteria. Value presented is EPA's LOEC.

- 1) Within 10 percent of the distance (in any spatial direction) from the discharge to the edge of the Regulatory Mixing Zone used for the application of the chronic criteria.
- 2) Within a distance of 50 times at the "discharge length scale" (i.e., 50 times the square root of the cross-sectional area of the discharge outlet).
- 3) Within a distance of 5 times the local water depth in any horizontal direction from the discharge outlet.

Thus, the region where the acute criteria are applied, sometimes called the "Toxic Dilution Zone", extends outward from the discharge to a distance no larger than any of the three lengths cited above. Moreover, the Toxic Dilution Zone must not occupy more than 5 percent of the cross-sectional area of the receiving waters. Finally, the acute criteria are applied under flow conditions determined from site-specific data for the mean low water elevation, the minimum daily averaged 1-h tidal velocity and, when appropriate, design stream flow conditions. Such tidal stages are deemed "critical" (minimum dilution) conditions.

5.2.3.2 Site Applicability

The regulatory conditions relevant to mixing zones were developed for outfall diffusers. Their application to an area-wide groundwater discharge is somewhat uncertain due to the unknown extent of the area. It seems likely that the discharge area is confined within the Regulatory Mixing Zone, which for Beach Point, extends approximately 150 m (\approx 500 ft) from the river bank. However, the extent of the Toxic Dilution Zone is more poorly defined because it partially depends on the configuration of the groundwater discharge. Thus, the applicability of the mixing zone concept must be further examined.

There are three approaches used for the assessment of toxic substance discharges in the United States (U.S. EPA, 1993b). The concept of a mixing zone provides an intermediate approach to the assessment of toxic pollutant impacts. The two other approaches generally in use are "end of pipe" and "fully mixed". The "end of pipe" approach allows for zero dilution; aquatic life criteria are applied directly to the discharge water quality. This simplified approach yields the most stringent results which, in turn, most ensures the protection of aquatic biota. However, as Najarian Associates, Inc. (1994) points out, it may lack a physical or biological basis in terms of receiving water impacts. Thus, its results are not considered to be defensible. It is effectively a non-technically based policy decision.

Conversely, the "fully mixed" (or far-field) approach considers the total volume of a receiving water segment to be

instantly available for the dilution of a discharge. This approach is generally far less stringent than the "end of pipe" approach. However, it may not adequately address biological impacts in the receiving water due to near-field conditions. The concept of mixing zones evolved to address these limitations. Mixing zones allow consideration of the initial dilution process. This approach also protects aquatic life by limiting the potential for exposure to elevated concentrations to minor portions of the waterway. Since only a portion of the available dilution is considered in this approach, mixing zones provide a much more stringent basis for regulation than the "fully mixed" approach.

In the application of the mixing zone concept to the Bush River, it was necessary to conceptualize the area-wide groundwater discharge from Beach Point as a line source (diffuser). Because of this approach, potential impacts could be assessed in a technically defensible manner. However, the approach had a side effect of concentrating the groundwater impact into a smaller area. That is, the approach minimized the available near-field dilution and projected higher receiving water concentrations than would actually occur. Thus, the mixing zone analysis produced an overly conservative result. In the near-field, actual receiving water concentrations would be less than those projected due to the more dispersed nature of the groundwater plume. In the near-field, projected and actual receiving water concentrations should be comparable. However, far-field concentrations would always be less than the near-field projections.

5.2.3.3 Conformance with Aquatic Life Criteria

The anticipated near-field receiving water concentrations are compared in Table 6 with the water quality criteria for the contaminants in well CC-33B. As can be seen in Table 6, no exceedences of the criteria are projected. It is important to point out that no exceedences of the criteria are projected for groundwater discharge pollutants which were assumed to have the highest concentrations measured in well CC-33B. The assumption that the groundwater discharge would have concentrations as high as those in well CC-33B is very conservative.

There would be detectable concentrations of some of the chlorinated organics in the receiving stream if the contaminant distribution assumptions in Section 4.2.4 are used in the near-field projection. With the exception of trichloroethene for human health fish consumption, the State of Maryland has not implemented criteria for the remaining organics present in the groundwater.

5.2.3.4 Re-evaluation of Hazard Assessment Assumptions as an Alternative to a Mixing Zone

It is argued above that the State of Maryland could consider a mixing zone for the discharge of Beach Point groundwater to the Bush River. The argument for a mixing zone, however, is based on the assumption that rather large volumes of groundwater with high concentrations of contaminants may enter the Bush River. As discussed in Sections 4.2.3 and 4.2.4, the volume of groundwater and the concentrations of contaminants in the groundwater that were assumed to occur in the near-field mixing zone were based on a number of very conservative assumptions. It is highly unlikely that the contaminant loads assumed in the near-field dilution model will ever occur in the Bush River. Thus, the assumptions that were used in the dilution model should be evaluated further before a decision is made to request a mixing zone. A refinement of the assumptions would reduce the volume of the discharge as well as the concentration of contaminants that may enter the receiving stream. Finally, the State may not consider a mixing zone for a groundwater discharge as a policy to reduce contaminant loading in the Bush River.

A better estimate of the discharge rate of the surficial aquifer is needed. Several of the conservative assumptions discussed in Section 4.2.3 concerning the discharge rate could be re-evaluated to obtain better estimates that could be used in the discharge rate calculation. The following could be examined: 1) Is the horizontal transport value V_{hor} of 0.27 m/d (0.89 ft/d), which was taken from one 25-h study in an intermediate depth well (CC-33B.1), representative of the entire aquifer? 2) Is the aquifer homogeneous? 3) Does isotropic flow occur throughout the aquifer? 4) Is the average saturated thickness of the aquifer 18.8 m (61.8 ft)? 5) Does the aquifer discharge over the entire length of Beach Point Peninsula fronting the Bush River?

The following assumptions concerning the concentrations of contaminants in the groundwater (Section 4.2.4) may be considered: 1) What is the extent of heavy metal retardation via adsorption onto solid surfaces or trapping by clays through ion exchange? 2) How important is the precipitation of metals when the pH shifts from 4 to neutrality as the groundwater moves through the sediments into the receiving stream? 3) How are the DNAPL concentrations changed as abiotic (chemical) and biotic (microbial) transformations occur? 4) The assumption was made that the heavy metals and residual DNAPLs were homogeneously mixed throughout the complete aquifer. The concentrations of the materials in the shallow and intermediate wells clearly show that this is not the case. Thus, better estimates of the distribution (concentration) of the materials in the aquifer are needed. This is particularly important for vertical flux estimates of the DNAPLs.

SECTION 6

CONCLUSIONS

6.1 Biomonitoring Evaluation

Several EPA priority pollutant heavy metals were found in the groundwater. The concentrations of cadmium, copper, nickel, and zinc exceeded one or more of EPA's numerical water quality criteria for the specific metal. A number of chlorinated aliphatic compounds were also found in the groundwater, several of which are EPA priority pollutants. None of the priority pollutant organics found in the groundwater have EPA numerical water quality criteria values because insufficient data exist to develop criteria. 1,1,2,2-Tetrachlorethane concentrations found in the groundwater exceeded EPA's freshwater chronic LOEC value. None of the organics found in the groundwater have octanol water partition coefficients ($\log K_{ow}$ or $\log P$) >3 ; thus, bioaccumulation is not a potential toxicological problem.

Toxicity was detected at various groundwater concentrations by all biomonitoring systems at pH 4 with the exception of two assays. The Ames assay for mutagenicity was negative in all cases (1, 10, and 100% groundwater by volume). Japanese medaka growth was not affected by 9 months of exposure to 1 and 10% groundwater by volume in the chronic histopathology assay. No significant lesions were found in the Japanese medaka exposed to groundwater concentrations up to 10% groundwater by volume. A positive response was found for 100% groundwater in the SCE assay when the groundwater sample was concentrated $\approx 50,000X$. The SCE assay was negative for unconcentrated 100% groundwater. The positive SCE response in the 50,000X concentrated sample was judged not to be important to aquatic organisms in the vicinity of the groundwater discharge.

The lowest concentration of groundwater that caused no observable adverse effect at pH 4, in the test systems in which the NOEC value could be determined, was 10% groundwater by volume. A NOEC of 10% groundwater by volume occurred in 3 out of 4 tests in both the cladoceran and FETAX assays; it occurred once in the fathead minnow test. The NOEC concentration was higher at pH 7 in both the fathead minnow and FETAX assays. The 10% groundwater by volume NOEC for cladoceran at pH 4, however, did not change when the organism was exposed to buffered groundwater at pH 7.

The data for the alga, invertebrate, and fish in the biomonitoring study suggest that the groundwater would not be harmful at a concentration of 10% groundwater by volume. Likewise, no genotoxicity, developmental toxicity, or chronic histopathology occurred at 10% groundwater by volume. Thus, the

biomonitoring results suggest that chronic toxicity may not occur in the receiving stream at a dilution of 10:1. Chronic toxicity would be predicted to occur in the Bush River if the groundwater entered the receiving stream at the concentrations which occur in well CC-33B.

6.2 Hazard Assessment Evaluation

The groundwater may be considered a potentially hazardous material to the benthic biota of the Bush River when conservative groundwater assumptions are used (e.g., homogeneously mixed heavy metals and chlorinated organics, no chemical or biotic transformations of the DNAPLs, large release volumes through the sediments to the Bush River, etc.). The hazard to water column aquatic biota would rapidly dissipate as the groundwater materials are mixed in the receiving stream. Because the potential water quality impacts were judged to be minimal, a mixing zone approach by the State of Maryland which allows for local exceedences of water quality standards may be pursued.

The near-field (ULINE model) and far-field (dye-tracer model) screening level dilution models suggested that a total near-field dilution of approximately 42:1 for the application of Maryland's acute aquatic life criteria and a near-field dilution level of 168:1 for the application of chronic criteria would occur in the Bush River. Thus, contaminants introduced via Beach Point groundwater into Bush River receiving waters at a concentration of 42 $\mu\text{g/L}$ would be diluted locally to a concentration of approximately 1 $\mu\text{g/L}$ or less. When the dilution factors were applied to groundwater quality at Beach Point, none of the heavy metals or chlorinated aliphatic compounds exceeded Maryland's current acute or chronic aquatic life criteria. The dilution study showed that detectable concentrations of some of the chlorinated organics would occur in the receiving stream when the conservative assumptions concerning the groundwater contaminants were used in the model.

Although an argument can be made for a mixing zone, it is highly unlikely that the contaminant loads assumed in the near-field dilution model will ever occur in the Bush River. The conservative assumptions that were used in the hazard assessment should be evaluated further before a decision is made that the groundwater entering the Bush River is an environmental hazard and, thus, a mixing zone should be considered. A refinement of the assumptions would reduce the uncertainty regarding the volume of the discharge as well as the concentration of contaminants that may enter the receiving stream.

SECTION 7

REFERENCES

- Ames, B.N., W.E. Durston, E. Yamasaki, and F.D. Lee. 1973. Carcinogens are mutagens: A single test system combining liver homogenate for activation and bacteria for detection. *Proc. Natl. Acad. Sci. (USA)* 70:2281-2286.
- APHA et al. (American Public Health Association, American Water Works Association, and Water Pollution Control Federation). 1989. Standard methods for the examination of water and wastewater, 17th ed. Amer. Public Hlth. Assoc., Washington, DC.
- ASTM (American Society for Testing and Materials). 1992a. Standard guide for conducting the frog embryo teratogenesis assay-Xenopus (FETAX). ASTM Designation E 1439-91. Pages 1199-1209 in: 1992 Annual book of ASTM standards, Vol. 11.04. Amer. Soc. Testing Materials, Philadelphia.
- ASTM (American Society for Testing and Materials). 1992b. Standard guide for assessing the hazard of a material to aquatic organisms and their uses. ASTM Designation E 1023-84 (Reapproved 1988). Pages 669-684 in: 1992 Annual book of ASTM standards, Vol. 11.04. Amer. Soc. Testing Materials, Philadelphia.
- Baker, J.P., D.P. Bernard, S.W. Christensen, M.J. Safe, J. Freda, K.J. Heltcher, D.R. Marmorek, L. Rowe, P.F. Scanlon, G.W. Suter II, W.J. Warren-Hicks, and P.M. Welbourn. 1990. Biological effects of changes in surface water acid-base chemistry. NAPAP Report 13. in: Acidic deposition: State of science and technology, Vol. II Aquatic processes and effects. National Acid Precipitation Assessment Program, U.S. Environmental Protection Agency, Washington, DC.
- Bantle, J.A. 1994. Personal communication. Oklahoma State Univ., Stillwater, OK.
- Bantle, J.A., J.N. Dumont, R.A. Finch, and G. Linder. 1991. Atlas of abnormalities a guide for the performance of FETAX. Oklahoma State Univ., Stillwater, OK.
- Barnhouse, L.W., G.W. Suter, S.M. Bartell, J.J. Beauchamp, R.H. Gardner, R.V. O'Neill, and A.E. Rosen. 1986. Users manual for ecological risk assessment. ORNL-6251. Oak Ridge National Lab, Oak Ridge, TN.

- Bunton, T.E. 1990. Hepatopathology of diethylnitrosamine in the medaka (Oryzias latipes) following short-term exposure. *Toxicol. Pathol.* 18:313-323.
- Cairns, J., Jr. (ed.). 1986. Community toxicity testing. ASTM STP 920. Amer. Soc. Testing Materials, Philadelphia, PA.
- Cairns, J., Jr. 1990. The genesis of biomonitoring in aquatic ecosystems. *Environ. Professional* 12:169-176.
- Carter, H.H. 1976. Simple one dimensional kinematic model results for the Bush River and Romney Creek. Spec. Rep. 49. Johns Hopkins Univ. Chesapeake Bay Inst., Baltimore.
- Cohen, R.M. and J.W. Mercer. 1993. DNAPL site evaluation. CRC Press, Boca Raton, FL.
- Dawson, D.A., C.A. McCormick, and J.A. Bantle. 1985. Detection of teratogenic substances in acidic mine water samples using the frog embryo teratogenesis assay-Xenopus (FETAX). *J. Appl. Toxicol.* 5:234-244.
- de March, B.G.E. 1988. Acute toxicity of binary mixtures of five cations (Cu^{2+} , Cd^{2+} , Zn^{2+} , Mg^{2+} , and K^{+}) to the freshwater amphipod Gammarus lacustris (Sars): Alternative description models. *Can. J. Fish. Aquat. Sci.* 45:625-633.
- Dickson, K.L., A.W. Maki, and J.Cairns, Jr. (eds.). 1979. Analyzing the hazard evaluation process. Amer. Fish. Soc., Bethesda, MD.
- Dutka, B.J. and K.K. Kwan. 1988. Battery of screening tests approach applied to sediment extracts. *Toxicol. Assessment* 3:303-314.
- Experimental Pathology Laboratories, Inc. 1994. U.S. Army Biomedical Research and Development Laboratory test 410-001R Beach Point groundwater study with medaka, Vol. 1-3. EPL Project No. 406-027. Exp. Pathology Lab., Inc., Herndon, VA.
- Faust, M., R. Altenburger, W. Boedeker, and L.H. Grimme. 1994. Algal toxicity of binary combinations of pesticides. *Bull. Environ. Contam. Toxicol.* 53:134-141.
- Fisher, H.B., E.J. List, R.C.Y. Koh, J. Imberger, and N.H. Brooks. 1979. Mixing in inland and coastal waters. Academic Press, New York.
- Goodman, L.R. 1986. Proposed new standard guide for conducting early life-stage toxicity tests with fishes. Draft No. 10. Amer. Soc. Testing Materials, Philadelphia.

- Gulley, D.H., A.M. Boelter, and H.L. Bergman. 1989. Toxstat release 3.0. Univ. Wyoming, Laramie, WY.
- Hawkins, W.E. 1991. Development of carcinogenesis bioassay models: Response of small fish species to various classes of carcinogens. Contract No. DAMD17-88-C-8050. Gulf Coast Res. Lab., Ocean Springs, MS.
- Hawkins, W.E. 1994. Personal communication. Gulf Coast Res. Lab., Ocean Springs, MS.
- Hawkins, W.E., R.M. Overstreet, and W.W. Walker. 1988. Small fish models for identifying carcinogens in the aqueous environment. Water Resour. Bull. 24:941-949.
- Herricks, E.E. and D.J. Schaffer. 1985. Can we optimize biomonitoring? Environ. Management 9:487-492.
- Herriott, R.S. and D.T. Burton. 1992. U.S. Army Biomedical Research and Development Laboratory aquatic biomonitoring trailer version 1.0: Operations manual. AD A260733. Defense Tech. Information Center, Alexander, VA.
- Hinton, D.E., J.A. Hampton, and P.A. McCuskey. 1985. Japanese medaka liver tumor model: Review of literature and new findings. Pages 439-450 in: Jolley, R.L, R.J. Bull, W.P. Davis, S. Kata, M.H. Roberts, Jr., and V.A. Jacobs, eds. Water chlorination chemistry, environmental impact and health effects, Vol. 5. Lewis Publ., Inc., Ann Arbor, MI.
- Hinton, D.E., J.A. Couch, S.J. Teh, and L.A. Courtney. 1988. Cytological changes during progression of neoplasia in selected fish species. Aquatic Toxicol. 11:77-112.
- Hoffman, G.R. 1991. Genetic toxicology. Pages 201-225 in: Amdur, M.O., J. Doull, and C.D. Klaassen, eds. Casarett and Doull's toxicology the basic science of poisons, 4th ed. Pergamon Press, New York.
- Horning, W.B. and C.I. Weber (eds.). 1985. Short-term methods for estimating the chronic toxicity of effluents and receiving waters to freshwater organisms. EPA/600/4-85/014. U.S. Environmental Protection Agency, Environmental Monitoring and Support Laboratory, Cincinnati, OH.
- Howard, P.H. (ed.). 1989. Handbook of environmental fate and exposure data for organic chemicals, Vol. 1 Large production and priority pollutants. Lewis Publ., Inc., Chelsea, MI.
- Howard, P.H. (ed.). 1989. Handbook of environmental fate and exposure data for organic chemicals, Vol. 2 Solvents. Lewis Publ., Inc., Chelsea, MI.

- Hughes, J.S., M.M. Alexander, and K. Balu. 1988. An evaluation of appropriate expressions of toxicity in aquatic plant bioassays as demonstrated by the effects of atrazine on algae and duckweed. Pages 531-547 in: Adams, W.J., G.A. Chapman, and W.G. Landis, eds. Aquatic toxicology and environmental fate, Vol. 10. ASTM STP 971. Amer. Soc. Testing Materials, Philadelphia.
- Ishikawa, T. and S. Takayama. 1979. Importance of hepatic neoplasms in lower vertebrate animals as a tool in cancer research. J. Toxicol. Environ. Hlth. 5:537-550.
- JEG (Jacobs Engineering Group, Inc.). 1993. Beach Point test site Aberdeen Proving Ground - Edgewood Area, Maryland focused feasibility study final project work plan. Master Agreement No. 071914-A-D7, Task Order No. 142133, Supplement No. 8. Jacobs Engineering Group, Inc., Washington, DC.
- JEG (Jacobs Engineering Group, Inc.). 1994. Estimated discharge rates in the surficial aquifer Beach Point (Peninsula) study area Aberdeen Proving Ground-Edgewood Area, Maryland. Memorandum by Scott Treherne, May 23, 1994. No document reference number. Jacobs Engineering Group, Inc., Washington, DC.
- Kaiser, K.L.E. and J.M. Ribo. 1988. Photobacterium phosphoreum toxicity bioassay. II. Toxicity data compilation. Toxicity Assessment: An Internat. J. 3:195-237.
- Klaunig, J.E., B.A. Barut, and P.J. Goldblatt. 1984. Preliminary studies on the usefulness of medaka, Oryzias latipes, embryos in carcinogenicity testing. Natl. Cancer Inst. Monogr. 65:155-161.
- Knox, R.C., D.A. Sabatini, and L.W. Canter. 1993. Subsurface transport and fate processes. Lewis Publ., Ann Arbor, MI.
- K-V Associates, Inc. 1994. Hydrogeologic study of Beach Point, Edgewood Area Aberdeen Proving Grounds, Maryland. Project No. 1016. K-V Associates, Inc., Falmouth, MA.
- Lauren, D.J., S.J. Teh, and D.E. Hinton. 1990. Cytotoxicity phase of diethylnitrosamine-induced hepatic neoplasia in medaka. Cancer Res. 50:5504-5514.
- Lee, G.F. 1973. Chemical aspects of bioassay techniques for establishing water quality criteria. Water Res. 7:1525-1546.
- Lorah, M.M. 1994. Personal communication. U.S. Geological Survey, Towson, MD.

- Lorah, M.M. and J.S. Clark. 1992. Contamination of ground water, surface water, and soil and evaluation of selected pumpage scenarios in the Canal Creek Area of Aberdeen Proving Ground, Maryland. Draft Open File Report 92-. U.S. Geological Survey, Towson, MD.
- Lorah, M.M. and D.A. Vroblesky. 1989. Inorganic and organic ground-water chemistry in the Canal Creek Area of Aberdeen Proving Ground, Maryland. USGS Water-Resources Investigations Report 89-4002. U.S. Geological Survey, Towson, MD.
- Mackay, D.W., P.J. Holmes, and C.J. Redshaw. 1989. The application of bioassay techniques to water pollution problems - The United Kingdom experience. *Hydrobiologia* 188/189:77-86.
- Mackay, D., W.Y. Shiu, and K.C. Ma. 1993. Illustrated handbook of physical-chemical properties and environmental fate for organic chemicals, Vol. III Volatile organic chemicals. Lewis Publ., Inc., Ann Arbor, MI.
- Marking, L.L. 1985. Toxicity of chemical mixtures. Pages 164-176 in: Rand, G.M. and S.R. Petrocelli, eds. *Fundamentals of aquatic toxicology methods and applications*. Hemisphere Publ. Corp., New York.
- McGinnis, L.D., C.R. Daudt, M.D. Thompson, S.F. Miller, W.A. Mandell, and J. Wrobel. 1994. Environmental geophysics at Beach Point Aberdeen Proving Ground, Maryland. No document reference number. Argonne National Laboratory, Argonne, IL.
- Metcalfe, C.D. 1989. Tests for predicting carcinogenicity in fish. *Rev. Aquatic Sci.* 1:111-129.
- Microtox®. 1988. Microtox® Model 500 Toxicity Test System Manual. Microbics Corp., Carlsbad, CA.
- Miller, W.E., J.C. Greene, and T. Shiroyama. 1978. The Selenastrum capricornutum Printz algal assay bottle test. Experimental design, application and data interpretation protocol. EPA-600/9-78-018. U.S. Environmental Protection Agency, Environmental Research Laboratory, Corvallis, OR.
- Muellenhoff, W.P., A.M. Soldate, D.J. Baumgartner, M.D. Schuldt, L.R. Davis, and W.E. Frick. 1985. Initial mixing characteristics of municipal ocean discharges: Vol. 2 Procedures and applications. EPA 600/3-85-073a. U.S. Environmental Protection Agency, Environmental Research Laboratory, Narragansett, RI.

- Najarian Associates, Inc. 1994. Screening-level assessment of the near-field dilution of groundwater discharges from the Beach Point Site. Job 1437. Najarian Assoc., Inc., Eatontown, NJ.
- National Research Council. 1981. Testing for effects of chemicals on ecosystems. National Academy Press, Washington, DC.
- Nemath, G. 1989. RCRA facility assessment report Edgewood Area Aberdeen Proving Ground, Maryland. Rep. No. 39-26-0490-90. U.S. Army Environ. Hygiene Agency, Aberdeen Proving Ground, MD.
- Neuhold, J.M. 1986. Toward a meaningful interaction between ecology and aquatic toxicology. Pages 11-21 in: Poston, T.M. and R. Purdy, eds. Aquatic toxicology and environmental fate, Vol. 10. ASTM STP 921. Amer. Soc. Testing Materials, Philadelphia.
- Oliveros, J.P. and D.A. Vroblesky. 1989. Hydrogeology of the Canal Creek Area, Aberdeen Proving Ground, Maryland. USGS Water-Resources Investigations Report 89-4021. U.S. Geological Survey, Towson, MD.
- Peltier, W.H. and C.I. Weber. 1985. Methods for measuring the acute toxicity of effluents to freshwater and marine organisms, 3rd ed. EPA/600/4-85/013. U.S. Environmental Protection Agency, Environmental Monitoring and Support Laboratory, Cincinnati, OH.
- Perry, P.E. and H.J. Evans. 1975. Cytological detection of mutagens/carcinogens exposure by sister chromatid exchange. Nature 258:121-125.
- Pritchard, D.W. 1976. Distribution of contaminants and excess heat in the Bush River and Romney Creek from the proposed Perryman Electric Power Plant. Spec. Rep. 58. Johns Hopkins Univ. Chesapeake Bay Inst., Baltimore.
- Potts, K.W. 1994. Personal communication. U.S. Environmental Protection Agency, Office of Water, Heath and Ecological Criteria, Water Quality Criteria, Washington, DC.
- Qureshi, A.A., K.W. Flood, S.R. Thompson, S.M. Janhurst, C.S. Inniss, and D.A. Rokosh. 1982. Comparison of a luminescent bacterial test with other bioassays for determining toxicity of pure compounds and complex effluents. Pages 179-195 in: Pearson, J.G., R.B. Foster, and W.E. Bishop, eds. Aquatic toxicology and hazard assessment, Vol. 5. ASTM STP 776. Amer. Soc. Testing Materials, Philadelphia.

- Rand, G.M. and S.R. Petrocelli (eds.). 1985. Fundamentals of aquatic toxicology methods and applications. Hemisphere Publ. Corp., New York.
- SAS. 1987. SAS/STAT™ guide for personal computer, 6th ed. SAS Inst., Inc., Cary, NC.
- Schaeffer, D.J. and K.G. Janardan. 1987. Designing batteries of short-term tests with largest inter-tier correlation. *Ecotoxicol. Environ. Safety* 13:316-323.
- Sigma Stat. 1992. Sigma Stat statistical software for working scientists, version 1.01. Jandell Scientific, Inc., San Rafael, CA.
- Sprague, J.B. 1985. Factors that modify toxicity. Pages 123-163 in: Rand, G.M. and S.R. Petrocelli, eds. Fundamentals of aquatic toxicology methods and applications. Hemisphere Publ. Corp., New York.
- Stephen, C.E. 1978. EPA statistics program, version 1.4. U.S. Environmental Protection Agency, Environmental Research Laboratory, Duluth, MN.
- Suter, G.W., II. 1993a. Organism-level effects. Pages 175-246 in: Suter, G.W., II, ed. Ecological risk assessment. Lewis Publ., Ann Arbor, MI.
- Thurston, R.V., R.C. Russo, and K. Emerson. 1979. Aqueous ammonia equilibrium - Tabulation of percent ionized ammonia. EPA-600/3-79-091. U.S. Environmental Protection Agency, Environmental Research Laboratory, Duluth, MN.
- U.S. Biomedical Research and Development Laboratory. 1993. SOP for the determination of HMX, RDX, TNB, Tetryl, TNT, 2,6-DNT, and 2,4-DNT in water by high performance liquid chromatography (HPLC). No SOP Number. U.S. Biomedical Research and Development Laboratory, Ft. Detrick, Frederick, MD.
- U.S. EPA. 1980a. Ambient water quality criteria for nickel. EPA 440/5-80-060. U.S. Environmental Protection Agency, Washington, DC.
- U.S. EPA. 1980b. Ambient water quality criteria for chlorinated ethanes. EPA 440/5-80-029. U.S. Environmental Protection Agency, Washington, DC.
- U.S. EPA. 1984a. Ambient water quality criteria for cadmium - 1984. EPA 440/5-84-032. U.S. Environmental Protection Agency, Washington, DC.

- U.S. EPA. 1984b. Ambient water quality criteria for copper - 1984. EPA 440/5-84-031. U.S. Environmental Protection Agency, Washington, DC.
- U.S. EPA. 1985a. Toxic substances control act test guidelines; final rules. Federal Register 50:39252-39516.
- U.S. EPA. 1985b. Water quality assessment: A screening procedure for toxic and conventional pollutants. EPA/600/6-85/002b. U.S. Environmental Protection Agency, Athens, GA.
- U.S. EPA. 1986a. Revision of toxic substances control act test guidelines. Federal Register 51:1522-1543.
- U.S. EPA. 1986b. Quality criteria for water 1986. EPA/5-86-001. U.S. Environmental Protection Agency, Washington, DC.
- U.S. EPA. 1987. Ambient water quality criteria for zinc - 1987. EPA 440/5-87-003. U.S. Environmental Protection Agency, Washington, DC.
- U.S. EPA. 1991a. Technical support document for water quality-based toxics control. EPA/505/2-90-001. U.S. Environmental Protection Agency, Washington, DC.
- U.S. EPA. 1991b. Evaluation of dredged material proposed for ocean disposal. EPA-503/8-91/001. U.S. Environmental Protection Agency, Washington, DC.
- U.S. EPA. 1991c. Assessment and control of bioconcentratable contaminants in surface waters. 1991 Draft Report. U.S. Environmental Protection Agency, Washington, DC.
- U.S. EPA. 1993a. Technical basis for deriving sediment quality criteria for nonionic organic contaminants for the protection of benthic organisms by using equilibrium partitioning. EPA-822-R-93-011. U.S. Environmental Protection Agency, Washington, DC.
- U.S. EPA. 1993b. Water quality standards handbook, 2nd ed. EPA-823-B-93-002. U.S. Environmental Protection Agency, Washington, DC.
- Walker, W.W., C.S. Manning, R.R. Overstreet, and W.E. Hawkins. 1985. Development of aquarium fish models for environmental carcinogenesis: an intermittent-flow exposure system for volatile hydrophobic chemicals. J. Appl. Toxicol. 5:255-260.

Weber, C.I. (ed.). 1991. Methods for measuring the acute toxicity of effluents and receiving waters to freshwater and marine organisms, 4th ed. EPA/600/4-90/027. U.S. Environmental Protection Agency, Environmental Monitoring and Support Laboratory, Cincinnati, OH.

Weber, C.I., W.H. Peltier, T.J. Norberg-King, W.B. Horning, II, F.A. Kessler, J.R. Menkedick, T.W. Neiheisel, P.A. Lewis, D.J. Klemm, Q.H. Pickering, E.L. Robinson, J.M. Lazorchak, L.J. Wymer, and R.W. Freyberg. 1989. Short-term methods for estimating the chronic toxicity of effluents and receiving waters to freshwater organisms, 2nd ed. EPA/600/4-89/001. U.S. Environmental Protection Agency, Environmental Monitoring and Support Laboratory, Cincinnati, OH.

Weis, J.S. and P. Weis. 1989. Effects of environmental pollutants on early fish development. Rev. Aquat. Sci. 1:45-73.

Williamson, R.L., D.T. Burton, J.H. Clarke, and L.E. Fleming. 1993. Gathering danger: The urgent need to regulate substances that can bioaccumulate. Ecol. Law Quart. 20:605-720.

Winner, R.W. 1989. Multigeneration life-span tests of the nutritional adequacy of several diets and culture waters for Ceriodaphnia dubia. Environ. Toxicol. Chem. 8:513-520.

APPENDIX 1

MICROTOX® ASSAYS CONDUCTED ON RAW (pH \approx 4) BEACH POINT
GROUNDWATER (WELL CC-33B), CHRONIC HISTOPATHOLOGY
EXPOSURE TANKS, AND DILUENT WATER

TABLE A1-1. MICROTOX® TEST RESULTS ON GRAB SAMPLES OF BEACH POINT 100% GROUNDWATER (pH
 ~4) TAKEN FROM WELL CC-33B^a

| Date of Sample | Microtox® 5-Min Readings | | | Microtox® 15-Min Readings | | |
|-------------------|--------------------------|------------------------|-------|---------------------------|------------------------|-------|
| | 5-min EC50 | 95% Fiducial Limits | Slope | 15-min EC50 | 95% Fiducial Limits | Slope |
| Mar 24 | 15.7 | 12.45-19.71 | 1.07 | 19.5 | 12.35-30.81 | 1.20 |
| 26 | 11.3 | 08.67-14.69 | 0.94 | 18.0 | 11.15-28.97 | 1.00 |
| 31 | 13.3 | 11.02-16.14 | 0.98 | 21.1 | 10.47-42.55 | 1.10 |
| Apr 02 | 11.8 | 07.50-18.46 | 0.80 | 19.0 | 08.69-41.47 | 0.82 |
| 05 | 13.3 | 09.57-18.51 | 0.81 | 21.8 | 07.79-61.22 | 0.89 |
| 07 | 12.4 | 09.32-16.40 | 0.83 | 20.1 | 10.21-39.53 | 0.95 |
| 09 | 14.7 | 11.70-18.45 | 0.82 | 22.9 | 10.57-49.61 | 0.89 |
| 12 | 14.4 | 10.75-19.16 | 0.93 | 23.0 | 11.32-46.61 | 1.06 |
| 14 | 14.9 | 11.29-19.56 | 0.98 | 22.5 | 08.73-58.01 | 1.01 |
| 16 | 14.1 | 09.88-20.21 | 0.90 | 21.5 | 08.02-57.54 | 1.02 |
| 19 | 10.7 | 07.10-16.05 | 0.90 | 17.0 | 07.38-39.01 | 0.97 |
| 21 | 11.8 | 08.91-15.62 | 0.85 | 19.3 | 09.48-39.40 | 0.98 |
| 23 | 12.3 | 09.55-15.79 | 0.90 | 18.2 | 07.97-41.51 | 0.90 |
| 26 | 13.7 | 11.78-15.83 | 0.92 | 21.4 | 10.65-43.14 | 1.00 |
| 28 | 12.6 | 09.31-17.15 | 0.86 | 19.6 | 10.34-37.08 | 0.94 |
| 30 | 12.4 | 09.53-16.20 | 0.89 | 20.4 | 09.22-45.15 | 1.03 |
| May 03 | 15.1 | 10.69-21.25 | 1.01 | 24.9 | 09.18-67.57 | 1.09 |
| 05 | 15.2 | 08.18-28.25 | 0.94 | 23.6 | 06.13-90.84 | 1.13 |
| 07 | 11.0 | 08.62-14.10 | 0.79 | 17.2 | 08.64-34.06 | 0.87 |
| 10 | 12.4 | 10.73-14.39 | 0.87 | 18.8 | 10.47-33.93 | 0.89 |
| 12 | 14.9 | 14.48-15.36 | 1.00 | 20.9 | 10.75-40.66 | 1.02 |
| 14 | 10.6 | 07.55-14.81 | 0.79 | 20.5 | 15.70-26.86 | 1.02 |
| 17 | 12.9 | 12.01-13.89 | 0.88 | 20.3 | 08.79-47.08 | 0.86 |
| 19 | 13.9 | 11.33-16.97 | 0.84 | 22.1 | 10.19-48.04 | 0.92 |
| 21 | 13.5 | 11.27-16.27 | 0.88 | 20.6 | 08.67-48.76 | 0.95 |
| 24 | 12.0 | 08.01-18.06 | 0.81 | 22.2 | 10.40-47.16 | 0.93 |
| 26 | 13.6 | 12.90-14.32 | 0.96 | 20.7 | 11.36-37.63 | 1.04 |
| 28 | 11.5 | 08.94-14.91 | 0.88 | 20.4 | 10.87-38.38 | 1.04 |
| Jun 01 | 12.6 | 08.78-18.10 | 0.92 | 20.0 | 11.70-34.37 | 1.05 |
| 02 | 12.2 | 09.43-15.80 | 0.85 | 19.2 | 11.23-32.82 | 1.04 |

TABLE A1-1. (CONTINUED)

| Date of Sample | Microtox® 5-Min Readings | | | Microtox® 15-Min Readings | | |
|-------------------|--------------------------|------------------------|-------|---------------------------|------------------------|-------|
| | 5-min EC50 | 95% Fiducial Limits | Slope | 15-min EC50 | 95% Fiducial Limits | Slope |
| Jun 04 | 10.8 | 07.65-15.37 | 0.82 | | | |
| 07 | ^b | | | ^b | | |
| 09 | 14.7 | 13.43-16.11 | 1.04 | 21.4 | 11.09-41.33 | 1.12 |
| 10 | 13.6 | 11.89-15.51 | 0.99 | 20.3 | 10.57-38.86 | 0.98 |
| 15 | 13.0 | 09.71-17.43 | 0.94 | 19.7 | 09.15-42.35 | 0.99 |
| 16 | 14.1 | 12.23-16.32 | 0.95 | 20.0 | 11.25-35.65 | 1.00 |
| 18 | 16.0 | 12.97-19.63 | 0.90 | 21.8 | 11.46-41.67 | 0.86 |
| 21 | 15.7 | 14.41-17.14 | 1.05 | 23.0 | 16.04-32.87 | 1.02 |
| 23 | 14.1 | 11.31-17.54 | 1.02 | 22.4 | 12.09-41.49 | 1.03 |
| 25 | 15.4 | 10.14-23.52 | 0.95 | 22.0 | 08.32-57.97 | 0.86 |
| 28 | 17.3 | 14.52-20.64 | 0.97 | 24.2 | 09.90-59.13 | 0.88 |
| 30 | 16.8 | 14.09-20.13 | 1.05 | 25.3 | 11.26-56.62 | 0.94 |
| Jul 02 | 19.5 | 14.57-26.03 | 1.08 | 26.4 | 10.25-67.97 | 0.95 |
| 05 | ^c | | | ^c | | |
| 07 | ^c | | | ^c | | |
| 09 | ^c | | | ^c | | |
| 12 | 16.2 | 12.50-21.01 | 0.91 | 23.3 | 10.41-51.94 | 0.93 |
| 14 | 14.8 | 11.59-18.82 | 0.87 | 23.4 | 09.20-59.61 | 0.80 |
| 16 | 17.7 | 14.17-22.15 | 1.01 | 24.2 | 11.78-49.89 | 0.87 |
| 19 | 11.6 | 10.02-13.46 | 0.87 | 18.7 | 07.97-43.69 | 0.80 |
| 21 | 16.3 | 12.25-21.66 | 0.88 | 24.9 | 09.14-67.78 | 1.07 |
| 23 | 17.5 | 15.15-20.11 | 0.98 | 24.8 | 12.05-50.98 | 0.99 |
| 26 | 18.1 | 16.58-19.80 | 0.91 | 25.7 | 11.13-59.10 | 1.02 |
| 28 | ^b | | | ^b | | |
| 30 | 17.0 | 11.08-26.24 | 1.24 | 23.1 | 13.36-39.81 | 1.26 |
| Aug 02 | 16.1 | 14.61-17.77 | 0.86 | 23.8 | 12.26-46.21 | 1.00 |
| 05 | 15.6 | 13.71-17.70 | 0.79 | 22.7 | 09.53-54.16 | 0.86 |
| 06 | 17.2 | 13.86-21.33 | 0.85 | 28.8 | 15.58-53.17 | 1.21 |
| 09 | 19.6 | 17.98-21.31 | 1.02 | 30.1 | 12.93-70.19 | 0.95 |
| 11 | 20.8 | 18.16-23.89 | 0.93 | 32.4 | 11.26-93.20 | 0.82 |

TABLE A1-1. (CONTINUED)

| Date of Sample | Microtox® 5-Min Readings | | | Microtox® 15-Min Readings | | |
|-------------------|--------------------------|------------------------|-------|---------------------------|------------------------|-------|
| | 5-min EC50 | 95% Fiducial Limits | Slope | 15-min EC50 | 95% Fiducial Limits | Slope |
| Aug 12 | 20.9 | 17.73-24.57 | 0.95 | 32.0 | 10.21-100.56 | 0.86 |
| 16 | 21.0 | 14.51-30.35 | 1.02 | 33.0 | 11.24-96.77 | 0.92 |
| 18 | 17.4 | 13.18-23.04 | 0.90 | 29.6 | 21.19-41.37 | 0.93 |
| 20 | 23.7 | 15.90-35.51 | 0.83 | 32.7 | 14.11-75.79 | 0.92 |
| 24 | 23.8 | 21.14-26.89 | 1.12 | 34.6 | 15.63-76.62 | 1.16 |
| 25 | 21.4 | 17.72-25.92 | 1.07 | 33.0 | 18.00-60.43 | 1.14 |
| 27 | 24.1 | 16.02-36.29 | 0.88 | 33.7 | 13.32-85.10 | 0.92 |
| 30 | 21.0 | 16.85-26.16 | 0.94 | 28.8 | 11.34-73.26 | 1.02 |
| Sep 01 | 29.1 | 21.56-39.19 | 1.13 | 39.2 | 15.80-97.27 | 1.02 |
| 03 | 26.9 | 21.13-34.34 | 0.83 | 33.8 | 25.35-44.96 | 1.03 |
| 07 | 26.9 | 20.69-34.99 | 0.93 | 35.4 | 08.68-145.15 | 0.81 |
| 08 | 22.2 | 18.59-26.43 | 0.85 | 27.8 | 12.49-62.07 | 0.82 |
| 09 | 25.9 | 18.65-35.92 | 0.91 | 31.0 | 16.10-59.63 | 0.91 |
| 13 | 22.2 | 16.46-29.86 | 0.85 | 29.5 | 11.53-75.74 | 0.83 |
| 14 | 25.4 | 17.16-37.66 | 0.87 | 32.5 | 07.40-142.39 | 0.88 |
| 15 | 26.8 | 20.51-35.11 | 0.85 | 28.7 | 08.41-97.92 | 0.93 |
| 20 | 22.0 | 14.99-32.35 | 0.87 | 27.8 | 10.71-72.04 | 0.88 |
| 21 | 22.8 | 19.25-26.97 | 1.06 | 29.7 | 15.06-58.44 | 1.18 |
| 22 | 19.9 | 17.06-23.21 | 0.93 | 26.1 | 10.52-65.00 | 1.01 |
| 27 | 25.6 | 18.60-35.17 | 1.13 | 34.2 | 15.10-77.48 | 1.18 |
| 28 | 23.5 | 13.30-41.68 | 0.81 | 32.5 | 09.76-108.07 | 0.85 |
| 29 | 24.5 | 17.63-34.05 | 0.98 | 33.3 | 15.19-72.95 | 1.04 |
| Oct 04 | 21.7 | 15.45-30.42 | 0.94 | 32.7 | 10.06-106.37 | 0.85 |
| 06 | 23.1 | 17.77-29.96 | 0.98 | 30.8 | 11.10-85.53 | 0.85 |
| 08 | 22.1 | 21.24-23.05 | 1.04 | 29.2 | 10.61-80.54 | 1.11 |
| 11 | 22.9 | 18.61-28.24 | 0.94 | 30.6 | 10.88-86.27 | 0.95 |
| 13 | 27.1 | 17.56-41.71 | 0.92 | 36.2 | 08.15-160.86 | 0.83 |
| 15 | 21.3 | 16.92-26.91 | 0.93 | 30.4 | 08.52-108.22 | 0.88 |
| 18 | 23.0 | 15.02-35.23 | 0.99 | 33.1 | 11.59-94.23 | 1.00 |
| 19 | 19.7 | 14.70-26.52 | 0.86 | 26.7 | 09.67-73.96 | 0.84 |
| 20 | 27.1 | 19.65-37.29 | 1.16 | d | | |

TABLE A1-1. (CONTINUED)

| Date of Sample | Microtox® 5-Min Readings | | | Microtox® 15-Min Readings | | |
|-------------------|--------------------------|------------------------|-------|---------------------------|------------------------|-------|
| | 5-min EC50 | 95% Fiducial Limits | Slope | 15-min EC50 | 95% Fiducial Limits | Slope |
| Oct 26 | 22.5 | 17.55-28.92 | 0.97 | 34.1 | 22.19-52.27 | 1.07 |
| 27 | 23.7 | 19.69-28.49 | 1.02 | 34.9 | 15.18-80.33 | 1.10 |
| 29 | 22.0 | 16.02-30.24 | 0.85 | 36.7 | 23.45-57.37 | 1.15 |
| Nov 01 | 20.0 | 17.83-22.33 | 0.67 | 38.2 | 12.02-121.33 | 0.94 |
| 03 | 19.3 | 17.04-21.89 | 0.86 | 35.8 | 30.83-41.52 | 1.26 |
| 05 | 22.4 | 21.16-23.64 | 0.90 | 38.4 | 08.77-168.47 | 1.73 |
| 08 | 19.1 | 14.29-25.44 | 0.69 | 37.0 | 09.61-142.56 | 0.86 |
| 10 | 18.9 | 15.46-23.07 | 0.79 | 33.7 | 10.97-103.34 | 0.99 |
| 12 | 22.5 | 20.02-25.24 | 0.82 | 36.9 | 10.69-127.60 | 1.05 |
| 15 | 24.1 | 18.81-30.75 | 0.90 | 40.7 | 12.11-136.78 | 0.98 |
| 17 | 27.0 | 24.43-29.94 | 0.89 | 45.2 | 12.04-169.60 | 1.06 |
| 19 | 25.4 | 17.11-37.62 | 0.92 | 40.2 | 05.67-285.40 | 1.00 |
| 22 | 21.8 | 17.12-27.68 | 0.84 | 39.4 | 12.71-122.21 | 0.93 |
| 24 | 17.2 | 14.62-20.16 | 0.68 | 33.6 | 13.92-81.10 | 0.78 |
| 26 | 30.3 | 27.34-33.68 | 0.94 | 47.1 | 06.54-338.68 | 1.16 |
| Dec 01 | 20.4 | 15.79-26.47 | 0.79 | 36.8 | 16.79-80.87 | 0.90 |
| 03 | 20.8 | 20.17-21.50 | 0.90 | 39.3 | 24.17-64.04 | 1.24 |
| 06 | 25.4 | 11.39-56.51 | 1.05 | 30.2 | 07.97-114.48 | 1.09 |

- ^a EC50s and 95% fiducial limits are expressed as percent groundwater by volume.
^b Microtox® instrument malfunctioned; sent to Microbics Corporation for repair.
^c Microtox® instrument was not available for use.
^d Lost data - dropped cuvette.

TABLE A1-2. MICROTOX® TEST RESULTS ON GRAB SAMPLES OF BEACH POINT 10% GROUNDWATER TAKEN FROM THE CHRONIC HISTOPATHOLOGY STUDY TANKS^a

| Date of Sample | Microtox® 5-Min Readings | | Microtox® 15-Min Readings | |
|----------------|--------------------------|-------------------------|---------------------------|---------------------|
| | 5-min EC50 | 95% Fiducial Limits | 15-min EC50 | 95% Fiducial Limits |
| Mar 24 | - | | - | |
| Mar 26 | - | | - | |
| Mar 31 | - | | - | |
| Apr 02 | - | | - | |
| Apr 05 | - | | - | |
| Apr 07 | - | | - | |
| Apr 09 | - | | - | |
| Apr 12 | - | | - | |
| Apr 14 | - | | - | |
| Apr 16 | - | | - | |
| Apr 19 | - | | - | |
| Apr 21 | 98.6 | 00.00-1x10 ⁶ | - | |
| Apr 23 | - | | - | |
| Apr 26 | - | | - | |
| Apr 28 | - | | - | |
| Apr 30 | - | | - | |
| May 03 | - | | - | |
| May 05 | - | | - | |
| May 07 | - | | - | |
| May 10 | - | | - | |
| May 12 | - | | - | |
| May 14 | - | | - | |
| May 17 | - | | - | |
| May 19 | - | | - | |
| May 21 | - | | - | |
| May 24 | - | | - | |
| May 26 | - | | - | |
| May 28 | - | | - | |
| Jun 01 | - | | - | |

TABLE A1-2. (CONTINUED)

| Date of Sample | Microtox® 5-Min Readings | | | Microtox® 15-Min Readings | | |
|----------------|--------------------------|---------------------|-------|---------------------------|---------------------|-------|
| | 5-min EC50 | 95% Fiducial Limits | Slope | 15-min EC50 | 95% Fiducial Limits | Slope |
| Jun 02 | - | | | - | | |
| 04 | - | | | b | | |
| 07 | b | | | b | | |
| 09 | - | | | - | | |
| 10 | - | | | - | | |
| 15 | - | | | - | | |
| 16 | - | | | - | | |
| 18 | - | | | - | | |
| 21 | - | | | - | | |
| 23 | - | | | - | | |
| 25 | - | | | - | | |
| 28 | - | | | - | | |
| 30 | - | | | - | | |
| Jul 02 | - | | | - | | |
| 05 | c | | | c | | |
| 07 | c | | | c | | |
| 09 | c | | | c | | |
| 12 | - | | | - | | |
| 14 | - | | | - | | |
| 16 | - | | | - | | |
| 19 | - | | | - | | |
| 21 | - | | | - | | |
| 23 | - | | | - | | |
| 26 | - | | | - | | |
| 28 | b | | | b | | |
| 30 | - | | | - | | |
| Aug 02 | - | | | - | | |
| 05 | - | | | - | | |
| 06 | - | | | - | | |
| 09 | - | | | - | | |

TABLE A1-2. (CONTINUED)

| Date of Sample | Microtox® 5-Min Readings | | | Microtox® 15-Min Readings | | |
|-------------------|--------------------------|------------------------|-------|---------------------------|------------------------|-------|
| | 5-min EC50 | 95% Fiducial Limits | Slope | 15-min EC50 | 95% Fiducial Limits | Slope |
| Aug 11 | - | | | - | | |
| 12 | - | | | - | | |
| 16 | - | | | - | | |
| 18 | - | | | - | | |
| 20 | - | | | - | | |
| 24 | - | | | - | | |
| 25 | - | | | - | | |
| 27 | - | | | - | | |
| 30 | - | | | - | | |
| Sep 01 | - | | | - | | |
| 03 | - | | | - | | |
| 07 | - | | | - | | |
| 08 | - | | | - | | |
| 09 | - | | | - | | |
| 13 | - | | | - | | |
| 14 | - | | | - | | |
| 15 | - | | | - | | |
| 20 | - | | | - | | |
| 21 | - | | | - | | |
| 22 | - | | | - | | |
| 27 | - | | | - | | |
| 28 | - | | | - | | |
| 29 | - | | | - | | |
| Oct 04 | - | | | - | | |
| 06 | - | | | - | | |
| 08 | - | | | - | | |
| 11 | - | | | - | | |
| 13 | - | | | - | | |
| 15 | - | | | - | | |
| 18 | - | | | - | | |

TABLE A1-2. (CONTINUED)

| Date of Sample | Microtox® 5-Min Readings | | | Microtox® 15-Min Readings | | |
|-------------------|--------------------------|------------------------|-------|---------------------------|------------------------|-------|
| | 5-min EC50 | 95% Fiducial Limits | Slope | 15-min EC50 | 95% Fiducial Limits | Slope |
| 19 | - | | | - | | |
| 20 | - | | | - | | |
| 26 | - | | | - | | |
| 27 | - | | | - | | |
| 29 | - | | | - | | |
| Nov 01 | - | | | - | | |
| 03 | - | | | - | | |
| 05 | - | | | - | | |
| 08 | - | | | - | | |
| 10 | - | | | - | | |
| 12 | - | | | - | | |
| 15 | - | | | - | | |
| 17 | - | | | - | | |
| 19 | - | | | - | | |
| 22 | - | | | - | | |
| 24 | - | | | - | | |
| 26 | - | | | - | | |
| Dec 01 | - | | | - | | |
| 03 | - | | | - | | |
| 06 | - | | | - | | |

- ^a EC50s and 95% fiducial limits are expressed as percent groundwater by volume.
^b Microtox® instrument malfunctioned; sent to Microbics Corporation for repair.
^c Microtox® instrument was not available for use.

TABLE A1-3. MICROTOX® TEST RESULTS ON GRAB SAMPLES OF BEACH POINT 1% GROUNDWATER TAKEN FROM THE CHRONIC HISTOPATHOLOGY STUDY TANKS^a

| Date of Sample | Microtox® 5-Min Readings | | | Microtox® 15-Min Readings | | |
|----------------|--------------------------|---------------------|-------|---------------------------|---------------------|-------|
| | 5-min EC50 | 95% Fiducial Limits | Slope | 15-min EC50 | 95% Fiducial Limits | Slope |
| Mar 24 | - | | | - | | |
| 26 | - | | | - | | |
| 31 | - | | | - | | |
| Apr 02 | - | | | - | | |
| 05 | - | | | - | | |
| 07 | - | | | - | | |
| 09 | - | | | - | | |
| 12 | - | | | - | | |
| 14 | - | | | - | | |
| 16 | - | | | - | | |
| 19 | - | | | - | | |
| 21 | - | | | - | | |
| 23 | - | | | - | | |
| 26 | - | | | - | | |
| 28 | - | | | - | | |
| 30 | - | | | - | | |
| May 03 | - | | | - | | |
| 05 | - | | | - | | |
| 07 | - | | | - | | |
| 10 | - | | | - | | |
| 12 | - | | | - | | |
| 14 | - | | | - | | |
| 17 | - | | | - | | |
| 19 | - | | | - | | |
| 21 | - | | | - | | |
| 24 | - | | | - | | |
| 26 | - | | | - | | |
| 28 | - | | | - | | |
| Jun 01 | - | | | - | | |

TABLE A1-3. (CONTINUED)

| Date of Sample | Microtox® 5-Min Readings | | | Microtox® 15-Min Readings | | |
|-------------------|--------------------------|------------------------|-------|---------------------------|------------------------|-------|
| | 5-min EC50 | 95% Fiducial Limits | Slope | 15-min EC50 | 95% Fiducial Limits | Slope |
| Jun 02 | - | | | - | | |
| 04 | - | | | b | | |
| 07 | b | | | b | | |
| 09 | - | | | - | | |
| 10 | - | | | - | | |
| 15 | - | | | - | | |
| 16 | - | | | - | | |
| 18 | - | | | - | | |
| 21 | - | | | - | | |
| 23 | - | | | - | | |
| 25 | - | | | - | | |
| 28 | - | | | - | | |
| 30 | - | | | - | | |
| Jul 02 | - | | | - | | |
| 05 | c | | | c | | |
| 07 | c | | | c | | |
| 09 | c | | | c | | |
| 12 | - | | | - | | |
| 14 | - | | | - | | |
| 16 | - | | | - | | |
| 19 | - | | | - | | |
| 21 | - | | | - | | |
| 23 | - | | | 52.7 ^d | 07.90-351.71 | 1.49 |
| 26 | - | | | - | | |
| 28 | b | | | b | | |
| 30 | - | | | - | | |
| Aug 02 | - | | | - | | |
| 05 | - | | | - | | |
| 06 | - | | | - | | |
| 09 | - | | | - | | |

TABLE A1-3. (CONTINUED)

| Date of Sample | Microtox® 5-Min Readings | | | Microtox® 15-Min Readings | | |
|-------------------|--------------------------|------------------------|-------|---------------------------|------------------------|-------|
| | 5-min EC50 | 95% Fiducial Limits | Slope | 15-min EC50 | 95% Fiducial Limits | Slope |
| Aug 11 | - | | | - | | |
| 12 | - | | | - | | |
| 16 | - | | | - | | |
| 18 | - | | | - | | |
| 20 | - | | | - | | |
| 24 | - | | | - | | |
| 25 | - | | | - | | |
| 27 | - | | | - | | |
| 30 | - | | | - | | |
| Sep 01 | - | | | - | | |
| 03 | - | | | - | | |
| 07 | - | | | - | | |
| 08 | - | | | - | | |
| 09 | - | | | - | | |
| 13 | - | | | - | | |
| 14 | - | | | - | | |
| 15 | - | | | - | | |
| 20 | - | | | - | | |
| 21 | - | | | - | | |
| 22 | - | | | - | | |
| 27 | - | | | - | | |
| 28 | - | | | - | | |
| 29 | - | | | - | | |
| Oct 04 | - | | | - | | |
| 06 | - | | | - | | |
| 08 | - | | | - | | |
| 11 | - | | | - | | |
| 13 | - | | | - | | |
| 15 | - | | | - | | |
| 18 | - | | | - | | |

TABLE A1-3. (CONTINUED)

| Date of Sample | Microtox® 5-Min Readings | | | Microtox® 15-Min Readings | | |
|-------------------|--------------------------|------------------------|-------|---------------------------|------------------------|-------|
| | 5-min EC50 | 95% Fiducial Limits | Slope | 15-min EC50 | 95% Fiducial Limits | Slope |
| 19 | - | | | - | | |
| 20 | - | | | - | | |
| 26 | - | | | - | | |
| 27 | - | | | - | | |
| 29 | - | | | - | | |
| Nov 01 | - | | | - | | |
| 03 | - | | | - | | |
| 05 | - | | | - | | |
| 08 | - | | | - | | |
| 10 | - | | | - | | |
| 12 | - | | | - | | |
| 15 | - | | | - | | |
| 17 | - | | | - | | |
| 19 | - | | | - | | |
| 22 | - | | | - | | |
| 24 | - | | | - | | |
| 26 | - | | | - | | |
| Dec 01 | - | | | - | | |
| 03 | - | | | - | | |
| 06 | - | | | - | | |

- ^a EC50s and 95% fiducial limits are expressed as percent groundwater by volume.
^b Microtox® instrument malfunctioned; sent to Microbics Corporation for repair.
^c Microtox® instrument was not available for use.
^d Microtox® instrument malfunctioning; reading may be incorrect.

TABLE A1-4. MICROTOX® TEST RESULTS ON GRAB SAMPLES OF APG-EA DILUENT WATER^a

| Date of Sample | Microtox® 5-Min Readings | | | Microtox® 15-Min Readings | | |
|-------------------|--------------------------|------------------------|-------|---------------------------|------------------------|-------|
| | 5-min EC50 | 95% Fiducial Limits | Slope | 15-min EC50 | 95% Fiducial Limits | Slope |
| Mar 26 | - | | | - | | |
| Mar 31 | - | | | - | | |
| Apr 02 | - | | | - | | |
| Apr 05 | - | | | - | | |
| Apr 07 | - | | | - | | |
| Apr 09 | - | | | - | | |
| Apr 14 | - | | | - | | |
| Apr 21 | - | | | - | | |
| Apr 28 | - | | | - | | |
| May 07 | - | | | - | | |
| May 12 | - | | | - | | |
| May 19 | - | | | - | | |
| May 26 | - | | | - | | |
| Jun 04 | b | | | b | | |
| Jun 09 | - | | | - | | |
| Jun 16 | - | | | - | | |
| Jun 23 | - | | | - | | |
| Jun 30 | - | | | - | | |
| Jul 07 | c | | | c | | |
| Jul 14 | - | | | - | | |
| Jul 21 | - | | | - | | |
| Jul 28 | b | | | b | | |
| Aug 06 | - | | | - | | |
| Aug 11 | - | | | - | | |
| Aug 18 | - | | | - | | |
| Aug 25 | - | | | - | | |
| Sep 01 | - | | | - | | |
| Sep 07 | - | | | - | | |
| Sep 15 | - | | | - | | |

TABLE A1-4. (CONTINUED)

| Date of Sample | Microtox® 5-Min Readings | | | Microtox® 15-Min Readings | | |
|-------------------|--------------------------|------------------------|-------|---------------------------|------------------------|-------|
| | 5-min EC50 | 95% Fiducial Limits | Slope | 15-min EC50 | 95% Fiducial Limits | Slope |
| Sep 22 | - | | | - ^d | | |
| Sep 27 | - | | | - | | |
| Oct 06 | - | | | - | | |
| Oct 13 | - | | | - | | |
| Oct 20 | - | | | - | | |
| Oct 27 | - | | | - | | |
| Nov 03 | - | | | - | | |
| Nov 10 | - | | | - | | |
| Nov 19 | - | | | - | | |
| Nov 24 | - | | | - | | |
| Dec 01 | - | | | - | | |

^a EC50s and 95% fiducial limits are expressed as percent groundwater by volume.

^b Microtox® instrument malfunctioned; sent to Microbics Corporation for repair.

^c Microtox® instrument was not available for use.

^d Electrical power surge caused an interrupt in the circuit; therefore, the 15-min readings could not be obtained.

APPENDIX 2

MICROTOX® ASSAYS CONDUCTED ON BEACH POINT
BUFFERED (pH \approx 7) GROUNDWATER (WELL CC-33B)
AND pH OF SAMPLES USED IN ASSAYS

TABLE A2-1. MICROTOX® TEST RESULTS ON GRAB SAMPLES OF BEACH POINT BUFFERED 100% GROUNDWATER (pH ≈7) TAKEN FROM WELL CC-33B^a

| Date of Sample | Rep | Microtox® 5-Min Readings | | | Microtox® 15-Min Readings | | |
|----------------|-----|--------------------------|---------------------|-------|---------------------------|---------------------|-------|
| | | 5-min EC50 | 95% Fiducial Limits | Slope | 15-min EC50 | 95% Fiducial Limits | Slope |
| Apr 28 | 1 | 24.8 | 23.24-26.46 | 1.02 | 29.3 | 24.36-35.25 | 1.05 |
| | 2 | 33.3 | 25.01-44.26 | 1.05 | 38.0 | 34.63-41.75 | 1.00 |
| May 07 | 1 | 16.9 | 07.85-36.22 | 0.76 | 18.4 | 09.59-35.13 | 0.72 |
| | 2 | 18.7 | 12.99-26.88 | 0.86 | 22.3 | 14.69-33.95 | 0.82 |
| 12 | 1 | 18.8 | 15.95-22.10 | 0.85 | 22.8 | 21.75-23.83 | 0.82 |
| | 2 | 22.0 | 20.81-23.22 | 0.90 | 26.6 | 20.09-35.33 | 0.81 |
| 19 | 1 | 17.5 | 13.04-23.54 | 0.86 | 21.8 | 16.30-29.14 | 0.85 |
| | 2 | 23.1 | 20.52-26.10 | 0.94 | 29.3 | 28.03-30.55 | 0.83 |
| 26 | 1 | 23.0 | 18.51-28.67 | 0.86 | 27.8 | 23.10-33.37 | 0.82 |
| | 2 | 22.4 _b | 18.50-27.18 | 0.77 | 25.2 _b | 21.14-29.92 | 0.75 |
| Jun 04 | 1 | _b | | | _b | | |
| | 2 | 22.3 | 20.89-23.77 | 0.82 | 26.8 | 22.07-32.45 | 0.82 |
| 09 | 1 | 23.1 | 19.89-26.75 | 0.95 | 28.4 | 21.91-36.72 | 0.92 |
| | 2 | 20.9 | 17.39-25.00 | 0.96 | 22.7 | 18.40-28.06 | 0.88 |
| 16 | 1 | 21.3 | 15.63-29.00 | 1.01 | 25.1 | 19.05-33.17 | 0.97 |
| | 2 | 27.0 | 24.92-29.34 | 0.84 | 32.8 | 26.74-40.18 | 0.81 |
| 23 | 1 | 24.8 | 19.61-31.38 | 0.90 | 28.0 | 21.37-36.64 | 0.82 |
| | 2 | 27.2 | 20.84-35.57 | 0.99 | 31.6 | 23.29-42.77 | 0.94 |
| 30 | 1 | 28.2 _c | 17.19-46.21 | 0.87 | 33.3 _c | 17.70-62.71 | 0.86 |
| | 2 | _c | | | _c | | |
| Jul 07 | 1 | 23.8 | 13.82-40.83 | 0.92 | 29.0 | 13.47-62.58 | 0.90 |
| | 2 | 22.7 | 21.45-24.05 | 0.84 | 28.3 | 25.34-31.56 | 0.83 |
| 14 | 1 | 21.7 | 18.00-26.23 | 0.80 | 25.5 | 23.88-27.34 | 0.76 |
| | 2 | 28.8 _b | 22.46-36.89 | 0.87 | 34.9 _b | 24.01-50.62 | 0.86 |
| 21 | 1 | _b | | | _b | | |
| | 2 | _b | | | _b | | |
| 28 | 1 | | | | | | |
| | 2 | | | | | | |

TABLE A2-1. (CONTINUED)

| Date of Sample | Rep | Microtox® 5-Min Readings | | | Microtox® 15-Min Readings | | |
|----------------|-----|--------------------------|---------------------|-------|---------------------------|---------------------|-------|
| | | 5-min EC50 | 95% Fiducial Limits | Slope | 15-min EC50 | 95% Fiducial Limits | Slope |
| Aug 06 | 1 | 27.7 | 24.00-31.98 | 0.83 | 37.3 | 29.64-47.01 | 0.82 |
| | 2 | 30.2 | 26.32-34.75 | 0.98 | 37.4 | 27.14-51.48 | 1.09 |
| 11 | 1 | 27.2 | 22.70-32.53 | 0.89 | 30.7 | 20.42-46.25 | 0.86 |
| | 2 | 28.8 | 17.35-47.88 | 0.79 | 37.1 | 18.51-74.25 | 0.69 |
| 18 | 1 | 21.6 | 13.40-34.75 | 1.10 | 23.8 | 14.11-40.09 | 1.04 |
| | 2 | 32.5 | 27.44-38.45 | 1.07 | 37.2 | 26.91-51.51 | 1.08 |
| 25 | 1 | 33.7 | 29.52-38.48 | 0.94 | 43.3 | 39.54-47.48 | 0.93 |
| | 2 | 45.6 | 23.96-86.94 | 0.77 | 62.2 | 28.33-136.53 | 0.76 |
| Sep 01 | 1 | 38.7 | 30.26-49.52 | 1.01 | 45.9 | 35.01-60.30 | 1.00 |
| | 2 | 46.8 | 22.72-96.55 | 0.80 | 70.1 | 30.92-158.88 | 0.68 |
| 07 | 1 | 41.6 | 23.19-74.56 | 0.96 | 49.6 | 28.50-86.25 | 1.03 |
| | 2 | 50.4 | 32.13-79.02 | 0.76 | 68.2 | 25.62-181.54 | 0.67 |
| 15 | 1 | 33.1 | 29.74-36.81 | 0.78 | 42.0 | 36.27-48.55 | 0.79 |
| | 2 | 38.1 | 32.91-44.08 | 0.96 | 47.9 | 32.20-71.21 | 0.97 |
| 22 | 1 | 38.9 | 21.19-71.41 | 0.73 | 47.3 | 15.33-145.90 | 0.61 |
| | 2 | 40.7 | 05.45-304.23 | 1.34 | 44.5 | 04.19-472.05 | 1.58 |
| 27 | 1 | 39.5 | 34.20-45.66 | 0.99 | d | | |
| | 2 | 43.0 | 20.66-89.66 | 1.28 | d | | |
| Oct 06 | 1 | 40.9 | 28.98-57.74 | 0.90 | 48.7 | 34.07-69.75 | 0.87 |
| | 2 | 43.1 | 32.96-56.41 | 0.75 | 54.9 | 33.61-89.66 | 0.67 |
| 13 | 1 | 52.3 | 45.56-60.19 | 0.85 | 65.2 | 59.34-71.73 | 0.82 |
| | 2 | 45.0 | 37.34-54.24 | 0.92 | 60.5 | 55.41-66.06 | 0.88 |
| 20 | 1 | 39.9 | 27.91-56.95 | 0.96 | 47.9 | 28.60-80.29 | 0.88 |
| | 2 | 43.7 | 30.48-62.69 | 0.88 | 57.7 | 52.74-63.09 | 0.96 |
| 27 | 1 | 39.1 | 33.67-45.44 | 0.95 | 53.8 | 40.25-71.82 | 0.96 |
| | 2 | 45.4 | 34.58-59.64 | 0.89 | 69.1 | 40.14-118.93 | 0.87 |

TABLE A2-1. (CONTINUED)

| Date of Sample | Rep | Microtox® 5-Min Readings | | | Microtox® 15-Min Readings | | |
|----------------|-----|--------------------------|---------------------|-------|---------------------------|---------------------|-------|
| | | 5-min EC50 | 95% Fiducial Limits | Slope | 15-min EC50 | 95% Fiducial Limits | Slope |
| Nov 03 | 1 | 39.2 | 23.70-64.95 | 0.91 | 53.2 | 30.03-94.12 | 0.84 |
| | 2 | 54.7 | 38.84-76.94 | 0.88 | 83.5 | 33.51-208.14 | 0.88 |
| | 10 | 41.7 | 31.91-54.48 | 1.02 | 52.0 | 39.93-67.61 | 1.07 |
| | 2 | 48.7 | 36.39-65.12 | 0.84 | 66.4 | 41.14-107.28 | 0.86 |
| 19 | 1 | 50.9 | 27.15-95.30 | 0.74 | 69.6 | 18.91-256.33 | 0.68 |
| | 2 | 54.9 | 51.79-58.15 | 0.74 | 89.0 | 53.86-147.02 | 0.63 |
| 24 | 1 | 40.5 | 35.50-46.11 | 1.01 | 55.7 | 46.12-67.28 | 0.89 |
| | 2 | 38.3 | 32.19-45.68 | 0.95 | 50.8 | 42.44-60.86 | 0.96 |
| Dec 01 | 1 | 46.5 | 30.01-72.09 | 0.75 | 67.4 | 44.82-101.45 | 0.78 |
| | 2 | 59.1 | 46.34-75.40 | 0.81 | 95.6 | 30.11-303.76 | 0.72 |

A2-4

- a EC50s and 95% fiducial limits are expressed as percent groundwater by volume.
 b Microtox® instrument malfunctioned; sent to Microbics Corporation for repair.
 c Microtox® instrument was not available for use.
 d Electrical power surge caused an interrupt in the circuit; consequently, the 15-min readings were lost.

TABLE A2-2. pH OF THE RAW AND BUFFERED 100% GROUNDWATER USED
IN THE MICROTOX® ASSAY

| Date of Sample | Rep | Raw Groundwater | <u>Buffered Groundwater</u> | |
|-------------------|-----|--------------------|-----------------------------|-------|
| | | | Initial | Final |
| Apr 28 | 1 | 4.04 | 7.07 | 7.16 |
| | 2 | 4.04 | 7.07 | 7.16 |
| May 07 | 1 | 3.99 | 7.13 | 7.28 |
| | 2 | 3.99 | 7.13 | 7.28 |
| 12 | 1 | 4.01 | 7.09 | 7.16 |
| | 2 | 4.01 | 7.09 | 7.16 |
| 19 | 1 | 3.97 | 7.05 | 7.12 |
| | 2 | 3.97 | 7.05 | 7.12 |
| 26 | 1 | 4.00 | 7.11 | 7.19 |
| | 2 | 4.00 | 7.11 | 7.19 |
| Jun 09 | 1 | 4.03 | 7.00 | 7.02 |
| | 2 | 4.03 | 7.00 | 7.02 |
| 16 | 1 | 4.06 | 7.05 | 7.07 |
| | 2 | 4.06 | 7.05 | 7.07 |
| 23 | 1 | 4.02 | 7.01 | 7.03 |
| | 2 | 4.02 | 7.01 | 7.03 |
| 30 | 1 | 4.08 | 7.01 | 7.11 |
| | 2 | 4.08 | 7.01 | 7.11 |
| Jul 07 | 1 | a | | |
| | 2 | a | | |
| 14 | 1 | 4.02 | 6.90 | 6.87 |
| | 2 | 4.02 | 6.90 | 6.87 |
| 21 | 1 | 4.06 | 6.99 | 7.07 |
| | 2 | 4.06 | 6.99 | 7.07 |
| 28 | 1 | b | | |
| | | | | |
| Aug 06 | 1 | 3.98 | 6.98 | 7.01 |
| | 2 | 3.98 | 6.98 | 7.01 |
| 11 | 1 | 3.98 | 6.94 | 7.01 |
| | 2 | 3.98 | 6.94 | 7.01 |
| 18 | 1 | 4.02 | 7.04 | 7.09 |
| | 2 | 4.02 | 7.04 | 7.09 |
| Aug 25 | 1 | 4.05 | 6.95 | 6.98 |
| | 2 | 4.05 | 6.95 | 6.98 |
| Sep 01 | 1 | 4.06 | 7.11 | 7.18 |
| | 2 | 4.06 | 7.11 | 7.18 |
| 07 | 1 | 4.01 | 6.93 | 6.96 |
| | 2 | 4.01 | 6.93 | 6.96 |
| 15 | 1 | 4.01 | 7.02 | 7.13 |
| | 2 | 4.01 | 7.02 | 7.13 |
| 22 | 1 | 3.99 | 7.00 | 7.07 |
| | 2 | 3.99 | 7.00 | 7.07 |
| 27 | 1 | 3.99 | 7.06 | 7.11 |
| | 2 | 3.99 | 7.06 | 7.11 |

TABLE A2-2. (CONTINUED)

| Date of Sample | Rep | Raw Groundwater | <u>Buffered Groundwater</u> | |
|----------------|-----|-----------------|-----------------------------|-------|
| | | | Initial | Final |
| Oct 06 | 1 | 4.02 | 7.01 | 7.11 |
| | 2 | 4.02 | 7.01 | 7.11 |
| | 1 | 4.03 | 6.97 | 7.12 |
| | 2 | 4.03 | 6.97 | 7.12 |
| | 1 | 4.07 | 7.09 | 7.18 |
| | 2 | 4.07 | 7.09 | 7.18 |
| | 1 | 4.08 | 7.00 | 7.10 |
| | 2 | 4.08 | 7.00 | 7.10 |
| | 1 | 4.00 | 6.94 | 7.02 |
| | 2 | 4.00 | 6.94 | 7.02 |
| Nov 03 | 1 | 4.03 | 7.00 | 7.10 |
| | 2 | 4.03 | 7.00 | 7.10 |
| | 1 | 3.99 | 6.93 | 7.09 |
| | 2 | 3.99 | 6.93 | 7.09 |
| | 1 | 4.02 | 7.04 | 7.12 |
| | 2 | 4.02 | 7.04 | 7.12 |
| Dec 01 | 1 | 4.05 | 7.01 | 7.09 |
| | 2 | 4.05 | 7.01 | 7.09 |

^a Microtox® instrument was not available for use.

^b Microtox® instrument malfunctioned and was sent to Microbics Corporation for repair.

APPENDIX 3

ROTIFER 24-H ACUTE TEST CONDUCTED ON BEACH POINT RAW (pH \approx 4) GROUNDWATER (WELL CC-33B) (TEST NO. 1)

| | |
|---------------------------|---|
| Test Method: | Rotifer Toxkit™ Screening Test (Bio-Response Systems, Halifax, Nova Scotia) |
| Type of Test: | Static |
| Date: | April 10-12, 1993 |
| Investigator: | S. D. Turley |
| Laboratory: | UMD/WREC |
| Groundwater: | |
| Source: | APG-EA Beach Point Well CC-33B |
| Chemical Characteristics: | See Appendix 48 |
| Test Medium: | Rotifer Toxkit™ synthetic medium |
| Test Organism: | |
| Scientific Name: | <u>Brachionus rubens</u> |
| Age at Start of Test: | <4 h after hatch |
| Source: | Rotifer Toxkit™ cyst |
| Experimental Chambers: | |
| Material: | Glass petri dish |
| Test Solution Volume: | 10 mL |
| No. Organisms/Replicate: | 10 |
| No. Organisms/Treatment: | 30 |
| Loading: | n/a |
| Lighting: | Fluorescent; 60-85 foot candles |
| Aeration: | None |
| Endpoint: | Mortality |

Water Quality:

| | |
|--|--|
| Dissolved Oxygen (mg/L): | Mean: 8.5 Range: 8.2-8.8 APHA Standards Methods (1989) |
| pH (std. units): | Mean: 5.8 Range: 4.0-6.7 APHA Standards Methods (1989) |
| Conductivity (μ mhos/cm): | APG-EA diluent water: 140 100% groundwater: 1500 |
| Alkalinity (mg/L as CaCO_3): | APG-EA diluent water: 50 100% groundwater: n/a |
| Hardness (mg/L as CaCO_3): | APG-EA diluent water: 48 100% groundwater: n/a |
| Temperature ($^{\circ}\text{C}$): | 25 \pm 0.5 |

Results:

100% Beach Point raw groundwater killed 80% of the rotifers in 24 h (Table A3-1). The 24-h LC50, which was determined by the moving average angle method, is as follows:

24-h LC50 = 71.9% raw groundwater by volume (95% confidence limits = 64.26-83.39).

TABLE A3-1. ROTIFER BEACH POINT RAW GROUNDWATER TOXICITY TEST
DATA (TEST NO. 1) - SURVIVAL AFTER 24 HOURS OF
EXPOSURE

| Concentration (% by Volume) | Rep | Number Tested | No. Alive at End of Test | Percent Alive |
|--------------------------------|-----|------------------|-----------------------------|------------------|
| Growth Medium | 1 | 10 | 10 | 100 |
| | 2 | 10 | 10 | 100 |
| | 3 | 10 | 10 | 100 |
| APG-EA Diluent Water | 1 | 10 | 10 | 100 |
| | 2 | 10 | 10 | 100 |
| | 3 | 10 | 10 | 100 |
| 18 | 1 | 10 | 10 | 100 |
| | 2 | 10 | 10 | 100 |
| | 3 | 10 | 9 | 90 |
| 56 | 1 | 10 | 8 | 80 |
| | 2 | 10 | 9 | 90 |
| | 3 | 10 | 10 | 100 |
| 100 | 1 | 10 | 3 | 30 |
| | 2 | 10 | 1 | 10 |
| | 3 | 10 | 2 | 20 |

APPENDIX 4

ROTIFER 24-H ACUTE TEST CONDUCTED ON BEACH POINT
BUFFERED (pH \approx 7) GROUNDWATER (WELL CC-33B)
(TEST NO. 1)

| | |
|---------------------------|---|
| Test Method: | Rotifer Toxkit™ Screening Test (Bio-Response Systems, Halifax, Nova Scotia) |
| Type of Test: | Static |
| Date: | April 10-12, 1993 |
| Investigator: | S. D. Turley |
| Laboratory: | UMD/WREC |
| Groundwater: | |
| Source: | APG-EA Beach Point Well CC-33B |
| Chemical Characteristics: | See Appendix 48 |
| Test Medium: | Rotifer Toxkit™ synthetic medium |
| Test Organism: | |
| Scientific Name: | <u>Brachionus rubens</u> |
| Age at Start of Test: | <4 h after hatch |
| Source: | Rotifer Toxkit™ cyst |
| Experimental Chambers: | |
| Material: | Glass Petri dish |
| Test Solution Volume: | 10 mL |
| No. Organisms/Replicate: | 10 |
| No. Organisms/Treatment: | 30 |
| Loading: | n/a |
| Lighting: | Fluorescent; 60-85 foot candles |
| Aeration: | None |
| pH Buffer: | 10 N NaOH |
| Endpoint: | Mortality |

Water Quality:

| | |
|--|--|
| Dissolved Oxygen (mg/L): | Mean: 8.4 Range: 8.1-8.8 APHA Standards Methods (1989) |
| pH (std. units): | Mean: 6.8 Range: 6.6-7.0 APHA Standards Methods (1989) |
| Conductivity (μ mhos/cm): | APG-EA diluent water: 140 100% groundwater: 1700 |
| Alkalinity (mg/L as CaCO_3): | APG-EA diluent water: 50 100% groundwater: 80 |
| Hardness (mg/L as CaCO_3): | APG-EA diluent water: 48 100% groundwater: 280 |
| Temperature ($^{\circ}\text{C}$): | 25 \pm 0.5 |

Results:

The buffered groundwater did not affect rotifer survival. The data are summarized in Table A4-1.

TABLE A4-1. ROTIFER BEACH POINT BUFFERED GROUNDWATER TOXICITY
TEST DATA (TEST NO. 1) - SURVIVAL AFTER 24 HOURS OF
EXPOSURE

| Concentration (% by Volume) | Rep | Number Tested | No. Alive at End of Test | Percent Alive |
|--------------------------------|-----|------------------|-----------------------------|------------------|
| Growth Medium | 1 | 10 | 10 | 100 |
| | 2 | 10 | 10 | 100 |
| | 3 | 10 | 10 | 100 |
| 56 | 1 | 10 | 9 | 90 |
| | 2 | 10 | 9 | 90 |
| | 3 | 10 | 10 | 100 |
| 100 | 1 | 10 | 8 | 80 |
| | 2 | 10 | 8 | 80 |
| | 3 | 10 | 9 | 90 |

APPENDIX 5

ROTIFER 24-H ACUTE TEST CONDUCTED ON BEACH POINT RAW (pH \approx 4) GROUNDWATER (WELL CC-33B) (TEST NO. 2)

| | |
|---------------------------|---|
| Test Method: | Rotifer Toxkit™ Screening Test (Bio-Response Systems, Halifax, Nova Scotia) |
| Type of Test: | Static |
| Date: | June 2-4, 1993 |
| Investigator: | S. D. Turley |
| Laboratory: | UMD/WREC |
| Groundwater: | |
| Source: | APG-EA Beach Point Well CC-33B |
| Chemical Characteristics: | See Appendix 48 |
| Test Medium: | Rotifer Toxkit™ synthetic medium |
| Test Organism: | |
| Scientific Name: | <u>Brachionus rubens</u> |
| Age at Start of Test: | <4 h after hatch |
| Source: | Rotifer Toxkit™ cyst |
| Experimental Chambers: | |
| Material: | Glass petri dish |
| Test Solution Volume: | 10 mL |
| No. Organisms/Replicate: | 10 |
| No. Organisms/Treatment: | 30 |
| Loading: | n/a |
| Lighting: | Fluorescent; 60-85 foot candles |
| Aeration: | None |
| Endpoint: | Mortality |

Water Quality:

| | |
|--|--|
| Dissolved Oxygen (mg/L): | Mean: 8.3 Range: 8.1-8.4 APHA Standards Methods (1989) |
| pH (std. units): | Mean: 5.9 Range: 3.9-6.8 APHA Standards Methods (1989) |
| Conductivity (μ mhos/cm): | APG-EA diluent water: 150 100% groundwater: 1400 |
| Alkalinity (mg/L as CaCO_3): | APG-EA diluent water: 55 100% groundwater: n/a |
| Hardness (mg/L as CaCO_3): | APG-EA diluent water: 50 100% groundwater: n/a |
| Temperature ($^{\circ}\text{C}$): | 25 \pm 0.5 |

Results:

100% Beach Point raw groundwater killed 87% of the rotifers in 24 h (Table A5-1). The 24-h LC50, which was determined by the moving average angle method, is as follows:

24-h LC50 = 64.4% raw groundwater by volume (95% confidence limits = 57.64-73.54).

TABLE A5-1. ROTIFER BEACH POINT RAW GROUNDWATER TOXICITY TEST
DATA (TEST NO. 2) - SURVIVAL AFTER 24 HOURS OF
EXPOSURE

| Concentration (% by Volume) | Rep | Number Tested | No. Alive at End of Test | Percent Alive |
|--------------------------------|-----|------------------|-----------------------------|------------------|
| Growth Medium | 1 | 10 | 10 | 100 |
| | 2 | 10 | 10 | 100 |
| | 3 | 10 | 10 | 100 |
| APG-EA Diluent Water | 1 | 10 | 10 | 100 |
| | 2 | 10 | 10 | 100 |
| | 3 | 10 | 10 | 100 |
| 18 | 1 | 10 | 10 | 100 |
| | 2 | 10 | 10 | 100 |
| | 3 | 10 | 10 | 100 |
| 32 | 1 | 10 | 9 | 90 |
| | 2 | 10 | 10 | 100 |
| | 3 | 10 | 9 | 90 |
| 56 | 1 | 10 | 8 | 80 |
| | 2 | 10 | 7 | 70 |
| | 3 | 10 | 7 | 70 |
| 100 | 1 | 10 | 1 | 10 |
| | 2 | 10 | 0 | 0 |
| | 3 | 10 | 3 | 30 |

APPENDIX 6

ROTIFER 24-H ACUTE TEST CONDUCTED ON BEACH POINT BUFFERED (pH \approx 7) GROUNDWATER (WELL CC-33B) (TEST NO. 2)

| | |
|---------------------------|---|
| Test Method: | Rotifer Toxkit™ Screening Test (Bio-Response Systems, Halifax, Nova Scotia) |
| Type of Test: | Static |
| Date: | June 2-4, 1993 |
| Investigator: | S. D. Turley |
| Laboratory: | UMD/WREC |
| Groundwater: | |
| Source: | APG-EA Beach Point Well CC-33B |
| Chemical Characteristics: | See Appendix 48 |
| Test Medium: | Rotifer Toxkit™ synthetic medium |
| Test Organism: | |
| Scientific Name: | <u>Brachionus rubens</u> |
| Age at Start of Test: | <4 h after hatch |
| Source: | Rotifer Toxkit™ cyst |
| Experimental Chambers: | |
| Material: | Glass Petri dish |
| Test Solution Volume: | 10 mL |
| No. Organisms/Replicate: | 10 |
| No. Organisms/Treatment: | 30 |
| Loading: | n/a |
| Lighting: | Fluorescent; 60-85 foot candles |
| Aeration: | None |
| pH Buffer: | 10 N NaOH |
| Endpoint: | Mortality |

Water Quality:

| | |
|--|--|
| Dissolved Oxygen (mg/L): | Mean: 7.5 Range: 6.5-8.3 APHA Standards Methods (1989) |
| pH (std. units): | Mean: 6.7 Range: 6.3-7.0 APHA Standards Methods (1989) |
| Conductivity (μ mhos/cm): | APG-EA diluent water: 150 100% groundwater: 1400 |
| Alkalinity (mg/L as CaCO_3): | APG-EA diluent water: 55 100% groundwater: 100 |
| Hardness (mg/L as CaCO_3): | APG-EA diluent water: 50 100% groundwater: 170 |
| Temperature ($^{\circ}\text{C}$): | 25 \pm 0.5 |

Results:

The 100% buffered groundwater killed 33% of the rotifers; therefore, an LC50 could not be obtained since <50% mortality occurred. A summary of the mortality data is given in Table A6-1.

TABLE A6-1. ROTIFER BEACH POINT BUFFERED GROUNDWATER TOXICITY
TEST DATA (TEST NO. 2) - SURVIVAL AFTER 24 HOURS OF
EXPOSURE

| Concentration (% by Volume) | Rep | Number Tested | No. Alive at End of Test | Percent Alive |
|--------------------------------|-----|------------------|-----------------------------|------------------|
| Growth Medium | 1 | 10 | 10 | 100 |
| | 2 | 10 | 10 | 100 |
| | 3 | 10 | 10 | 100 |
| 32 | 1 | 10 | 10 | 100 |
| | 2 | 10 | 10 | 100 |
| | 3 | 10 | 9 | 90 |
| 56 | 1 | 10 | 9 | 90 |
| | 2 | 10 | 8 | 80 |
| | 3 | 10 | 9 | 90 |
| 100 | 1 | 10 | 5 | 50 |
| | 2 | 10 | 7 | 70 |
| | 3 | 10 | 8 | 80 |

APPENDIX 7

ROTIFER 24-H ACUTE TEST CONDUCTED ON BEACH POINT RAW (pH \approx 4) GROUNDWATER (WELL CC-33B) (TEST NO. 3)

| | |
|---------------------------|---|
| Test Method: | Rotifer Toxkit™ Screening Test (Bio-Response Systems, Halifax, Nova Scotia) |
| Type of Test: | Static |
| Date: | August 24-26, 1993 |
| Investigator: | S. D. Turley |
| Laboratory: | UMD/WREC |
| Groundwater: | |
| Source: | APG-EA Beach Point Well CC-33B |
| Chemical Characteristics: | See Appendix 48 |
| Test Medium: | Rotifer Toxkit™ synthetic medium |
| Test Organism: | |
| Scientific Name: | <u>Brachionus rubens</u> |
| Age at Start of Test: | <4 h after hatch |
| Source: | Rotifer Toxkit™ cyst |
| Experimental Chambers: | |
| Material: | Glass petri dish |
| Test Solution Volume: | 10 mL |
| No. Organisms/Replicate: | 10 |
| No. Organisms/Treatment: | 30 |
| Loading: | n/a |
| Lighting: | Fluorescent; 60-85 foot candles |
| Aeration: | None |
| Endpoint: | Mortality |

Water Quality:

| | |
|--|--|
| Dissolved Oxygen (mg/L): | Mean: 8.3 Range: 8.0-8.7 APHA Standards Methods (1989) |
| pH (std. units): | Mean: 5.9 Range: 3.9-6.7 APHA Standards Methods (1989) |
| Conductivity (μ mhos/cm): | APG-EA diluent water: 150 100% groundwater: 1800 |
| Alkalinity (mg/L as CaCO_3): | APG-EA diluent water: 50 100% groundwater: n/a |
| Hardness (mg/L as CaCO_3): | APG-EA diluent water: 52 100% groundwater: n/a |
| Temperature ($^{\circ}\text{C}$): | 25 \pm 0.5 |

Results:

100% Beach Point raw groundwater killed 80% of the rotifers in 24 h (Table A7-1). The 24-h LC50, which was determined by the moving average angle method, is as follows:

24-h LC50 = 70.2% raw groundwater by volume (95% confidence limits = 62.37-82.03).

TABLE A7-1. ROTIFER BEACH POINT RAW GROUNDWATER TOXICITY TEST
DATA (TEST NO. 3) - SURVIVAL AFTER 24 HOURS OF
EXPOSURE

| Concentration (% by Volume) | Rep | Number Tested | No. Alive at End of Test | Percent Alive |
|--------------------------------|-----|------------------|-----------------------------|------------------|
| Growth Medium | 1 | 10 | 10 | 100 |
| | 2 | 10 | 9 | 90 |
| | 3 | 10 | 10 | 100 |
| APG-EA Diluent Water | 1 | 10 | 10 | 100 |
| | 2 | 10 | 10 | 100 |
| | 3 | 10 | 10 | 100 |
| 10 | 1 | 10 | 10 | 100 |
| | 2 | 10 | 10 | 100 |
| | 3 | 10 | 9 | 90 |
| 18 | 1 | 10 | 10 | 100 |
| | 2 | 10 | 10 | 100 |
| | 3 | 10 | 10 | 100 |
| 32 | 1 | 10 | 9 | 90 |
| | 2 | 10 | 10 | 100 |
| | 3 | 10 | 10 | 100 |
| 56 | 1 | 10 | 8 | 80 |
| | 2 | 10 | 7 | 70 |
| | 3 | 10 | 9 | 90 |
| 100 | 1 | 10 | 2 | 20 |
| | 2 | 10 | 3 | 30 |
| | 3 | 10 | 1 | 10 |

APPENDIX 8

ROTIFER 24-H ACUTE TEST CONDUCTED ON BEACH POINT BUFFERED (pH \approx 7) GROUNDWATER (WELL CC-33B) (TEST NO. 3)

| | |
|---------------------------|--|
| Test Method: | Rotifer Toxkit™ Screening Test (Bio-Response Systems, Halifax, Novia Scotia) |
| Type of Test: | Static |
| Date: | August 24-26, 1993 |
| Investigator: | S. D. Turley |
| Laboratory: | UMD/WREC |
| Groundwater: | |
| Source: | APG-EA Beach Point Well CC-33B |
| Chemical Characteristics: | See Appendix 48 |
| Test Medium: | Rotifer Toxkit™ synthetic medium |
| Test Organism: | |
| Scientific Name: | <u>Brachionus rubens</u> |
| Age at Start of Test: | <4 h after hatch |
| Source: | Rotifer Toxkit™ cyst |
| Experimental Chambers: | |
| Material: | Glass Petri dish |
| Test Solution Volume: | 10 mL |
| No. Organisms/Replicate: | 10 |
| No. Organisms/Treatment: | 30 |
| Loading: | n/a |
| Lighting: | Fluorescent; 60-85 foot candles |
| Aeration: | None |
| pH Buffer: | 10 N NaOH |
| Endpoint: | Mortality |

Water Quality:

| | |
|--------------------------|-------------------------------|
| Dissolved Oxygen (mg/L): | Mean: 8.2 |
| | Range: 7.9-8.4 |
| | APHA Standards Methods (1989) |

| | |
|------------------|-------------------------------|
| pH (std. units): | Mean: 7.1 |
| | Range: 6.9-7.7 |
| | APHA Standards Methods (1989) |

| | |
|--------------------------------|---------------------------|
| Conductivity (μ mhos/cm): | APG-EA diluent water: 150 |
| | 100% groundwater: 1900 |

| | |
|--|--------------------------|
| Alkalinity (mg/L as CaCO_3): | APG-EA diluent water: 50 |
| | 100% groundwater: 100 |

| | |
|--------------------------------------|--------------------------|
| Hardness (mg/L as CaCO_3): | APG-EA diluent water: 48 |
| | 100% groundwater: 210 |

| | |
|-------------------------------------|--------------|
| Temperature ($^{\circ}\text{C}$): | 25 \pm 0.5 |
|-------------------------------------|--------------|

Results:

The buffered groundwater did not affect rotifer survival. The data are summarized in Table A8-1.

TABLE A8-1. ROTIFER BEACH POINT BUFFERED GROUNDWATER TOXICITY
TEST DATA (TEST NO. 3) - SURVIVAL AFTER 24 HOURS OF
EXPOSURE

| Concentration (% by Volume) | Rep | Number Tested | No. Alive at End of Test | Percent Alive |
|--------------------------------|-----|------------------|-----------------------------|------------------|
| Growth Medium | 1 | 10 | 10 | 100 |
| | 2 | 10 | 9 | 90 |
| | 3 | 10 | 10 | 100 |
| 18 | 1 | 10 | 10 | 100 |
| | 2 | 10 | 10 | 100 |
| | 3 | 10 | 10 | 100 |
| 32 | 1 | 10 | 9 | 90 |
| | 2 | 10 | 10 | 100 |
| | 3 | 10 | 10 | 100 |
| 56 | 1 | 10 | 9 | 90 |
| | 2 | 10 | 10 | 100 |
| | 3 | 10 | 8 | 80 |
| 100 | 1 | 10 | 8 | 80 |
| | 2 | 10 | 6 | 60 |
| | 3 | 10 | 7 | 70 |

APPENDIX 9

ROTIFER 24-H ACUTE TEST CONDUCTED ON BEACH POINT RAW (pH \approx 4) GROUNDWATER (WELL CC-33B) (TEST NO. 4)

| | |
|---------------------------|---|
| Test Method: | Rotifer Toxkit™ Screening Test (Bio-Response Systems, Halifax, Nova Scotia) |
| Type of Test: | Static |
| Date: | December 1-3, 1993 |
| Investigator: | S. D. Turley |
| Laboratory: | UMD/WREC |
| Groundwater: | |
| Source: | APG-EA Beach Point Well CC-33B |
| Chemical Characteristics: | See Appendix 48 |
| Test Medium: | Rotifer Toxkit™ synthetic medium |
| Test Organism: | |
| Scientific Name: | <u>Brachionus rubens</u> |
| Age at Start of Test: | <4 h after hatch |
| Source: | Rotifer Toxkit™ cyst |
| Experimental Chambers: | |
| Material: | Glass petri dish |
| Test Solution Volume: | 10 mL |
| No. Organisms/Replicate: | 10 |
| No. Organisms/Treatment: | 30 |
| Loading: | n/a |
| Lighting: | Fluorescent; 60-85 foot candles |
| Aeration: | None |
| Endpoint: | Mortality |

Water Quality:

| | |
|--|--|
| Dissolved Oxygen (mg/L): | Mean: 8.6 Range: 8.3-8.9 APHA Standards Methods (1989) |
| pH (std. units): | Mean: 6.1 Range: 3.8-7.7 APHA Standards Methods (1989) |
| Conductivity (μ mhos/cm): | APG-EA diluent water: 150 100% groundwater: 1750 APHA Standards Methods (1989) |
| Alkalinity (mg/L as CaCO_3): | APG-EA diluent water: 50 100% groundwater: n/a APHA Standards Methods (1989) |
| Hardness (mg/L as CaCO_3): | APG-EA diluent water: 54 100% groundwater: n/a APHA Standards Methods (1989) |
| Temperature ($^{\circ}\text{C}$): | 25 \pm 0.5 |

Results:

100% Beach Point raw groundwater killed 53% of the rotifers in 24 h (Table A9-1). The 24-h LC50, which was determined by the moving average angle method, is as follows:

24-h LC50 = 94.5% raw groundwater by volume (95% confidence limits = 75.84-147.56).

TABLE A9-1. ROTIFER BEACH POINT RAW GROUNDWATER TOXICITY TEST
DATA (TEST NO. 4) - SURVIVAL AFTER 24 HOURS OF
EXPOSURE

| Concentration (% by Volume) | Rep | Number Tested | No. Alive at End of Test | Percent Alive |
|--------------------------------|-----|------------------|-----------------------------|------------------|
| Growth Medium | 1 | 10 | 10 | 100 |
| | 2 | 10 | 9 | 90 |
| | 3 | 10 | 10 | 100 |
| APG-EA Diluent Water | 1 | 10 | 10 | 100 |
| | 2 | 10 | 10 | 100 |
| | 3 | 10 | 8 | 80 |
| 10 | 1 | 10 | 10 | 100 |
| | 2 | 10 | 10 | 100 |
| | 3 | 10 | 10 | 100 |
| 18 | 1 | 10 | 9 | 90 |
| | 2 | 10 | 10 | 100 |
| | 3 | 10 | 10 | 100 |
| 32 | 1 | 10 | 9 | 90 |
| | 2 | 10 | 9 | 90 |
| | 3 | 10 | 10 | 100 |
| 56 | 1 | 10 | 8 | 80 |
| | 2 | 10 | 10 | 100 |
| | 3 | 10 | 8 | 80 |
| 100 | 1 | 10 | 6 | 60 |
| | 2 | 10 | 4 | 40 |
| | 3 | 10 | 4 | 40 |

APPENDIX 10

ROTIFER 24-H ACUTE TEST CONDUCTED ON BEACH POINT
BUFFERED (pH \approx 7) GROUNDWATER (WELL CC-33B)
(TEST NO. 4)

| | |
|---------------------------|---|
| Test Method: | Rotifer Toxkit™ Screening Test (Bio-Response Systems, Halifax, Nova Scotia) |
| Type of Test: | Static |
| Date: | December 1-2, 1993 |
| Investigator: | S. D. Turley |
| Laboratory: | UMD/WREC |
| Groundwater: | |
| Source: | APG-EA Beach Point Well CC-33B |
| Chemical Characteristics: | See Appendix 48 |
| Test Medium: | Rotifer Toxkit™ synthetic medium |
| Test Organism: | |
| Scientific Name: | <u>Brachionus rubens</u> |
| Age at Start of Test: | <4 h after hatch |
| Source: | Rotifer Toxkit™ cyst |
| Experimental Chambers: | |
| Material: | Glass Petri dish |
| Test Solution Volume: | 10 mL |
| No. Organisms/Replicate: | 10 |
| No. Organisms/Treatment: | 30 |
| Loading: | n/a |
| Lighting: | Fluorescent; 60-85 foot candles |
| Aeration: | None |
| pH Buffer: | 10 N NaOH |
| Endpoint: | Mortality |

Water Quality:

| | |
|--|--|
| Dissolved Oxygen (mg/L): | Mean: 8.4 Range: 8.1-8.8 APHA Standards Methods (1989) |
| pH (std. units): | Mean: 7.3 Range: 7.1-7.6 APHA Standards Methods (1989) |
| Conductivity (μ mhos/cm): | APG-EA diluent water: 190 100% groundwater: 1800 APHA Standards Methods (1989) |
| Alkalinity (mg/L as CaCO_3): | APG-EA diluent water: 50 100% groundwater: 60 APHA Standards Methods (1989) |
| Hardness (mg/L as CaCO_3): | APG-EA diluent water: 50 100% groundwater: 198 APHA Standards Methods (1989) |
| Temperature ($^{\circ}\text{C}$): | 25 \pm 0.5 |

Results:

The buffered groundwater did not affect rotifer survival.
The data are summarized in Table A10-1.

TABLE A10-1. ROTIFER BEACH POINT BUFFERED GROUNDWATER TOXICITY
TEST DATA (TEST NO. 4) - SURVIVAL AFTER 24 HOURS OF
EXPOSURE

| Concentration (% by Volume) | Rep | Number Tested | No. Alive at End of Test | Percent Alive |
|--------------------------------|-----|------------------|-----------------------------|------------------|
| Growth Medium | 1 | 10 | 10 | 100 |
| | 2 | 10 | 9 | 90 |
| | 3 | 10 | 10 | 100 |
| 10 | 1 | 10 | 9 | 90 |
| | 2 | 10 | 10 | 100 |
| | 3 | 10 | 10 | 100 |
| 18 | 1 | 10 | 10 | 100 |
| | 2 | 10 | 10 | 100 |
| | 3 | 10 | 10 | 100 |
| 32 | 1 | 10 | 9 | 90 |
| | 2 | 10 | 10 | 100 |
| | 3 | 10 | 10 | 100 |
| 56 | 1 | 10 | 8 | 80 |
| | 2 | 10 | 9 | 90 |
| | 3 | 10 | 10 | 100 |
| 100 | 1 | 10 | 7 | 70 |
| | 2 | 10 | 9 | 90 |
| | 3 | 10 | 8 | 80 |

APPENDIX 11

GREEN ALGAL 96-H GROWTH TEST CONDUCTED ON BEACH POINT RAW (pH \approx 4) GROUNDWATER (WELL CC-33B) (TEST NO. 1)

Test Method: EPA/600/4-89/001
(Weber et al., 1989)

Type of Test: Static

Date: April 10-14, 1993

Investigator: S. D. Turley

Laboratory: UMD/WREC

Groundwater:

Source: APG-EA Beach Point Well CC-33B

Chemical Characteristics: See Appendix 48

Test Medium: Double strength "AAP" medium
(Miller et al., 1978) with P
added to achieve a 20:1 N:P
atomic ratio

Test Organism:

Scientific Name: Selenastrum capricornutum

Age at Start of Test: Log growth

Source: UMD/WREC culture

Experimental Chambers:

Material: 250 mL glass culture flasks
with cheesecloth/cotton
stoppers

Test Solution Volume: 100 mL

Initial Cell Density: 1×10^4 cells/mL

No. Replicates per Treatment: 3

Lighting: Fluorescent; cool white;
continuous; \approx 300 foot candles

Shaking Rate: 100 cpm continuously

Endpoint: Reduction in growth relative
to control

Temperature:

25 ± 0.2 °C

Results:

100% raw Beach Point groundwater killed all algal cells during the first 24 h of exposure (Table A11-1). Growth (cell density) was not affected by exposure to concentrations up to 56% raw groundwater by volume (see Tables A11-1 and A11-2). The NOEC and LOEC for reduction in growth are as follows:

NOEC = 56% raw groundwater by volume.

LOEC = 100% raw groundwater by volume.

The 96-h EC50 (reduction in growth), which was determined by the moving average angle method, is as follows:

96-h EC50 = 59.4% raw groundwater by volume (95% confidence limits = 57.01-61.86).

TABLE A11-1. GREEN ALGA BEACH POINT RAW GROUNDWATER TOXICITY
DATA (TEST NO. 1) - MEAN CELL DENSITY (CELLS/ML)
AFTER 96 HOURS OF EXPOSURE

| Conc (% by Vol) | Rep | Mean Cell Density | | | | |
|----------------------------|-----|-------------------|-------|--------|--------|---------|
| | | 0H | 24H | 48H | 72H | 96H |
| Growth Medium | 1 | 10120 | 60120 | 328400 | 866210 | 1333333 |
| | 2 | 10120 | 64240 | 341080 | 828800 | 1482400 |
| | 3 | 10120 | 63180 | 336080 | 808640 | 1480400 |
| APG-EA Diluent Water | 1 | 10320 | 62000 | 325000 | 820200 | 1320000 |
| | 2 | 10320 | 65200 | 341300 | 822800 | 1510000 |
| | 3 | 10320 | 60020 | 333333 | 811100 | 1453200 |
| 10 | 1 | 10430 | 61000 | 339020 | 816400 | 1395000 |
| | 2 | 10430 | 64180 | 327800 | 802120 | 1341400 |
| | 3 | 10430 | 59300 | 331400 | 808300 | 1380800 |
| 18 | 1 | 10000 | 62340 | 348000 | 820140 | 1404100 |
| | 2 | 10000 | 62000 | 326200 | 798000 | 1290900 |
| | 3 | 10000 | 60040 | 330120 | 804600 | 1320400 |
| 32 | 1 | 10480 | 63220 | 336120 | 818800 | 1330360 |
| | 2 | 10480 | 61040 | 330080 | 810100 | 1288200 |
| | 3 | 10480 | 60080 | 340000 | 820800 | 1380400 |
| 56 | 1 | 9900 | 58600 | 326800 | 798800 | 1278000 |
| | 2 | 9900 | 59800 | 322600 | 790400 | 1299200 |
| | 3 | 9900 | 57960 | 330140 | 812200 | 1268200 |
| 100 | 1 | 10120 | DEAD | | | |
| | 2 | 10120 | DEAD | | | |
| | 3 | 10120 | DEAD | | | |

Table A11-2. GREEN ALGA BEACH POINT RAW GROUNDWATER TOXICITY
TEST STATISTICAL ANALYSIS (TEST NO. 1) - MEAN CELL
DENSITY (CELLS/ML)^a

Data Transformation:

None

Chi-Square Test for Normality:

| | |
|----------------------------|---|
| Calculated test statistic: | 5.15 |
| Alpha value: | 0.01 |
| Critical value: | 13.28 |
| Conclusion: | Fail to reject the null hypothesis that the data are normally distributed |

Bartlett's Test for Homogeneity of Variances:

| | |
|----------------------------|--|
| Calculated test statistic: | 5.99 |
| Alpha value: | 0.01 |
| Critical value: | 15.09 |
| Conclusion: | Fail to reject the null hypothesis that the variances are homogenous |

ANOVA:

| | |
|----------------------------|--|
| Calculated test statistic: | 2.63 |
| Alpha value: | 0.05 |
| Critical value: | 3.11 |
| Conclusion: | Fail to reject the null hypothesis that all groups are equal |

^a The 100% raw Beach Point groundwater treatment was not included in the statistical analyses because all cells died during the test.

APPENDIX 12

GREEN ALGAL 96-H GROWTH TEST CONDUCTED ON BEACH POINT BUFFERED (pH \approx 7) GROUNDWATER (WELL CC-33B) (TEST NO. 1)

| | |
|-------------------------------|---|
| Test Method: | EPA/600/4-89/001 (Weber et al., 1989) |
| Type of Test: | Static |
| Date: | April 10-14, 1993 |
| Investigator: | S. D. Turley |
| Laboratory: | UMD/WREC |
| Groundwater: | |
| Source: | APG-EA Beach Point Well CC-33B |
| Chemical Characteristics: | See Appendix 48 |
| Test Medium: | Double strength "AAP" medium (Miller et al., 1978) with P added to achieve a 20:1 N:P atomic ratio |
| Test Organism: | |
| Scientific Name: | <u>Selenastrum capricornutum</u> |
| Age at Start of Test: | Log growth |
| Source: | UMD/WREC culture |
| Experimental Chambers: | |
| Material: | 250 mL glass culture flasks with cheesecloth/cotton stoppers |
| Test Solution Volume: | 100 mL |
| Initial Cell Density: | 1×10^4 cells/mL |
| No. Replicates per Treatment: | 3 |
| Lighting: | Fluorescent; cool white; continuous; \approx 300 foot candles |
| Shaking Rate: | 100 cpm continuously |
| pH Buffer: | 10 N NaOH |
| Endpoint: | Reduction in growth relative |

to control

Temperature:

25 \pm 0.2 °C

Results:

Significant ($\alpha = 0.05$) reductions in growth (cell density) occurred in 100% buffered Beach Point groundwater (see Tables A12-1, A12-2 and A12-3). Growth was not affected by exposure to concentrations up to 56% buffered groundwater by volume. The NOEC and LOEC for reduction in growth are as follows:

NOEC = 56% buffered groundwater by volume.

LOEC = 100% buffered groundwater by volume.

A 96-h EC50 (reduction in growth) could not be calculated because <50% inhibition of growth occurred in the test.

Table A12-1. GREEN ALGA BEACH POINT BUFFERED GROUNDWATER
TOXICITY DATA (TEST NO. 1) - MEAN CELL DENSITY
(CELLS/ML) AFTER 96 HOURS OF EXPOSURE

| Conc (% by Vol) | Rep | Mean Cell Density | | | | |
|-----------------------|-----|-------------------|-------|--------|--------|---------|
| | | 0H | 24H | 48H | 72H | 96H |
| Growth Medium | 1 | 10120 | 60120 | 328400 | 814000 | 1333333 |
| | 2 | 10120 | 64240 | 341080 | 828800 | 1482400 |
| | 3 | 10120 | 63180 | 336080 | 808640 | 1480400 |
| 32 | 1 | 10560 | 61120 | 340000 | 822600 | 1399300 |
| | 2 | 10560 | 58400 | 330860 | 806200 | 1326940 |
| | 3 | 10560 | 60200 | 333120 | 810200 | 1352200 |
| 56 | 1 | 10600 | 59980 | 328760 | 812200 | 1340000 |
| | 2 | 10600 | 59120 | 341080 | 826400 | 1394400 |
| | 3 | 10600 | 61120 | 324800 | 788800 | 1290460 |
| 100 | 1 | 10040 | 58680 | 318800 | 786240 | 1246400 |
| | 2 | 10040 | 59980 | 330600 | 818100 | 1258800 |
| | 3 | 10040 | 60120 | 326420 | 800800 | 1300300 |

Table A12-2. GREEN ALGA BEACH POINT BUFFERED GROUNDWATER
TOXICITY TEST STATISTICAL ANALYSIS (TEST NO. 1) -
MEAN CELL DENSITY (CELLS/ML)

Data Transformation:

None

Chi-Square Test for Normality:

| | |
|----------------------------|---|
| Calculated test statistic: | 4.08 |
| Alpha value: | 0.01 |
| Critical value: | 13.28 |
| Conclusion: | Fail to reject the null hypothesis that the data are normally distributed |

Bartlett's Test for Homogeneity of Variance:

| | |
|----------------------------|--|
| Calculated test statistic: | 2.30 |
| Alpha value: | 0.01 |
| Critical value: | 11.34 |
| Conclusion: | Fail to reject the null hypothesis that the variances are homogenous |

ANOVA:

| | |
|----------------------------|--|
| Calculated test statistic: | 4.45 |
| Alpha value: | 0.05 |
| Critical value: | 4.07 |
| Conclusion: | Reject the null hypothesis that all groups are equal |

Dunnett's Test:

| | |
|----------------------------|--|
| Calculated test statistic: | See Table A12-3 |
| Alpha value: | 0.05 |
| Critical value: | 2.42 |
| Conclusion: | Reject the null hypothesis that all groups are equal |

TABLE A12-3. GREEN ALGA BEACH POINT BUFFERED GROUNDWATER
TOXICITY TEST STATISTICAL ANALYSIS (TEST NO. 1) -
RESULTS OF DUNNETT'S TEST ON MEAN CELL DENSITY
(CELLS/ML)

| Conc (% by Vol) | No. of Reps | Mean Cell Density | T Statistic | Significance |
|-----------------------|-------------------|----------------------|-------------|--------------|
| Growth Medium | 3 | 1432044 | | |
| 32 | 3 | 1359480 | 1.61 | |
| 56 | 3 | 1341620 | 2.01 | |
| 100 | 3 | 1268500 | 3.63 | * |

* Significantly different at $\alpha = 0.05$ (Dunnett's critical value = 2.42).

APPENDIX 13

GREEN ALGAL 96-H GROWTH TEST CONDUCTED ON BEACH POINT RAW (pH \approx 4) GROUNDWATER (WELL CC-33B) (TEST NO. 2)

| | |
|-------------------------------|---|
| Test Method: | EPA/600/4-89/001 (Weber et al., 1989) |
| Type of Test: | Static |
| Date: | June 3-7, 1993 |
| Investigator: | S. D. Turley |
| Laboratory: | UMD/WREC |
| Groundwater: | |
| Source: | APG-EA Beach Point Well CC-33B |
| Chemical Characteristics: | See Appendix 48 |
| Test Medium: | Double strength "AAP" medium (Miller et al., 1978) with P added to achieve a 20:1 N:P atomic ratio |
| Test Organism: | |
| Scientific Name: | <u>Selenastrum capricornutum</u> |
| Age at Start of Test: | Log growth |
| Source: | UMD/WREC culture |
| Experimental Chambers: | |
| Material: | 250 mL glass culture flasks with cheesecloth/cotton stoppers |
| Test Solution Volume: | 100 mL |
| Initial Cell Density: | 1×10^4 cells/mL |
| No. Replicates per Treatment: | 3 |
| Lighting: | Fluorescent; cool white; continuous; \approx 300 foot candles |
| Shaking Rate: | 100 cpm continuously |
| Endpoint: | Reduction in growth relative to control |

Temperature:

25 ± 0.2 °C

Results:

100% raw Beach Point groundwater killed all algal cells during the first 24 h of exposure (Table A13-1). A significant ($\alpha = 0.05$) reduction in cell density occurred in algae exposed to 32% and 56% raw groundwater by volume (see Tables A13-1, A13-2 and A13-3). The NOEC and LOEC for reduction in algal growth are as follows:

NOEC = 18% raw groundwater by volume.

LOEC = 32% raw groundwater by volume.

The 96-h EC50 (reduction in growth), which was determined by the moving average angle method, is as follows:

96-h EC50 = 51.3% raw groundwater by volume (95% confidence limits = 45.15-58.19).

TABLE A13-1. GREEN ALGA BEACH POINT RAW GROUNDWATER TOXICITY
DATA (TEST NO. 2) - MEAN CELL DENSITY (CELLS/ML)
AFTER 96 HOURS OF EXPOSURE

| Conc (% by Vol) | Rep | Mean Cell Density | | | | |
|----------------------------|-----|-------------------|-------|--------|--------|---------|
| | | 0H | 24H | 48H | 72H | 96H |
| Growth Medium | 1 | 9880 | 56500 | 313200 | 788880 | 1369800 |
| | 2 | 9880 | 59480 | 309980 | 791140 | 1371640 |
| | 3 | 9880 | 54980 | 315420 | 800000 | 1402760 |
| APG-EA Diluent Water | 1 | 10400 | 57780 | 315800 | 801160 | 1409400 |
| | 2 | 10400 | 55560 | 310400 | 780080 | 1359480 |
| | 3 | 10400 | 56120 | 313600 | 790100 | 1376600 |
| 10 | 1 | 9920 | 60080 | 318000 | 781400 | 1359600 |
| | 2 | 9920 | 55000 | 309600 | 789860 | 1381640 |
| | 3 | 9920 | 54120 | 307980 | 761480 | 1363300 |
| 18 | 1 | 10120 | 58880 | 311120 | 780420 | 1363420 |
| | 2 | 10120 | 51600 | 304880 | 769920 | 1349120 |
| | 3 | 10120 | 59400 | 315580 | 784240 | 1372480 |
| 32 | 1 | 10300 | 51110 | 309400 | 788120 | 1240800 |
| | 2 | 10300 | 51940 | 299980 | 741760 | 1190980 |
| | 3 | 10300 | 53680 | 307340 | 760740 | 1209980 |
| 56 | 1 | 11000 | 47420 | 286124 | 639560 | 980880 |
| | 2 | 11000 | 44800 | 251160 | 618540 | 926140 |
| | 3 | 11000 | 50120 | 270280 | 630680 | 950460 |
| 100 | 1 | 10320 | DEAD | | | |
| | 2 | 10320 | DEAD | | | |
| | 3 | 10320 | DEAD | | | |

Table A13-2. GREEN ALGA BEACH POINT RAW GROUNDWATER TOXICITY
TEST STATISTICAL ANALYSIS (TEST NO. 2) - MEAN CELL
DENSITY (CELLS/ML)^a

Data Transformation:

None

Chi-Square Test for Normality:

| | |
|----------------------------|---|
| Calculated test statistic: | 5.15 |
| Alpha value: | 0.01 |
| Critical value: | 13.28 |
| Conclusion: | Fail to reject the null hypothesis that the data are normally distributed |

Bartlett's Test for Homogeneity of Variances:

| | |
|----------------------------|--|
| Calculated test statistic: | 2.17 |
| Alpha value: | 0.01 |
| Critical value: | 15.09 |
| Conclusion: | Fail to reject the null hypothesis that the variances are homogenous |

ANOVA:

| | |
|----------------------------|--|
| Calculated test statistic: | 199.31 |
| Alpha value: | 0.05 |
| Critical value: | 3.11 |
| Conclusion: | Reject the null hypothesis that all groups are equal |

Dunnett's Test:

| | |
|----------------------------|--|
| Calculated test statistic: | See Table A13-3 |
| Alpha value: | 0.05 |
| Critical value: | 2.50 |
| Conclusion: | Reject the null hypothesis that all groups are equal |

^a The 100% raw Beach Point groundwater treatment was not included in the statistical analyses because all cells died during the test.

Table A13-3. GREEN ALGA BEACH POINT RAW GROUNDWATER TOXICITY
TEST STATISTICAL ANALYSIS (TEST NO. 2) - RESULTS OF
DUNNETT'S TEST ON MEAN CELL DENSITY (CELLS/ML)

| Conc (% by Vol) | No. of Reps | Mean Cell Density | T Statistic | Significance |
|----------------------------|-------------------|----------------------|-------------|--------------|
| Growth Medium | 3 | 1381400 | | |
| APG-EA Diluent Water | 3 | 1381827 | -0.03 | |
| 10 | 3 | 1368180 | 0.77 | |
| 18 | 3 | 1361673 | 1.15 | |
| 32 | 3 | 1213920 | 9.76 | * |
| 56 | 3 | 952493 | 25.00 | * |

* Significantly different at $\alpha = 0.05$ (Dunnett's critical
value = 2.50).

APPENDIX 14

GREEN ALGAL 96-H GROWTH TEST CONDUCTED ON BEACH POINT BUFFERED (pH \approx 7) GROUNDWATER (WELL CC-33B) (TEST NO. 2)

Test Method: EPA/600/4-89/001
(Weber et al., 1989)

Type of Test: Static

Date: June 3-7, 1993

Investigator: S. D. Turley

Laboratory: UMD/WREC

Groundwater:

Source: APG-EA Beach Point Well CC-33B
Chemical Characteristics: See Appendix 48

Test Medium: Double strength "AAP" medium
(Miller et al., 1978) with P
added to achieve a 20:1 N:P
atomic ratio

Test Organism:

Scientific Name: Selenastrum capricornutum
Age at Start of Test: Log growth
Source: UMD/WREC culture

Experimental Chambers:

Material: 250 mL glass culture flasks
with cheesecloth/cotton
stoppers

Test Solution Volume: 100 mL

Initial Cell Density: 1×10^4 cells/mL

No. Replicates per Treatment: 3

Lighting: Fluorescent; cool white;
continuous; \approx 300 foot candles

Shaking Rate: 100 cpm continuously

pH Buffer: 10 N NaOH

Endpoint: Reduction in growth relative

to control

Temperature:

25 \pm 0.2 °C

Results:

100% buffered Beach Point groundwater killed all algal cells during the first 24 h of exposure (see Table A14-1). A significant ($\alpha = 0.05$) reduction in cell density occurred in algae exposed to 56% buffered groundwater by volume (see Tables A14-1, A14-2 and A14-3). The NOEC and LOEC for reduction in growth are as follows:

NOEC = 32% buffered groundwater by volume.

LOEC = 56% buffered groundwater by volume.

The 96-h EC50 (reduction in growth), which was determined by the moving average angle method, is as follows:

96-h EC50 = 58.2% raw groundwater by volume (95% confidence limits = 55.81-60.70).

Table A14-1. GREEN ALGA BEACH POINT BUFFERED GROUNDWATER
TOXICITY DATA (TEST NO. 2) - MEAN CELL DENSITY
(CELLS/ML) AFTER 96 HOURS OF EXPOSURE

| Conc (% by Vol) | Rep | Mean Cell Density | | | | |
|-----------------------|-----|-------------------|-------|--------|--------|---------|
| | | 0H | 24H | 48H | 72H | 96H |
| Growth Medium | 1 | 9880 | 56500 | 313200 | 788880 | 1369800 |
| | 2 | 9880 | 59480 | 309980 | 791140 | 1371640 |
| | 3 | 9880 | 54980 | 315420 | 800000 | 1402760 |
| 32 | 1 | 10600 | 56500 | 316600 | 779690 | 1360340 |
| | 2 | 10600 | 57980 | 317700 | 784600 | 1379960 |
| | 3 | 10600 | 54000 | 304880 | 759980 | 1359980 |
| 56 | 1 | 11200 | 50000 | 281460 | 638080 | 960840 |
| | 2 | 11200 | 48640 | 260480 | 627580 | 919860 |
| | 3 | 11200 | 52180 | 290100 | 649590 | 1000800 |
| 100 | 1 | 10040 | DEAD | | | |
| | 2 | 10040 | DEAD | | | |
| | 3 | 10040 | DEAD | | | |

Table A14-2. GREEN ALGA BEACH POINT BUFFERED GROUNDWATER
TOXICITY TEST STATISTICAL ANALYSIS (TEST NO. 2) -
MEAN CELL DENSITY (CELLS/ML)^a

Data Transformation:

None

Chi-Square Test for Normality:

| | |
|----------------------------|---|
| Calculated test statistic: | 6.90 |
| Alpha value: | 0.01 |
| Critical value: | 13.28 |
| Conclusion: | Fail to reject the null hypothesis that the data are normally distributed |

Bartlett's Test for Homogeneity of Variance:

| | |
|----------------------------|--|
| Calculated test statistic: | 2.55 |
| Alpha value: | 0.01 |
| Critical value: | 9.21 |
| Conclusion: | Fail to reject the null hypothesis that the variances are homogenous |

ANOVA:

| | |
|----------------------------|--|
| Calculated test statistic: | 243.24 |
| Alpha value: | 0.05 |
| Critical value: | 5.14 |
| Conclusion: | Reject the null hypothesis that all groups are equal |

Dunnett's Test:

| | |
|----------------------------|--|
| Calculated test statistic: | See Table A14-3 |
| Alpha value: | 0.05 |
| Critical value: | 2.34 |
| Conclusion: | Reject the null hypothesis that all groups are equal |

^a The 100% buffered Beach Point groundwater treatment was not included in the statistical analyses because all cells died during the test.

TABLE A14-3. GREEN ALGA BEACH POINT BUFFERED GROUNDWATER
TOXICITY TEST STATISTICAL ANALYSIS (TEST NO. 2) -
RESULTS OF DUNNETT'S TEST ON MEAN CELL DENSITY
(CELLS/ML)

| Conc (% by Vol) | No. of Reps | Mean Cell Density | T Statistic | Significance |
|-----------------------|-------------------|----------------------|-------------|--------------|
| Growth Medium | 3 | 1381400 | | |
| 32 | 3 | 1366760 | 0.68 | |
| 56 | 3 | 960500 | 19.43 | * |

* Significantly different at $\alpha = 0.05$ (Dunnett's critical value = 2.34).

APPENDIX 15

GREEN ALGAL 96-H GROWTH TEST CONDUCTED ON BEACH POINT RAW (pH \approx 4) GROUNDWATER (WELL CC-33B) (TEST NO. 3)

Test Method: EPA/600/4-89/001
(Weber et al., 1989)

Type of Test: Static

Date: August 24-28, 1993

Investigator: S. D. Turley

Laboratory: UMD/WREC

Groundwater:

Source: APG-EA Beach Point Well CC-33B
Chemical Characteristics: See Appendix 48

Test Medium: Double strength "AAP" medium
(Miller et al., 1978) with P
added to achieve a 20:1 N:P
atomic ratio

Test Organism:

Scientific Name: Selenastrum capricornutum
Age at Start of Test: Log growth
Source: UMD/WREC culture

Experimental Chambers:

Material: 250 mL glass culture flasks
with cheesecloth/cotton
stoppers

Test Solution Volume: 100 mL

Initial Cell Density: 1×10^4 cells/mL

No. Replicates per Treatment: 3

Lighting: Fluorescent; cool white;
continuous; \approx 300 foot candles

Shaking Rate: 100 cpm continuously

Endpoint: Reduction in growth relative
to control

Temperature:

25 ± 0.2 °C

Results:

100% raw Beach Point groundwater killed all algal cells during the first 24 h of exposure (Table A15-1). A significant ($\alpha = 0.05$) reduction in cell density occurred in algae exposed to 32% and 56% raw groundwater by volume (see Tables A15-1, A15-2 and A15-3). The NOEC and LOEC for reduction in growth are as follows:

NOEC = 18% raw groundwater by volume.

LOEC = 32% raw groundwater by volume.

The 96-h EC50 (reduction in growth), which was determined by the moving average angle method, is as follows:

96-h EC50 = 56.5% raw groundwater by volume (95% confidence limits = 54.04-59.04).

TABLE A15-1. GREEN ALGA BEACH POINT RAW GROUNDWATER TOXICITY
DATA (TEST NO. 3) - MEAN CELL DENSITY (CELLS/ML)
AFTER 96 HOURS OF EXPOSURE

| Conc (% by Vol) | Rep | Mean Cell Density | | | | |
|----------------------------|-----|-------------------|-------|--------|--------|---------|
| | | 0H | 24H | 48H | 72H | 96H |
| Growth Medium | 1 | 10100 | 51900 | 278100 | 681040 | 1182860 |
| | 2 | 10100 | 54800 | 259880 | 666460 | 1159680 |
| | 3 | 10100 | 57950 | 267140 | 670680 | 1176840 |
| APG-EA Diluent Water | 1 | 10400 | 55000 | 270080 | 680080 | 1180000 |
| | 2 | 10400 | 49600 | 255840 | 659980 | 1157980 |
| | 3 | 10400 | 53480 | 259480 | 669840 | 1173640 |
| 10 | 1 | 9900 | 52400 | 272280 | 655780 | 1159180 |
| | 2 | 9900 | 49580 | 270140 | 667740 | 1169240 |
| | 3 | 9900 | 53980 | 256420 | 650640 | 1146140 |
| 18 | 1 | 10000 | 50000 | 251480 | 653880 | 1143640 |
| | 2 | 10000 | 49860 | 263640 | 661880 | 1147160 |
| | 3 | 10000 | 54380 | 269120 | 666360 | 1167260 |
| 32 | 1 | 11000 | 52000 | 264400 | 646680 | 1137640 |
| | 2 | 11000 | 51040 | 260080 | 629860 | 1107480 |
| | 3 | 11000 | 53460 | 267460 | 634980 | 1118240 |
| 56 | 1 | 10800 | 45080 | 232000 | 550840 | 820140 |
| | 2 | 10800 | 48600 | 238200 | 541000 | 804400 |
| | 3 | 10800 | 44320 | 229480 | 505080 | 775180 |
| 100 | 1 | 10800 | DEAD | | | |
| | 2 | 10800 | DEAD | | | |
| | 3 | 10800 | DEAD | | | |

Table A15-2. GREEN ALGA BEACH POINT RAW GROUNDWATER TOXICITY
TEST STATISTICAL ANALYSIS (TEST NO. 3) - MEAN CELL
DENSITY (CELLS/ML)^a

Data Transformation:

None

Chi-Square Test for Normality:

| | |
|----------------------------|---|
| Calculated test statistic: | 3.76 |
| Alpha value: | 0.01 |
| Critical value: | 13.28 |
| Conclusion: | Fail to reject the null hypothesis that the data are normally distributed |

Bartlett's Test for Homogeneity of Variances:

| | |
|----------------------------|--|
| Calculated test statistic: | 1.42 |
| Alpha value: | 0.01 |
| Critical value: | 15.09 |
| Conclusion: | Fail to reject the null hypothesis that the variances are homogenous |

ANOVA:

| | |
|----------------------------|--|
| Calculated test statistic: | 290.41 |
| Alpha value: | 0.05 |
| Critical value: | 3.11 |
| Conclusion: | Reject the null hypothesis that all groups are equal |

Dunnett's Test:

| | |
|----------------------------|--|
| Calculated test statistic: | See Table A15-3 |
| Alpha value: | 0.05 |
| Critical value: | 2.50 |
| Conclusion: | Reject the null hypothesis that all groups are equal |

^a The 100% raw Beach Point groundwater treatment was not included in the statistical analyses because all cells died during the test.

Table A15-3. GREEN ALGA BEACH POINT RAW GROUNDWATER TOXICITY
TEST STATISTICAL ANALYSIS (TEST NO. 3) - RESULTS OF
DUNNETT'S TEST ON MEAN CELL DENSITY (CELLS/ML)

| Conc (% by Vol) | No. of Reps | Mean Cell Density | T Statistic | Significance |
|----------------------------|-------------------|----------------------|-------------|--------------|
| Growth Medium | 3 | 1173127 | | |
| APG-EA Diluent Water | 3 | 1170480 | 0.218 | |
| 10 | 3 | 1158187 | 1.231 | |
| 18 | 3 | 1152627 | 1.690 | |
| 32 | 3 | 1121120 | 4.286 | * |
| 56 | 3 | 799907 | 30.761 | * |

* Significantly different at alpha = 0.05 (Dunnett's critical
value = 2.50).

APPENDIX 16

GREEN ALGAL 96-H GROWTH TEST CONDUCTED ON BEACH POINT BUFFERED (pH \approx 7) GROUNDWATER (WELL CC-33B) (TEST NO. 3)

Test Method: EPA/600/4-89/001
(Weber et al., 1989)

Type of Test: Static

Date: August 24-28, 1993

Investigator: S. D. Turley

Laboratory: UMD/WREC

Groundwater:

Source: APG-EA Beach Point Well CC-33B
Chemical Characteristics: See Appendix 48

Test Medium: Double strength "AAP" medium
(Miller et al., 1978) with P
added to achieve a 20:1 N:P
atomic ratio

Test Organism:

Scientific Name: Selenastrum capricornutum
Age at Start of Test: Log growth
Source: UMD/WREC culture

Experimental Chambers:

Material: 250 mL glass culture flasks
with cheesecloth/cotton
stoppers

Test Solution Volume: 100 mL

Initial Cell Density: 1×10^4 cells/mL

No. Replicates per Treatment: 3

Lighting: Fluorescent; cool white;
continuous; \approx 300 foot candles

Shaking Rate: 100 cpm continuously

pH Buffer: 10 N NaOH

| | |
|--------------|---|
| Endpoint: | Reduction in growth relative to control |
| Temperature: | 25 ± 0.2 °C |

Results:

100% buffered Beach Point groundwater killed all algal cells during the first 24 h of exposure (Table A16-1). A significant ($\alpha = 0.05$) reduction in cell density occurred in algae exposed to 32% and 56% raw groundwater by volume (see Tables A16-1, A16-2 and A16-3). The NOEC and LOEC for reduction in growth are as follows:

NOEC = 18% buffered groundwater by volume.
LOEC = 32% buffered groundwater by volume.

The 96-h EC50 (reduction in growth), which was determined by the moving average angle method, is as follows:

96-h EC50 = 54.5% raw groundwater by volume (95% confidence limits = 51.06-58.52).

Table A16-1. GREEN ALGA BEACH POINT BUFFERED GROUNDWATER
TOXICITY DATA (TEST NO. 3) - MEAN CELL DENSITY
(CELLS/ML) AFTER 96 HOURS OF EXPOSURE

| Conc (% by Vol) | Rep | Mean Cell Density | | | | |
|-----------------------|-----|-------------------|-------|--------|--------|---------|
| | | 0H | 24H | 48H | 72H | 96H |
| Growth Medium | 1 | 10100 | 51900 | 278100 | 681040 | 1182860 |
| | 2 | 10100 | 54800 | 259880 | 666460 | 1159680 |
| | 3 | 10100 | 57950 | 267140 | 670680 | 1176840 |
| 18 | 1 | 10500 | 55480 | 270120 | 670000 | 1160580 |
| | 2 | 10500 | 51640 | 264120 | 663020 | 1155440 |
| | 3 | 10500 | 52860 | 261440 | 659860 | 1147640 |
| 32 | 1 | 10100 | 54800 | 251240 | 636460 | 1119080 |
| | 2 | 10100 | 50860 | 245240 | 619960 | 1097680 |
| | 3 | 10100 | 51000 | 251660 | 627680 | 1106980 |
| 56 | 1 | 10300 | 46880 | 230420 | 535400 | 805140 |
| | 2 | 10300 | 41640 | 209400 | 470840 | 714980 |
| | 3 | 10300 | 43640 | 222980 | 459240 | 702140 |
| 100 | 1 | 9800 | DEAD | | | |
| | 2 | 9800 | DEAD | | | |
| | 3 | 9800 | DEAD | | | |

Table A16-2. GREEN ALGA BEACH POINT BUFFERED GROUNDWATER
TOXICITY TEST STATISTICAL ANALYSIS (TEST NO. 3) -
MEAN CELL DENSITY (CELLS/ML)^a

Data Transformation:

None

Chi-Square Test for Normality:

| | |
|----------------------------|---|
| Calculated test statistic: | 2.51 |
| Alpha value: | 0.01 |
| Critical value: | 13.28 |
| Conclusion: | Fail to reject the null hypothesis that the data are normally distributed |

Bartlett's Test for Homogeneity of Variance:

| | |
|----------------------------|--|
| Calculated test statistic: | 9.13 |
| Alpha value: | 0.01 |
| Critical value: | 11.34 |
| Conclusion: | Fail to reject the null hypothesis that the variances are homogenous |

ANOVA:

| | |
|----------------------------|--|
| Calculated test statistic: | 144.74 |
| Alpha value: | 0.05 |
| Critical value: | 4.07 |
| Conclusion: | Reject the null hypothesis that all groups are equal |

Dunnett's Test:

| | |
|----------------------------|--|
| Calculated test statistic: | See Table A16-3 |
| Alpha value: | 0.05 |
| Critical value: | 2.42 |
| Conclusion: | Reject the null hypothesis that all groups are equal |

^a The 100% buffered Beach Point groundwater treatment was not included in the statistical analyses because all cells died during the test.

TABLE A16-3. GREEN ALGA BEACH POINT BUFFERED GROUNDWATER
TOXICITY TEST STATISTICAL ANALYSIS (TEST NO. 3) -
RESULTS OF DUNNETT'S TEST ON MEAN CELL DENSITY
(CELLS/ML)

| Conc (% by Vol) | No. of Reps | Mean Cell Density | T Statistic | Significance |
|-----------------------|-------------------|----------------------|-------------|--------------|
| Growth Medium | 3 | 1173127 | | |
| 18 | 3 | 1154553 | 0.774 | |
| 32 | 3 | 1107913 | 2.719 | * |
| 56 | 3 | 740753 | 18.024 | * |

* Significantly different at alpha = 0.05 (Dunnett's critical
value = 2.47).

APPENDIX 17

GREEN ALGAL 96-H GROWTH TEST CONDUCTED ON BEACH POINT RAW (pH \approx 4) GROUNDWATER (WELL CC-33B) (TEST NO. 4)

| | |
|-------------------------------|---|
| Test Method: | EPA/600/4-89/001 (Weber et al., 1989) |
| Type of Test: | Static |
| Date: | December 1-5, 1993 |
| Investigator: | S. D. Turley |
| Laboratory: | UMD/WREC |
| Groundwater: | |
| Source: | APG-EA Beach Point Well CC-33B |
| Chemical Characteristics: | See Appendix 48 |
| Test Medium: | Double strength "AAP" medium (Miller et al., 1978) with P added to achieve a 20:1 N:P atomic ratio |
| Test Organism: | |
| Scientific Name: | <u>Selenastrum capricornutum</u> |
| Age at Start of Test: | Log growth |
| Source: | UMD/WREC culture |
| Experimental Chambers: | |
| Material: | 250 mL glass culture flasks with cheesecloth/cotton stoppers |
| Test Solution Volume: | 100 mL |
| Initial Cell Density: | 1×10^4 cells/mL |
| No. Replicates per Treatment: | 3 |
| Lighting: | Fluorescent; cool white; continuous; \approx 300 foot candles |
| Shaking Rate: | 100 cpm continuously |
| Endpoint: | Reduction in growth relative to control |

Temperature:

25 ± 0.2 °C

Results:

100% raw Beach Point groundwater killed all algal cells during the first 24 h of exposure (Table A17-1). A significant ($\alpha = 0.05$) reduction in cell density occurred in algae exposed to 32% and 56% raw groundwater by volume (see Tables A17-1, A17-2 and A17-3). The NOEC and LOEC for reduction in growth are as follows:

NOEC = 18% raw groundwater by volume.

LOEC = 32% raw groundwater by volume.

The 96-h EC50 (reduction in growth), which was determined by the moving average angle method, is as follows:

96-h EC50 = 54.6% raw groundwater by volume (95% confidence limits = 51.24-58.51).

TABLE A17-1. GREEN ALGA BEACH POINT RAW GROUNDWATER TOXICITY
DATA (TEST NO. 4) - MEAN CELL DENSITY (CELLS/ML)
AFTER 96 HOURS OF EXPOSURE

| Conc (% by Vol) | Rep | Mean Cell Density | | | | |
|----------------------------|-----|-------------------|-------|--------|--------|---------|
| | | 0H | 24H | 48H | 72H | 96H |
| Growth Medium | 1 | 10400 | 66800 | 270240 | 570840 | 1510840 |
| | 2 | 10400 | 58660 | 278880 | 582040 | 1601000 |
| | 3 | 10400 | 56660 | 264660 | 560220 | 1496460 |
| APG-EA Diluent Water | 1 | 10600 | 59880 | 276880 | 606800 | 1580000 |
| | 2 | 10600 | 59600 | 265840 | 579980 | 1557980 |
| | 3 | 10600 | 56480 | 269480 | 569440 | 1503440 |
| 10 | 1 | 10860 | 70000 | 288440 | 590080 | 1622000 |
| | 2 | 10860 | 58800 | 270240 | 576400 | 1509460 |
| | 3 | 10860 | 60240 | 264240 | 280260 | 1501200 |
| 18 | 1 | 11100 | 62240 | 270760 | 577240 | 1468960 |
| | 2 | 11100 | 64200 | 266240 | 570200 | 1555800 |
| | 3 | 11100 | 66380 | 260080 | 560000 | 1548980 |
| 32 | 1 | 9880 | 54980 | 230880 | 538080 | 1380640 |
| | 2 | 9880 | 55640 | 241680 | 550680 | 1478000 |
| | 3 | 9880 | 59880 | 226320 | 529400 | 1398240 |
| 56 | 1 | 12100 | 50400 | 200420 | 486260 | 999880 |
| | 2 | 12100 | 55600 | 222240 | 499780 | 1112660 |
| | 3 | 12100 | 57680 | 199980 | 460560 | 1001240 |
| 100 | 1 | 10000 | DEAD | | | |
| | 2 | 10000 | DEAD | | | |
| | 3 | 10000 | DEAD | | | |

Table A17-2. GREEN ALGA BEACH POINT RAW GROUNDWATER TOXICITY
TEST STATISTICAL ANALYSIS (TEST NO. 4) - MEAN CELL
DENSITY (CELLS/ML)^a

Data Transformation:

None

Chi-Square Test for Normality:

| | |
|----------------------------|---|
| Calculated test statistic: | 9.25 |
| Alpha value: | 0.01 |
| Critical value: | 13.28 |
| Conclusion: | Fail to reject the null hypothesis that the data are normally distributed |

Bartlett's Test for Homogeneity of Variances:

| | |
|----------------------------|--|
| Calculated test statistic: | 0.63 |
| Alpha value: | 0.01 |
| Critical value: | 15.09 |
| Conclusion: | Fail to reject the null hypothesis that the variances are homogenous |

ANOVA:

| | |
|----------------------------|--|
| Calculated test statistic: | 38.98 |
| Alpha value: | 0.05 |
| Critical value: | 3.11 |
| Conclusion: | Reject the null hypothesis that all groups are equal |

Dunnett's Test:

| | |
|----------------------------|--|
| Calculated test statistic: | See Table A17-3 |
| Alpha value: | 0.05 |
| Critical value: | 2.50 |
| Conclusion: | Reject the null hypothesis that all groups are equal |

^a The 100% raw Beach Point groundwater treatment was not included in the statistical analyses because all cells died during the test.

Table A17-3. GREEN ALGA BEACH POINT RAW GROUNDWATER TOXICITY
TEST STATISTICAL ANALYSIS (TEST NO. 4) - RESULTS OF
DUNNETT'S TEST ON MEAN CELL DENSITY (CELLS/ML)

| Conc (% by Vol) | No. of Reps | Mean Cell Density | T Statistic | Significance |
|----------------------------|-------------------|----------------------|-------------|--------------|
| Growth Medium | 3 | 1536100 | | |
| APG-EA Diluent Water | 3 | 1547140 | -0.243 | |
| 10 | 3 | 1544220 | -0.179 | |
| 18 | 3 | 1524580 | 0.254 | |
| 32 | 3 | 1418960 | 2.582 | * |
| 56 | 3 | 1037927 | 10.979 | * |

* Significantly different at alpha = 0.05 (Dunnett's critical
value = 2.50).

APPENDIX 18

GREEN ALGAL 96-H GROWTH TEST CONDUCTED ON BEACH POINT BUFFERED (pH \approx 7) GROUNDWATER (WELL CC-33B) (TEST NO. 4)

| | |
|-------------------------------|---|
| Test Method: | EPA/600/4-89/001 (Weber et al., 1989) |
| Type of Test: | Static |
| Date: | December 1-5, 1993 |
| Investigator: | S. D. Turley |
| Laboratory: | UMD/WREC |
| Groundwater: | |
| Source: | APG-EA Beach Point Well CC-33B |
| Chemical Characteristics: | See Appendix 48 |
| Test Medium: | Double strength "AAP" medium (Miller et al., 1978) with P added to achieve a 20:1 N:P atomic ratio |
| Test Organism: | |
| Scientific Name: | <u>Selenastrum capricornutum</u> |
| Age at Start of Test: | Log growth |
| Source: | UMD/WREC culture |
| Experimental Chambers: | |
| Material: | 250 mL glass culture flasks with cheesecloth/cotton stoppers |
| Test Solution Volume: | 100 mL |
| Initial Cell Density: | 1×10^4 cells/mL |
| No. Replicates per Treatment: | 3 |
| Lighting: | Fluorescent; cool white; continuous; \approx 300 foot candles |
| Shaking Rate: | 100 cpm continuously |
| pH Buffer: | 10 N NaOH |

| | |
|--------------|---|
| Endpoint: | Reduction in growth relative to control |
| Temperature: | 25 ± 0.2 °C |

Results:

100% buffered Beach Point groundwater killed all algal cells during the first 24 h of exposure (Table A18-1). A significant ($\alpha = 0.05$) reduction in cell density occurred in algae exposed to 32% and 56% raw groundwater by volume (see Tables A18-1, A18-2 and A18-3). The NOEC and LOEC for reduction in growth are as follows:

NOEC = 18% buffered groundwater by volume.
LOEC = 32% buffered groundwater by volume.

The 96-h EC50 (reduction in growth), which was determined by the moving average angle method, is as follows:

96-h EC50 = 49.7% raw groundwater by volume (95% confidence limits = 46.56-53.22).

Table A18-1. GREEN ALGA BEACH POINT BUFFERED GROUNDWATER
TOXICITY DATA (TEST NO. 4) - MEAN CELL DENSITY
(CELLS/ML) AFTER 96 HOURS OF EXPOSURE

| Conc (% by Vol) | Rep | Mean Cell Density | | | | |
|-----------------------|-----|-------------------|-------|--------|--------|---------|
| | | 0H | 24H | 48H | 72H | 96H |
| Growth Medium | 1 | 10400 | 66800 | 270240 | 570840 | 1510840 |
| | 2 | 10400 | 58660 | 278880 | 582040 | 1601000 |
| | 3 | 10400 | 56660 | 264660 | 560220 | 1496460 |
| 10 | 1 | 10400 | 63000 | 254240 | 555460 | 1481300 |
| | 2 | 10400 | 60360 | 248860 | 526780 | 1468340 |
| | 3 | 10400 | 62760 | 259980 | 566960 | 1444440 |
| 18 | 1 | 12400 | 66000 | 268860 | 587240 | 1561000 |
| | 2 | 12400 | 60780 | 265660 | 570240 | 1460000 |
| | 3 | 12400 | 57700 | 260000 | 548480 | 1448660 |
| 32 | 1 | 10420 | 57000 | 222300 | 536240 | 1219920 |
| | 2 | 10420 | 59980 | 209240 | 538000 | 1267080 |
| | 3 | 10420 | 61000 | 231880 | 521000 | 1300480 |
| 56 | 1 | 9680 | 50240 | 196880 | 451240 | 898200 |
| | 2 | 9680 | 58140 | 190180 | 455980 | 918460 |
| | 3 | 9680 | 59000 | 189040 | 425000 | 953440 |
| 100 | 1 | 9800 | DEAD | | | |
| | 2 | 9800 | DEAD | | | |
| | 3 | 9800 | DEAD | | | |

Table A18-2. GREEN ALGA BEACH POINT BUFFERED GROUNDWATER
TOXICITY TEST STATISTICAL ANALYSIS (TEST NO. 4) -
MEAN CELL DENSITY (CELLS/ML)^a

Data Transformation:

None

Chi-Square Test for Normality:

| | |
|----------------------------|---|
| Calculated test statistic: | 3.13 |
| Alpha value: | 0.01 |
| Critical value: | 13.28 |
| Conclusion: | Fail to reject the null hypothesis that the data are normally distributed |

Bartlett's Test for Homogeneity of Variance:

| | |
|----------------------------|--|
| Calculated test statistic: | 2.77 |
| Alpha value: | 0.01 |
| Critical value: | 13.28 |
| Conclusion: | Fail to reject the null hypothesis that the variances are homogenous |

ANOVA:

| | |
|----------------------------|--|
| Calculated test statistic: | 97.90 |
| Alpha value: | 0.05 |
| Critical value: | 3.48 |
| Conclusion: | Reject the null hypothesis that all groups are equal |

Dunnett's Test:

| | |
|----------------------------|--|
| Calculated test statistic: | See Table A18-3 |
| Alpha value: | 0.05 |
| Critical value: | 2.47 |
| Conclusion: | Reject the null hypothesis that all groups are equal |

^a The 100% buffered Beach Point groundwater treatment was not included in the statistical analyses because all cells died during the test.

TABLE A18-3. GREEN ALGA BEACH POINT BUFFERED GROUNDWATER
TOXICITY TEST STATISTICAL ANALYSIS (TEST NO. 4) -
RESULTS OF DUNNETT'S TEST ON MEAN CELL DENSITY
(CELLS/ML)

| Conc (% by Vol) | No. of Reps | Mean Cell Density | T Statistic | Significance |
|-----------------------|-------------------|----------------------|-------------|--------------|
| Growth Medium | 3 | 1536100 | | |
| 10 | 3 | 1464693 | 1.975 | |
| 18 | 3 | 1489887 | 1.278 | |
| 32 | 3 | 1262493 | 7.567 | * |
| 56 | 3 | 923367 | 16.947 | * |

* Significantly different at $\alpha = 0.05$ (Dunnett's critical value = 2.47).

APPENDIX 19

CLADOCERAN ACUTE AND 7-DAY SURVIVAL AND REPRODUCTION TEST CONDUCTED ON BEACH POINT RAW (pH \approx 4) GROUNDWATER (WELL CC-33B) (TEST NO. 1)

| | |
|---------------------------|--|
| Test Method: | EPA/600/4-89/001 (Weber et al., 1989) |
| Type of Test: | Static renewal (every 24 h) |
| Date: | April 7-14, 1993 |
| Investigator: | S.D. Turley |
| Laboratory: | UMD/WREC |
| Groundwater: | |
| Source: | APG-EA Beach Point Well CC-33B |
| Chemical Characteristics: | See Appendix 48 |
| Dilution Water: | |
| Source: | 20% Perrier: 80% RO water |
| Chemical Characteristics: | See Table A19-1 |
| Test Organism: | |
| Scientific Name: | <u>Ceriodaphnia dubia</u> |
| Age at Start of Test: | <4 h |
| Source: | UMD/WREC culture |
| Experimental Chambers: | |
| Material: | 50 mL glass beaker |
| Test Solution Volume: | 25 mL |
| No. Organisms/Replicate: | 1 |
| No. Organisms/Treatment: | 10 |
| Loading: | 1 organism/beaker |
| Lighting: | Fluorescent; 60-85 foot candles |
| Aeration: | Prior to each renewal |

Endpoints:

Mortality of adults; number of neonates produced in 3 broods

Water Quality:

Table A19-1

Results:

Mortality:

48-h Exposure:

The data for the 48-h LC50 were obtained from the 7-d study. All organisms exposed to 100% raw groundwater died during the 48-h exposure (Table A19-2). The organisms in all other treatments lived. The 48-h LC50, which was determined by the moving average angle method, is as follows:

48-h LC50 = 64.7% raw groundwater by volume (95% confidence limits = 57.17-75.48).

7-d Exposure:

All organisms exposed to 100% raw groundwater died during the 7-d exposure (Table A19-3); organisms in all other treatments lived. The 7-d LC50, which was determined by the moving average angle method, is as follows:

7-d LC50 = 64.7% raw groundwater by volume (95% confidence limits = 57.17-75.48).

Neonate Production:

Raw groundwater significantly ($\alpha = 0.05$) reduced neonate production relative to the controls down to 32% raw groundwater by volume (see Tables A19-3, A19-4, and A19-5). The NOEC and LOEC for the daphnids, based on reduced neonate production, are as follows:

NOEC = 18% raw groundwater by volume.

LOEC = 32% raw groundwater by volume.

TABLE A19-1. SUMMARY OF THE BEACH POINT RAW GROUNDWATER BIOASSAY
WATER QUALITY DATA FOR THE CLADOCERAN 7-DAY TEST
(TEST NO. 1) - DISSOLVED OXYGEN (MG/L)

| | <u>Test Concentrations (Percent Groundwater by Volume)</u> | | | | | |
|--------------|--|-----|-----|-----|-----|-----|
| | 0 | 10 | 18 | 32 | 56 | 100 |
| <u>Day 0</u> | | | | | | |
| 0 H | 8.8 | 8.7 | 8.6 | 8.3 | 8.2 | 8.2 |
| <u>Day 1</u> | | | | | | |
| 0 H | 8.2 | 8.1 | 8.1 | 8.1 | 8.1 | 5.6 |
| 24 H | 8.1 | 8.1 | 8.1 | 7.8 | 7.8 | |
| <u>Day 2</u> | | | | | | |
| 0 H | 8.2 | 8.1 | 8.1 | 8.1 | 8.1 | |
| 24 H | 7.6 | 7.6 | 7.6 | 7.6 | 7.6 | |
| <u>Day 3</u> | | | | | | |
| 0 H | 8.4 | 8.2 | 8.1 | 8.2 | 8.1 | |
| 24 H | 7.6 | 7.5 | 7.5 | 7.5 | 7.5 | |
| <u>Day 4</u> | | | | | | |
| 0 H | 8.4 | 8.2 | 8.2 | 8.2 | 8.0 | |
| 24 H | 8.5 | 8.5 | 8.6 | 8.6 | 8.6 | |
| <u>Day 5</u> | | | | | | |
| 0 H | 8.3 | 8.3 | 8.3 | 8.1 | 8.1 | |
| 24 H | 8.5 | 8.4 | 8.4 | 8.5 | 8.5 | |
| <u>Day 6</u> | | | | | | |
| 0 H | 8.5 | 8.1 | 8.2 | 8.0 | 8.0 | |
| 24 H | 8.4 | 8.2 | 8.4 | 8.6 | 8.6 | |
| <u>Day 7</u> | | | | | | |
| 24 H | 8.1 | 8.2 | 8.1 | 8.1 | 8.1 | |

TABLE A19-1. (CONTINUED) - pH (STANDARD UNITS)

| | <u>Test Concentrations (Percent Groundwater by Volume)</u> | | | | | |
|--------------|--|------|------|------|------|------|
| | 0 | 10 | 18 | 32 | 56 | 100 |
| <u>Day 0</u> | | | | | | |
| 0 H | 6.83 | 6.81 | 6.81 | 6.71 | 5.93 | 3.93 |
| <u>Day 1</u> | | | | | | |
| 0 H | 6.68 | 6.77 | 6.81 | 6.70 | 6.84 | 3.74 |
| 24 H | 6.63 | 6.81 | 6.98 | 6.80 | 6.72 | |
| <u>Day 2</u> | | | | | | |
| 0 H | 6.75 | 6.79 | 6.82 | 6.77 | 6.26 | |
| 24 H | 6.52 | 6.71 | 6.91 | 7.02 | 7.08 | |
| <u>Day 3</u> | | | | | | |
| 0 H | 6.69 | 6.32 | 6.28 | 6.27 | 6.23 | |
| 24 H | 6.57 | 6.51 | 6.61 | 6.78 | 6.80 | |
| <u>Day 4</u> | | | | | | |
| 0 H | 6.49 | 6.61 | 6.64 | 6.62 | 6.56 | |
| 24 H | 6.38 | 6.53 | 6.67 | 6.82 | 6.91 | |
| <u>Day 5</u> | | | | | | |
| 0 H | 7.00 | 7.07 | 6.93 | 6.63 | 6.17 | |
| 24 H | 7.04 | 6.99 | 7.03 | 7.10 | 7.11 | |
| <u>Day 6</u> | | | | | | |
| 0 H | 7.03 | 6.99 | 6.88 | 6.71 | 6.28 | |
| 24 H | 7.09 | 6.87 | 7.01 | 7.01 | 7.00 | |
| <u>Day 7</u> | | | | | | |
| 24 H | 6.71 | 6.74 | 6.87 | 6.92 | 6.94 | |

TABLE A19-1. (CONTINUED) - CONDUCTIVITY (μ MHOS/CM)

| <u>Test Concentrations (Percent Groundwater by Volume)</u> | | |
|--|----------|------------|
| | <u>0</u> | <u>100</u> |
| <u>Day 0</u> | | |
| 0 H | 140 | 1500 |
| <u>Day 1</u> | | |
| 0 H | 170 | 1600 |
| <u>Day 2</u> | | |
| 0 H | 155 | |
| <u>Day 3</u> | | |
| 0 H | 155 | |
| <u>Day 4</u> | | |
| 0 H | 165 | |
| <u>Day 5</u> | | |
| 0 H | 160 | |
| <u>Day 6</u> | | |
| 0 H | 165 | |
| <u>Day 7</u> | | |
| 24 H | 158 | |

TABLE A19-1. (CONTINUED) - ALKALINITY (MG/L AS CaCO₃)

| | <u>Test Concentrations (Percent Groundwater by Volume)</u> | |
|--------------|--|--------------|
| | 0 | 100 |
| <u>Day 0</u> | | |
| 0 H | 50 | ^a |
| <u>Day 1</u> | | |
| 0 H | 50 | ^a |
| <u>Day 2</u> | | |
| 0 H | 50 | |
| <u>Day 3</u> | | |
| 0 H | 50 | |
| <u>Day 4</u> | | |
| 0 H | 50 | |
| <u>Day 5</u> | | |
| 0 H | 50 | |
| <u>Day 6</u> | | |
| 0 H | 50 | |
| <u>Day 7</u> | | |
| 24 H | 50 | |

^a Could not obtain measurement.

TABLE A19-1. (CONTINUED) - HARDNESS (MG/L AS CaCO₃)

| <u>Test Concentrations (Percent Groundwater by Volume)</u> | | |
|--|----------|------------|
| | <u>0</u> | <u>100</u> |
| <hr/> | | |
| <u>Day 0</u> | | |
| 0 H | 48 | a |
| <u>Day 1</u> | | |
| 0 H | 46 | a |
| <u>Day 2</u> | | |
| 0 H | 56 | |
| <u>Day 3</u> | | |
| 0 H | 50 | |
| <u>Day 4</u> | | |
| 0 H | 52 | |
| <u>Day 5</u> | | |
| 0 H | 54 | |
| <u>Day 6</u> | | |
| 0 H | 54 | |
| <u>Day 7</u> | | |
| 24 H | 51 | |

^a Could not obtain measurement.

TABLE A19-2. CLADOCERAN BEACH POINT RAW GROUNDWATER TOXICITY
TEST DATA (TEST NO. 1) - SURVIVAL AFTER 48 HOURS OF
EXPOSURE

| Concentration (% Groundwater by Volume) | Number Tested | No. Alive at 48 Hours | Percent Alive |
|---|------------------|--------------------------|------------------|
| UMD/WREC Control | 10 | 10 | 100 |
| 10 | 10 | 10 | 100 |
| 18 | 10 | 10 | 100 |
| 32 | 10 | 10 | 100 |
| 56 | 10 | 10 | 100 |
| 100 | 10 | 0 | 0 |

TABLE A19-3. CLADOCERAN BEACH POINT RAW GROUNDWATER TOXICITY DATA (TEST NO. 1) - SURVIVAL OF ADULTS, NUMBER OF YOUNG PRODUCED PER BROOD, TOTAL NUMBER OF YOUNG, AND MEAN NUMBER OF YOUNG PER BROOD AFTER 7 DAYS OF EXPOSURE

| Conc (% by Vol) | Rep | Brood No. 1 | Brood No. 2 | Brood No. 3 | Total Young | Mean Young Per Brood |
|----------------------------|-----|----------------|----------------|----------------|----------------|-------------------------|
| UMD/WREC Control | 1 | 6 | 13 | 13 | 32 | 10.7 |
| | 2 | 4 | 12 | 12 | 28 | 9.3 |
| | 3 | 7 | 11 | 11 | 29 | 9.7 |
| | 4 | 4 | 10 | 11 | 25 | 8.3 |
| | 5 | 3 | 14 | 13 | 30 | 10.0 |
| | 6 | 5 | 7 | 13 | 25 | 8.3 |
| | 7 | 7 | 11 | 11 | 29 | 9.7 |
| | 8 | | | | | |
| | 9 | a 4 | 14 | 15 | 33 | 11.0 |
| | 10 | 5 | 12 | 11 | 28 | 9.3 |
| APG-EA Diluent Water | 1 | 5 | 11 | 12 | 28 | 9.3 |
| | 2 | 4 | 10 | 13 | 27 | 9.0 |
| | 3 | 2 | 9 | 14 | 25 | 8.3 |
| | 4 | 5 | 9 | 14 | 28 | 9.3 |
| | 5 | 4 | 8 | 12 | 24 | 8.0 |
| | 6 | 5 | 10 | 11 | 26 | 8.7 |
| | 7 | 6 | 9 | 12 | 27 | 9.0 |
| | 8 | 6 | 10 | 11 | 27 | 9.0 |
| | 9 | 5 | 10 | 13 | 28 | 9.3 |
| | 10 | 5 | 11 | 15 | 31 | 10.3 |
| 10 | 1 | 7 | 10 | 13 | 30 | 10.0 |
| | 2 | 7 | 11 | 10 | 28 | 9.3 |
| | 3 | 6 | 12 | 10 | 28 | 9.3 |
| | 4 | 6 | 11 | 9 | 26 | 8.7 |
| | 5 | 5 | 11 | 9 | 25 | 8.3 |
| | 6 | 7 | 11 | 11 | 29 | 9.7 |
| | 7 | 4 | 8 | 9 | 21 | 7.0 |
| | 8 | 7 | 10 | 12 | 29 | 9.7 |
| | 9 | 5 | 11 | 12 | 28 | 9.3 |
| | 10 | 7 | 11 | 11 | 29 | 9.7 |

TABLE A19-3. (CONTINUED)

| Conc (% by Vol) | Rep | Brood No. 1 | Brood No. 2 | Brood No. 3 | Total Young | Mean Young Per Brood |
|-----------------------|-----|----------------|----------------|----------------|----------------|-------------------------|
| 18 | 1 | 5 | 10 | 8 | 23 | 7.7 |
| | 2 | a | | | | |
| | 3 | 6 | 9 | 8 | 23 | 7.7 |
| | 4 | 7 | 9 | 10 | 26 | 8.7 |
| | 5 | a | | | | |
| | 6 | 6 | 11 | 12 | 29 | 9.7 |
| | 7 | 6 | 11 | 12 | 29 | 9.7 |
| | 8 | 6 | 10 | 10 | 26 | 8.7 |
| | 9 | 6 | 8 | 10 | 24 | 8.0 |
| | 10 | 4 | 6 | 11 | 21 | 7.0 |
| 32 | 1 | 7 | 9 | 12 | 28 | 9.3 |
| | 2 | 7 | 0 | 9 | 16 | 8.0 |
| | 3 | 5 | 9 | 10 | 24 | 8.0 |
| | 4 | 6 | 9 | 9 | 24 | 8.0 |
| | 5 | 6 | 6 | 4 | 16 | 5.3 |
| | 6 | 4 | 7 | 8 | 19 | 6.3 |
| | 7 | 7 | 10 | 10 | 27 | 9.0 |
| | 8 | 6 | 9 | 11 | 26 | 8.7 |
| | 9 | 5 | 8 | 5 | 18 | 6.0 |
| | 10 | 6 | 8 | 7 | 21 | 7.0 |
| 56 | 1 | 2 | 3 | 12 | 17 | 5.7 |
| | 2 | 2 | 4 | 4 | 10 | 3.3 |
| | 3 | 4 | 3 | 7 | 14 | 4.7 |
| | 4 | 4 | 6 | 7 | 17 | 5.7 |
| | 5 | 5 | 9 | 8 | 22 | 7.3 |
| | 6 | 4 | 6 | 8 | 18 | 6.0 |
| | 7 | 5 | 6 | 8 | 19 | 6.3 |
| | 8 | 4 | 5 | 10 | 19 | 6.3 |
| | 9 | 4 | 4 | 3 | 11 | 3.7 |
| | 10 | 5 | 5 | 6 | 16 | 5.3 |

TABLE A19-3. (CONTINUED)

| Conc (% by Vol) | Rep | Brood No. 1 | Brood No. 2 | Brood No. 3 | Total Young | Mean Young Per Brood |
|-----------------------|-----|----------------|----------------|----------------|----------------|-------------------------|
| 100 | 1 | DEAD | | | | |
| | 2 | DEAD | | | | |
| | 3 | DEAD | | | | |
| | 4 | DEAD | | | | |
| | 5 | DEAD | | | | |
| | 6 | DEAD | | | | |
| | 7 | DEAD | | | | |
| | 8 | DEAD | | | | |
| | 9 | DEAD | | | | |
| | 10 | DEAD | | | | |

^a Male adult, which was included in survival analysis, was not included in the neonate production analysis.

TABLE A19-4. CLADOCERAN BEACH POINT RAW GROUNDWATER TOXICITY
TEST STATISTICAL ANALYSIS (TEST NO. 1) - NEONATE
PRODUCTION AFTER 7 DAYS OF EXPOSURE^a

Data Transformation:

None

Chi-square Test for Normality:

| | |
|----------------------------|---|
| Calculated test statistic: | 1.07 |
| Alpha value: | 0.01 |
| Critical value: | 13.28 |
| Conclusion | Fail to reject the null hypothesis that the data are normally distributed |

Bartlett's Test for Homogeneity of Variances:

| | |
|----------------------------|--|
| Calculated test statistic: | 6.97 |
| Alpha value: | 0.01 |
| Critical value: | 15.09 |
| Conclusion: | Fail to reject the null hypothesis that the variances are homogenous |

ANOVA:

| | |
|----------------------------|--|
| Calculated test statistic: | 20.54 |
| Alpha value: | 0.05 |
| Critical value: | 2.45 |
| Conclusion: | Reject the null hypothesis that all groups are equal |

Bonferroni's T-test:

| | |
|-----------------------------|--|
| Calculated test statistics: | See Table A19-5 |
| Alpha value: | 0.05 |
| Critical value: | 2.40 |
| Conclusion: | Reject the null hypothesis that all groups are equal |

^a The 100% raw Beach Point groundwater treatment was not included in the statistical analyses because all organisms died during the test.

TABLE A19-5. CLADOCERAN BEACH POINT RAW GROUNDWATER TOXICITY
TEST STATISTICAL ANALYSIS (TEST NO. 1) - RESULTS OF
BONFERRONI'S T-TEST ON MEAN NEONATE PRODUCTION
AFTER 7 DAYS OF EXPOSURE

| Conc (% by Vol) | No. of Reps | Mean No. Neonates Produced | T Statistic | Significance |
|----------------------------|-------------------|----------------------------------|-------------|--------------|
| UMD/WREC Control | 9 | 28.8 | | |
| APG-EA Diluent Water | 10 | 27.1 | 1.14 | |
| 10 | 10 | 27.3 | 1.01 | |
| 18 | 8 | 25.1 | 2.35 | |
| 32 | 10 | 21.9 | 4.68 | * |
| 56 | 10 | 16.3 | 8.49 | * |

* Significantly different at $\alpha = 0.05$ (Bonferroni's critical value = 2.40).

APPENDIX 20

CLADOCERAN ACUTE AND 7-DAY SURVIVAL AND REPRODUCTION TEST CONDUCTED ON BEACH POINT BUFFERED (pH \approx 7) GROUNDWATER (WELL CC-33B) (TEST NO. 1)

| | |
|---------------------------|--|
| Test Method: | EPA/600/4-89/001 (Weber et al., 1989) |
| Type of Test: | Static renewal (every 24 h) |
| Date: | April 7-14, 1993 |
| Investigator: | S. D. Turley |
| Laboratory: | UMD/WREC |
| Groundwater: | |
| Source: | APG-EA Beach Point Well CC-33B |
| Chemical Characteristics: | See Appendix 48 |
| Dilution Water: | |
| Source: | 20% Perrier; 80% RO water |
| Chemical Characteristics: | See Table A20-1 |
| Test Organism: | |
| Scientific Name: | <u>Ceriodaphnia dubia</u> |
| Age at Start of Test: | <4 h |
| Source: | UMD/WREC culture |
| Experimental Chambers: | |
| Material: | 50 mL glass beaker |
| Test Solution Volume: | 25 mL |
| No. Organisms/Replicate: | 1 |
| No. Organisms/Treatment: | 10 |
| Loading: | 1 organism/beaker |
| Lighting: | Fluorescent; 60-85 foot candles |
| Aeration: | Prior to each renewal |
| pH Buffer: | 10 N NaOH |

Endpoints:

Mortality of adults; number of neonates produced in 3 broods

Water Quality:

Table A20-1

Results:

Mortality:

48-h Exposure:

The data for the 48-h LC50 were obtained from the 7-d study. The buffered groundwater did not affect larval survival. The data are summarized in Table A20-2.

7-d Exposure:

Significant ($\alpha = 0.05$) mortality occurred to the daphnids exposed to 100% and 56% buffered groundwater by volume; statistically significant mortality did not occur at 32% buffered groundwater by volume (see Tables A20-3, A20-4, and A20-5). The 7-d LC50 for adult mortality, which was determined by the moving average angle method, is as follows:

7-d LC50 = 43.3% buffered groundwater by volume (95% confidence limits = 1.00-56.42).

Neonate Production:

A significant ($\alpha = 0.001$) reduction in neonate production relative to the controls occurred at 32% buffered groundwater by volume, which was the lowest concentration studied (see Tables A20-3 and A20-6). The NOEC and LOEC for exposure of the organism to buffered Beach Point groundwater could not be determined because a significant reduction in neonate production occurred at the lowest concentration studied. A comparison of the daphnid toxicity results between the raw and buffered Beach Point groundwater suggests that the buffered groundwater may have the same toxicity or may be more toxic than the raw groundwater. For example, the 7-d LC50 for mortality of the adults in the buffered groundwater was 43.3% groundwater by volume in contrast to 59.5% groundwater by volume in the raw groundwater. Neonate production was reduced in both sources down to 32% groundwater by volume. The reason(s) for this finding is not clear; however, it may be related to the fact that during the test, material precipitated out of solution which appeared to interfere with the daphnids as they foraged for food on the bottom of the exposure vessels.

TABLE A20-1. SUMMARY OF THE BEACH POINT BUFFERED GROUNDWATER
BIOASSAY WATER QUALITY DATA FOR THE CLADOCERAN 7-
DAY TEST (TEST NO. 1) - DISSOLVED OXYGEN (MG/L)

| | <u>Test Concentrations (Percent Groundwater by Volume)</u> | | | |
|--------------|--|-----|-----|-----|
| | 0 | 32 | 56 | 100 |
| <u>Day 0</u> | | | | |
| 0 H | 8.8 | 7.8 | 8.1 | 8.2 |
| <u>Day 1</u> | | | | |
| 0 H | 8.2 | 7.9 | 7.9 | 5.6 |
| 24 H | 8.1 | 8.1 | 8.1 | 8.0 |
| <u>Day 2</u> | | | | |
| 0 H | 8.2 | 8.1 | 8.0 | 6.6 |
| 24 H | 7.6 | 7.6 | 7.6 | 7.6 |
| <u>Day 3</u> | | | | |
| 0 H | 8.4 | 8.1 | 8.1 | 6.8 |
| 24 H | 7.6 | 7.5 | 7.5 | 7.5 |
| <u>Day 4</u> | | | | |
| 0 H | 8.4 | 8.2 | 8.0 | 6.2 |
| 24 H | 8.5 | 8.4 | 8.2 | 8.3 |
| <u>Day 5</u> | | | | |
| 0 H | 8.3 | 8.1 | 8.1 | 5.1 |
| 24 H | 8.5 | 8.3 | 8.1 | 8.2 |
| <u>Day 6</u> | | | | |
| 0 H | 8.5 | 8.0 | 8.1 | 5.9 |
| 24 H | 8.4 | 8.2 | 8.2 | 8.3 |
| <u>Day 7</u> | | | | |
| 24 H | 8.1 | 8.0 | 8.0 | 8.0 |

TABLE A20-1. (CONTINUED) - pH (STANDARD UNITS)

| | | <u>Test Concentrations (Percent Groundwater by Volume)</u> | | | |
|--------------|--|--|------|------|------|
| | | 0 | 32 | 56 | 100 |
| <u>Day 0</u> | | | | | |
| 0 H | | 6.83 | 6.88 | 6.97 | 3.93 |
| <u>Day 1</u> | | | | | |
| 0 H | | 6.68 | 6.92 | 7.03 | 6.69 |
| 24 H | | 6.63 | 7.23 | 7.36 | 7.52 |
| <u>Day 2</u> | | | | | |
| 0 H | | 6.75 | 7.18 | 7.24 | 7.05 |
| 24 H | | 6.52 | 7.19 | 7.43 | 7.54 |
| <u>Day 3</u> | | | | | |
| 0 H | | 6.69 | 6.88 | 6.89 | 6.98 |
| 24 H | | 6.57 | 6.94 | 7.10 | 7.36 |
| <u>Day 4</u> | | | | | |
| 0 H | | 6.49 | 6.80 | 6.97 | 7.06 |
| 24 H | | 6.38 | 7.09 | 7.20 | 7.37 |
| <u>Day 5</u> | | | | | |
| 0 H | | 7.00 | 6.59 | 6.67 | 7.00 |
| 24 H | | 7.04 | 7.19 | 7.22 | 7.42 |
| <u>Day 6</u> | | | | | |
| 0 H | | 7.03 | 6.63 | 6.81 | 7.01 |
| 24 H | | 7.09 | 7.08 | 7.14 | 7.36 |
| <u>Day 7</u> | | | | | |
| 24 H | | 6.71 | 7.12 | 7.24 | 7.43 |

TABLE A20-1. (CONTINUED) - CONDUCTIVITY (μ MHOS/CM)

| | <u>Test Concentrations (Percent Groundwater by Volume)</u> | |
|--------------|--|------------|
| | <u>0</u> | <u>100</u> |
| <u>Day 0</u> | | |
| 0 H | 140 | 1700 |
| <u>Day 1</u> | | |
| 0 H | 170 | 1700 |
| <u>Day 2</u> | | |
| 0 H | 155 | 1700 |
| <u>Day 3</u> | | |
| 0 H | 155 | 1700 |
| <u>Day 4</u> | | |
| 0 H | 165 | 1750 |
| <u>Day 5</u> | | |
| 0 H | 160 | 1800 |
| <u>Day 6</u> | | |
| 0 H | 165 | 1750 |
| <u>Day 7</u> | | |
| 24 H | 158 | 1729 |

TABLE A20-1. (CONTINUED) - ALKALINITY (MG/L AS CaCO₃)

| <u>Test Concentrations (Percent Groundwater by Volume)</u> | | |
|--|----------|------------|
| | <u>0</u> | <u>100</u> |
| <hr/> | | |
| <u>Day 0</u> | | |
| 0 H | 50 | 80 |
| <u>Day 1</u> | | |
| 0 H | 50 | 80 |
| <u>Day 2</u> | | |
| 0 H | 50 | 85 |
| <u>Day 3</u> | | |
| 0 H | 50 | 85 |
| <u>Day 4</u> | | |
| 0 H | 50 | 100 |
| <u>Day 5</u> | | |
| 0 H | 50 | 85 |
| <u>Day 6</u> | | |
| 0 H | 50 | 85 |
| <u>Day 7</u> | | |
| 24 H | 50 | 85 |

TABLE A20-1. (CONTINUED) - HARDNESS (MG/L AS CaCO₃)

| | <u>Test Concentrations (Percent Groundwater by Volume)</u> | |
|--------------|--|-----|
| | 0 | 100 |
| <u>Day 0</u> | | |
| 0 H | 48 | 280 |
| <u>Day 1</u> | | |
| 0 H | 46 | 170 |
| <u>Day 2</u> | | |
| 0 H | 56 | 136 |
| <u>Day 3</u> | | |
| 0 H | 50 | 148 |
| <u>Day 4</u> | | |
| 0 H | 52 | 138 |
| <u>Day 5</u> | | |
| 0 H | 54 | 160 |
| <u>Day 6</u> | | |
| 0 H | 54 | 148 |
| <u>Day 7</u> | | |
| 24 H | 51 | 168 |

TABLE A20-2 CLADOCERAN BEACH POINT BUFFERED GROUNDWATER TOXICITY
TEST DATA (TEST NO. 1) - SURVIVAL AFTER 48 HOURS OF
EXPOSURE

| Concentration (% Groundwater by Volume) | Number Tested | No. Alive at 48 Hours | Percent Alive |
|---|------------------|--------------------------|------------------|
| UMD/WREC Control | 10 | 10 | 100 |
| 10 | 10 | 10 | 100 |
| 18 | 10 | 10 | 100 |
| 32 | 10 | 10 | 100 |
| 56 | 10 | 10 | 100 |
| 100 | 10 | 9 | 90 |

TABLE A20-3. CLADOCERAN BEACH POINT BUFFERED GROUNDWATER TOXICITY DATA (TEST NO.1) - SURVIVAL OF ADULTS, NUMBER OF YOUNG PRODUCED PER BROOD, TOTAL NUMBER OF YOUNG, AND MEAN NUMBER OF YOUNG PRODUCED PER BROOD AFTER 7 DAYS OF EXPOSURE

| Conc (% by Vol) | Rep | Brood No. 1 | Brood No. 2 | Brood No. 3 | Total Young | Mean Young Per Brood |
|-----------------------|-----|----------------|----------------|----------------|----------------|-------------------------|
| UMD/WREC Control | 1 | 6 | 13 | 13 | 32 | 10.7 |
| | 2 | 4 | 12 | 12 | 28 | 9.3 |
| | 3 | 7 | 11 | 11 | 29 | 9.7 |
| | 4 | 4 | 10 | 11 | 25 | 8.3 |
| | 5 | 3 | 14 | 13 | 30 | 10.0 |
| | 6 | 5 | 7 | 13 | 25 | 8.3 |
| | 7 | 7 | 11 | 11 | 29 | 9.7 |
| | 8 | a | | | | |
| | 9 | 4 | 14 | 15 | 33 | 11.0 |
| | 10 | 5 | 12 | 11 | 28 | 9.3 |
| 32 | 1 | 7 | 3 | DEAD | 10 | 5.0 |
| | 2 | 5 | 2 | 10 | 17 | 5.7 |
| | 3 | 7 | 10 | 6 | 23 | 7.7 |
| | 4 | a | | | | |
| | 5 | 6 | 7 | DEAD | 13 | 6.5 |
| | 6 | 5 | 5 | 4 | 14 | 4.7 |
| | 7 | 6 | 9 | 5 | 20 | 6.7 |
| | 8 | 7 | DEAD | | 7 | 7.0 |
| | 9 | 7 | 12 | 10 | 29 | 9.7 |
| | 10 | a | | | | |
| 56 | 1 | 4 | 7 | DEAD | 11 | 5.5 |
| | 2 | 4 | 3 | DEAD | 7 | 3.5 |
| | 3 | 3 | 0 | 0 | 3 | 3.0 |
| | 4 | 6 | DEAD | | 6 | 6.0 |
| | 5 | 6 | 6 | 0 | 12 | 6.0 |
| | 6 | 5 | 5 | DEAD | 10 | 5.0 |
| | 7 | 4 | DEAD | | 4 | 4.0 |
| | 8 | 3 | 0 | 0 | 3 | 3.0 |
| | 9 | 5 | 0 | 0 | 5 | 5.0 |
| | 10 | 4 | DEAD | | 4 | 4.0 |

TABLE A20-3. (CONTINUED)

| Conc (% by Vol) | Rep | Brood No. 1 | Brood No. 2 | Brood No. 3 | Total Young | Mean Young Per Brood |
|-----------------------|-----|----------------|----------------|----------------|----------------|-------------------------|
| 100 | 1 | DEAD | | | | |
| | 2 | DEAD | | | | |
| | 3 | DEAD | | | | |
| | 4 | DEAD | | | | |
| | 5 | DEAD | | | | |
| | 6 | 2 | DEAD | | 2 | 2.0 |
| | 7 | DEAD | | | | |
| | 8 | DEAD | | | | |
| | 9 | DEAD | | | | |
| | 10 | DEAD | | | | |

^a Male adult, which was included in survival analysis, was not included in the neonate production analysis.

TABLE A20-4. CLADOCERAN BEACH POINT BUFFERED GROUNDWATER
TOXICITY TEST STATISTICAL ANALYSIS (TEST NO. 1) -
ADULT SURVIVAL AFTER 7 DAYS OF EXPOSURE^a

Data Transformation:

None

Fisher's Exact Test:

| | |
|----------------------------|---|
| Calculated test statistic: | See Table A14-4 |
| Alpha value: | 0.05 |
| Critical value: | 6 |
| Conclusion: | Reject the null hypothesis that all groups are equal. |

^a The 100% buffered Beach Point groundwater treatment was not included in the statistical analyses because all organisms died during the test.

TABLE A20-5. CLADOCERAN BEACH POINT BUFFERED GROUNDWATER
TOXICITY TEST STATISTICAL ANALYSIS (TEST NO. 1) -
RESULTS OF FISHER'S EXACT TEST ON ADULT SURVIVAL
AFTER 7 DAYS OF EXPOSURE

| Conc (% by Vol) | Number Alive | Number Dead | b Value | Significance |
|-----------------------|-----------------|----------------|------------|--------------|
| UMD/WREC Control | 10 | 0 | | |
| 32 | 7 | 3 | 7 | |
| 56 | 4 | 6 | 4 | * |

* Significantly different at $\alpha = 0.05$ (Fisher's critical value = 6).

TABLE A20-6. CLADOCERAN BEACH POINT BUFFERED GROUNDWATER
TOXICITY TEST STATISTICAL ANALYSIS (TEST NO. 1) -
NEONATE PRODUCTION AFTER 7 DAYS OF EXPOSURE^a

Data Transformation:

None

Chi-Square Test for Normality:

| | |
|----------------------------|---|
| Calculated test statistic: | 1.81 |
| Alpha value: | 0.01 |
| Critical value: | 13.28 |
| Conclusion: | Fail to reject the null hypothesis that the data are normally distributed |

Bartlett's Test for Homogeneity of Variances:

| | |
|----------------------------|--|
| Calculated test statistic: | 8.24 |
| Alpha value: | 0.01 |
| Critical value: | 9.21 |
| Conclusion: | Fail to reject the null hypothesis that the variances are homogenous |

Student's T-test:

| | |
|----------------------------|--|
| Calculated test statistic: | 4.72 |
| Alpha value: | 0.001 |
| Critical value: | 4.07 |
| Conclusion: | Reject the null hypothesis that all groups are equal |

^a The 56% buffered Beach Point groundwater treatment was not included in the statistical analyses because all concentrations above the NOEC for survival are excluded from the hypothesis test for reproductive effects.

APPENDIX 21

CLADOCERAN ACUTE AND 7-DAY SURVIVAL AND REPRODUCTION
TEST CONDUCTED ON BEACH POINT RAW (pH \approx 4)
GROUNDWATER (WELL CC-33B)
(TEST NO. 2)

Test Method: EPA/600/4-89/001
(Weber et al., 1989)

Type of Test: Static renewal (every 24 h)

Date: June 1-8, 1993

Investigator: S.D. Turley

Laboratory: UMD/WREC

Groundwater:

Source: APG-EA Beach Point Well CC-33B
Chemical Characteristics: See Appendix 48

Dilution Water:

Source: 20% Perrier: 80% RO water
Chemical Characteristics: See Table A21-1

Test Organism:

Scientific Name: Ceriodaphnia dubia
Age at Start of Test: <4 h
Source: UMD/WREC culture

Experimental Chambers:

Material: 50 mL glass beaker
Test Solution Volume: 25 mL

No. Organisms/Replicate: 1

No. Organisms/Treatment: 10

Loading: 1 organism/beaker

Lighting: Fluorescent; 60-85 foot
candles

Aeration: Prior to each renewal

Endpoints: Mortality of adults; number of neonates produced in 3 broods

Water Quality: Table A21-1

Results:

Mortality:

48-h Exposure:

The data for the 48-h LC50 were obtained from the 7-d study. All organisms exposed to 100% raw groundwater died during the 48-h exposure (Table A21-2). The organisms in all other treatments lived. The 48-h LC50, which was determined by the moving average angle method, is as follows:

48-h LC50 = 64.7% raw groundwater by volume (95% confidence limits = 57.17-75.48).

7-d Exposure:

Significant ($\alpha = 0.05$) mortality occurred in the adults exposed to 100% raw groundwater (see Table A21-3). The 7-d LC50, which was determined by the moving average angle method, is as follows:

7-d LC50 = 64.7% raw groundwater by volume (95% confidence limits = 57.17-75.48).

Neonate Production:

Raw groundwater significantly ($\alpha = 0.05$) reduced neonate production relative to the controls down to 18% raw groundwater by volume (see Tables A21-3, A21-4, and A21-5). The NOEC and LOEC for the daphnids, based on reduced neonate production, are as follows:

NOEC = 10% raw groundwater by volume.

LOEC = 18% raw groundwater by volume.

TABLE A21-1. SUMMARY OF THE BEACH POINT RAW GROUNDWATER BIOASSAY
WATER QUALITY DATA FOR THE CLADOCERAN 7-DAY TEST
(TEST NO. 2) - DISSOLVED OXYGEN (MG/L)

| | | <u>Test Concentrations (Percent Groundwater by Volume)</u> | | | | | |
|--------------|--|--|-----|-----|-----|-----|-----|
| | | 0 | 10 | 18 | 32 | 56 | 100 |
| <u>Day 0</u> | | | | | | | |
| 0 H | | 8.3 | 8.3 | 8.2 | 8.0 | 7.9 | 6.8 |
| <u>Day 1</u> | | | | | | | |
| 0 H | | 8.3 | 8.3 | 8.2 | 7.9 | 7.6 | 6.6 |
| 24 H | | 8.2 | 8.0 | 8.0 | 8.0 | 8.0 | 8.1 |
| <u>Day 2</u> | | | | | | | |
| 0 H | | 8.0 | 8.0 | 7.9 | 7.4 | 7.1 | 7.2 |
| 24 H | | 8.1 | 8.0 | 7.9 | 7.9 | 7.9 | 7.7 |
| <u>Day 3</u> | | | | | | | |
| 0 H | | 8.3 | 8.0 | 8.0 | 7.9 | 7.9 | 7.1 |
| 24 H | | 7.8 | 7.4 | 7.3 | 7.2 | 7.2 | |
| <u>Day 4</u> | | | | | | | |
| 0 H | | 8.3 | 8.2 | 8.1 | 8.1 | 7.8 | |
| 24 H | | 8.3 | 8.1 | 8.1 | 7.8 | 7.8 | |
| <u>Day 5</u> | | | | | | | |
| 0 H | | 8.2 | 8.1 | 8.0 | 7.8 | 7.6 | |
| 24 H | | 8.3 | 8.2 | 8.3 | 8.0 | 8.0 | |
| <u>Day 6</u> | | | | | | | |
| 0 H | | 8.5 | 8.4 | 8.2 | 8.2 | 8.1 | |
| 24 H | | 8.2 | 8.1 | 8.1 | 8.2 | 8.1 | |
| <u>Day 7</u> | | | | | | | |
| 24 H | | 8.1 | 8.0 | 8.0 | 8.0 | 8.0 | |

TABLE A21-1. (CONTINUED) - pH (STANDARD UNITS)

| | <u>Test Concentrations (Percent Groundwater by Volume)</u> | | | | | |
|--------------|--|------|------|------|------|------|
| | 0 | 10 | 18 | 32 | 56 | 100 |
| <u>Day 0</u> | | | | | | |
| 0 H | 7.49 | 7.33 | 6.96 | 6.70 | 6.27 | 3.97 |
| <u>Day 1</u> | | | | | | |
| 0 H | 6.93 | 6.96 | 6.92 | 6.62 | 6.25 | 4.06 |
| 24 H | 6.80 | 7.06 | 7.14 | 7.23 | 7.28 | 5.58 |
| <u>Day 2</u> | | | | | | |
| 0 H | 6.36 | 6.32 | 6.23 | 6.14 | 5.70 | 3.95 |
| 24 H | 7.39 | 7.43 | 7.51 | 7.58 | 7.59 | 5.49 |
| <u>Day 3</u> | | | | | | |
| 0 H | 6.35 | 6.31 | 6.31 | 6.19 | 5.85 | 3.96 |
| 24 H | 6.35 | 6.63 | 6.72 | 6.79 | 6.84 | |
| <u>Day 4</u> | | | | | | |
| 0 H | 6.29 | 6.24 | 6.25 | 6.20 | 6.13 | |
| 24 H | 6.23 | 6.15 | 6.29 | 6.39 | 6.44 | |
| <u>Day 5</u> | | | | | | |
| 0 H | 6.54 | 6.55 | 6.58 | 6.57 | 5.94 | |
| 24 H | 7.13 | 7.14 | 7.24 | 7.30 | 7.34 | |
| <u>Day 6</u> | | | | | | |
| 0 H | 6.43 | 6.62 | 6.67 | 6.67 | 6.19 | |
| 24 H | 6.97 | 7.12 | 7.24 | 7.35 | 7.41 | |
| <u>Day 7</u> | | | | | | |
| 24 H | 6.90 | 7.06 | 7.11 | 7.24 | 7.33 | |

TABLE A21-1. (CONTINUED) - CONDUCTIVITY (μ MHOS/CM)

| | <u>Test Concentrations (Percent Groundwater by Volume)</u> | |
|--------------|--|------------|
| | <u>0</u> | <u>100</u> |
| <u>Day 0</u> | | |
| 0 H | 180 | 1350 |
| <u>Day 1</u> | | |
| 0 H | 160 | 1600 |
| <u>Day 2</u> | | |
| 0 H | 80 | |
| <u>Day 3</u> | | |
| 0 H | 90 | |
| <u>Day 4</u> | | |
| 0 H | 90 | |
| <u>Day 5</u> | | |
| 0 H | 140 | |
| <u>Day 6</u> | | |
| 0 H | 140 | |
| <u>Day 7</u> | | |
| 24 H | 130 | |

TABLE A21-1. (CONTINUED) - ALKALINITY (MG/L AS CaCO₃)

| | <u>Test Concentrations (Percent Groundwater by Volume)</u> | |
|--------------|--|-----|
| | 0 | 100 |
| <u>Day 0</u> | | |
| 0 H | 50 | a |
| <u>Day 1</u> | | |
| 0 H | 65 | a |
| <u>Day 2</u> | | |
| 0 H | 25 | |
| <u>Day 3</u> | | |
| 0 H | 25 | |
| <u>Day 4</u> | | |
| 0 H | 25 | |
| <u>Day 5</u> | | |
| 0 H | 55 | |
| <u>Day 6</u> | | |
| 0 H | 60 | |
| <u>Day 7</u> | | |
| 24 H | 50 | |

a Could not obtain measurement.

TABLE A21-1. (CONTINUED) - HARDNESS (MG/L AS CaCO_3)

| <u>Test Concentrations (Percent Groundwater by Volume)</u> | | |
|--|----------|--------------|
| | <u>0</u> | <u>100</u> |
| <u>Day 0</u> | | |
| 0 H | 50 | ^a |
| <u>Day 1</u> | | |
| 0 H | 60 | ^a |
| <u>Day 2</u> | | |
| 0 H | 36 | |
| <u>Day 3</u> | | |
| 0 H | 24 | |
| <u>Day 4</u> | | |
| 0 H | 24 | |
| <u>Day 5</u> | | |
| 0 H | 50 | |
| <u>Day 6</u> | | |
| 0 H | 54 | |
| <u>Day 7</u> | | |
| 24 H | 50 | |

^a Could not obtain measurement.

TABLE A21-2. CLADOCERAN BEACH POINT RAW GROUNDWATER TOXICITY
TEST DATA (TEST NO. 1) - SURVIVAL AFTER 48 HOURS OF
EXPOSURE

| Concentration (% Groundwater by Volume) | Number Tested | No. Alive at 48 Hours | Percent Alive |
|---|------------------|--------------------------|------------------|
| UMD/WREC Control | 10 | 10 | 100 |
| 10 | 10 | 10 | 100 |
| 18 | 10 | 10 | 100 |
| 32 | 10 | 10 | 100 |
| 56 | 10 | 10 | 100 |
| 100 | 10 | 0 | 0 |

TABLE A21-3. CLADOCERAN BEACH POINT RAW GROUNDWATER TOXICITY DATA (TEST NO. 2) - SURVIVAL OF ADULTS, NUMBER OF YOUNG PRODUCED PER BROOD, TOTAL NUMBER OF YOUNG, AND MEAN NUMBER OF YOUNG PER BROOD AFTER 7 DAYS OF EXPOSURE

| Conc (% by Vol) | Rep | Brood No. 1 | Brood No. 2 | Brood No. 3 | Total Young | Mean Young Per Brood |
|----------------------------|-----|----------------|----------------|----------------|----------------|-------------------------|
| UMD/WREC Control | 1 | 4 | 10 | 22 | 36 | 12.0 |
| | 2 | 4 | 8 | 15 | 27 | 9.0 |
| | 3 | 4 | 8 | 15 | 27 | 9.0 |
| | 4 | 3 | 7 | 10 | 20 | 6.7 |
| | 5 | 6 | 7 | 14 | 27 | 9.0 |
| | 6 | 4 | 6 | 12 | 22 | 7.3 |
| | 7 | 4 | 7 | 16 | 27 | 9.0 |
| | 8 | 5 | 8 | 10 | 23 | 7.7 |
| | 9 | 6 | 8 | 12 | 26 | 8.7 |
| | 10 | 11 | 12 | 13 | 36 | 12.0 |
| APG-EA Diluent Water | 1 | 4 | 9 | 23 | 36 | 12.0 |
| | 2 | 3 | 10 | 19 | 32 | 10.7 |
| | 3 | 5 | 9 | 17 | 31 | 10.3 |
| | 4 | 5 | 7 | 19 | 31 | 10.3 |
| | 5 | 4 | 13 | 12 | 29 | 9.9 |
| | 6 | 6 | 9 | 13 | 28 | 9.3 |
| | 7 | 5 | 12 | 15 | 32 | 10.7 |
| | 8 | 5 | 8 | 8 | 21 | 7.0 |
| | 9 | 4 | 8 | 16 | 28 | 9.3 |
| | 10 | 4 | 9 | 16 | 29 | 9.7 |
| 10 | 1 | 4 | 5 | 12 | 20 | 6.7 |
| | 2 | 5 | 8 | 13 | 26 | 8.7 |
| | 3 | 5 | 6 | 14 | 25 | 8.3 |
| | 4 | 5 | 6 | 15 | 26 | 8.7 |
| | 5 | 4 | 9 | 16 | 29 | 9.7 |
| | 6 | 4 | 0 | 13 | 17 | 8.5 |
| | 7 | 4 | 3 | 14 | 21 | 7.0 |
| | 8 | 5 | 7 | 12 | 24 | 8.0 |
| | 9 | 4 | 8 | 18 | 30 | 10.0 |
| | 10 | 5 | 9 | 13 | 27 | 9.0 |

TABLE A21-3. (CONTINUED)

| Conc (% by Vol) | Rep | Brood No. 1 | Brood No. 2 | Brood No. 3 | Total Young | Mean Young Per Brood |
|-----------------------|-----|----------------|----------------|----------------|----------------|-------------------------|
| 18 | 1 | 2 | 7 | 13 | 22 | 7.3 |
| | 2 | 4 | 7 | 12 | 23 | 7.7 |
| | 3 | 4 | 6 | 10 | 20 | 6.7 |
| | 4 | 4 | 7 | 9 | 20 | 6.7 |
| | 5 | 3 | 6 | 9 | 18 | 6.0 |
| | 6 | 5 | 10 | 12 | 27 | 9.0 |
| | 7 | 3 | 6 | 15 | 24 | 8.0 |
| | 8 | 4 | 9 | 10 | 23 | 7.7 |
| | 9 | 4 | 1 | 12 | 17 | 5.7 |
| | 10 | 4 | 9 | 13 | 26 | 8.7 |
| 32 | 1 | 2 | 6 | 13 | 21 | 7.0 |
| | 2 | 4 | 8 | 9 | 21 | 7.0 |
| | 3 | 4 | 6 | 8 | 18 | 6.0 |
| | 4 | 3 | 4 | 9 | 16 | 5.3 |
| | 5 | 4 | 7 | 8 | 19 | 6.3 |
| | 6 | 3 | 5 | 6 | 14 | 4.7 |
| | 7 | 2 | 5 | 8 | 15 | 5.0 |
| | 8 | 4 | 7 | 8 | 19 | 6.3 |
| | 9 | 3 | 5 | 11 | 19 | 6.3 |
| | 10 | 3 | 7 | 5 | 15 | 5.0 |
| 56 | 1 | 1 | 1 | 2 | 4 | 1.3 |
| | 2 | 2 | 1 | 0 | 3 | 1.5 |
| | 3 | 1 | 1 | 0 | 2 | 1.0 |
| | 4 | 1 | 0 | 0 | 1 | 1.0 |
| | 5 | 2 | 1 | 4 | 7 | 2.3 |
| | 6 | 2 | 0 | 0 | 2 | 2.0 |
| | 7 | 2 | 0 | 3 | 5 | 2.5 |
| | 8 | 3 | 1 | 0 | 4 | 2.0 |
| | 9 | 3 | 0 | 0 | 3 | 3.0 |
| | 10 | 2 | 2 | 1 | 5 | 1.7 |

TABLE A21-3. (CONTINUED)

| Conc (% by Vol) | Rep | Brood No. 1 | Brood No. 2 | Brood No. 3 | Total Young | Mean Young Per Brood |
|-----------------------|-----|----------------|----------------|----------------|----------------|-------------------------|
| 100 | 1 | DEAD | | | | |
| | 2 | DEAD | | | | |
| | 3 | DEAD | | | | |
| | 4 | DEAD | | | | |
| | 5 | DEAD | | | | |
| | 6 | DEAD | | | | |
| | 7 | DEAD | | | | |
| | 8 | DEAD | | | | |
| | 9 | DEAD | | | | |
| | 10 | DEAD | | | | |

TABLE A21-4. CLADOCERAN BEACH POINT RAW GROUNDWATER TOXICITY
TEST STATISTICAL ANALYSIS (TEST NO. 2) - NEONATE
PRODUCTION AFTER 7 DAYS OF EXPOSURE^a

Data Transformation:

None

Chi-square Test for Normality:

| | |
|----------------------------|---|
| Calculated test statistic: | 0.70 |
| Alpha value: | 0.01 |
| Critical value: | 13.28 |
| Conclusion | Fail to reject the null hypothesis that the data are normally distributed |

Bartlett's Test for Homogeneity of Variances:

| | |
|----------------------------|--|
| Calculated test statistic: | 11.25 |
| Alpha value: | 0.01 |
| Critical value: | 15.09 |
| Conclusion: | Fail to reject the null hypothesis that the variances are homogenous |

ANOVA:

| | |
|----------------------------|--|
| Calculated test statistic: | 65.48 |
| Alpha value: | 0.05 |
| Critical value: | 2.45 |
| Conclusion: | Reject the null hypothesis that all groups are equal |

Dunnett's T-test:

| | |
|-----------------------------|--|
| Calculated test statistics: | See Table A21-5 |
| Alpha value: | 0.05 |
| Critical value: | 2.31 |
| Conclusion: | Reject the null hypothesis that all groups are equal |

^a The 100% raw Beach Point groundwater treatment was not included in the statistical analyses because all concentrations above the NOEC for survival are excluded from the hypothesis test for reproductive effects.

TABLE A21-5. CLADOCERAN BEACH POINT RAW GROUNDWATER TOXICITY
TEST STATISTICAL ANALYSIS (TEST NO. 2) - RESULTS OF
DUNNETT'S TEST ON MEAN NEONATE PRODUCTION AFTER 7
DAYS OF EXPOSURE

| Conc (% by Vol) | No. of Reps | Mean No. Neonates Produced | T Statistic | Significance |
|----------------------------|-------------------|----------------------------------|-------------|--------------|
| UMD/WREC Control | 10 | 27.1 | | |
| APG-EA Diluent Water | 10 | 29.5 | -1.47 | |
| 10 | 10 | 24.6 | 1.53 | |
| 18 | 10 | 22.0 | 3.12 | * |
| 32 | 10 | 17.7 | 5.76 | * |
| 56 | 10 | 3.6 | 14.39 | * |

* Significantly different at alpha = 0.05 (Dunnett's critical
value = 2.31).

APPENDIX 22

CLADOCERAN ACUTE AND 7-DAY SURVIVAL AND REPRODUCTION TEST CONDUCTED ON BEACH POINT BUFFERED (pH \approx 7) GROUNDWATER (WELL CC-33B) (TEST NO. 2)

| | |
|---------------------------|--|
| Test Method: | EPA/600/4-89/001 (Weber et al., 1989) |
| Type of Test: | Static renewal (every 24 h) |
| Date: | June 1-8, 1993 |
| Investigator: | S. D. Turley |
| Laboratory: | UMD/WREC |
| Groundwater: | |
| Source: | APG-EA Beach Point Well CC-33B |
| Chemical Characteristics: | See Appendix 48 |
| Dilution Water: | |
| Source: | 20% Perrier: 80% RO water |
| Chemical Characteristics: | See Table A22-1 |
| Test Organism: | |
| Scientific Name: | <u>Ceriodaphnia dubia</u> |
| Age at Start of Test: | <4 h |
| Source: | UMD/WREC culture |
| Experimental Chambers: | |
| Material: | 50 mL glass beaker |
| Test Solution Volume: | 25 mL |
| No. Organisms/Replicate: | 1 |
| No. Organisms/Treatment: | 10 |
| Loading: | 1 organism/beaker |
| Lighting: | Fluorescent; 60-85 foot candles |
| Aeration: | Prior to each renewal |
| pH Buffer: | 10 N NaOH |

Endpoints:

Mortality of adults; number of neonates produced in 3 broods

Water Quality:

Table A22-1

Results:

Mortality:

48-h Exposure:

The data for the 48-h LC50 were obtained from the 7-d study. The buffered groundwater did not affect survival. The data are summarized in Table A22-2.

7-d Exposure:

Significant ($\alpha = 0.05$) mortality occurred to the daphnids exposed to 100% buffered groundwater by volume; statistically significant mortality did not occur at 56% buffered groundwater by volume (see Tables A22-3, A22-4, and A22-5). The 7-d LC50 for adult mortality, which was determined by the moving average angle method, is as follows:

7-d LC50 = 71.8% buffered groundwater by volume (95% confidence limits = 53.77-155.81)

Neonate Production:

Significant ($\alpha = 0.05$) reductions in neonate production relative to the controls occurred at all concentrations down to 18% buffered groundwater by volume, which was the lowest concentration studied (see Tables A22-3, A22-6, and A22-7). The NOEC and LOEC for exposure of the organism to buffered Beach Point groundwater could not be determined because a significant reduction in neonate production occurred at the lowest concentration studied.

TABLE A22-1. SUMMARY OF THE BEACH POINT BUFFERED GROUNDWATER
BIOASSAY WATER QUALITY DATA FOR THE CLADOCERAN 7-
DAY TEST (TEST NO. 2) - DISSOLVED OXYGEN (MG/L)

| | | <u>Test Concentrations (Percent Groundwater by Volume)</u> | | | | |
|--------------|--|--|-----|-----|-----|-----|
| | | 0 | 18 | 32 | 56 | 100 |
| <u>Day 0</u> | | | | | | |
| 0 H | | 8.3 | 7.7 | 7.5 | 7.5 | 6.7 |
| <u>Day 1</u> | | | | | | |
| 0 H | | 8.3 | 8.2 | 8.0 | 7.6 | 6.5 |
| 24 H | | 8.1 | 8.0 | 8.1 | 8.1 | 8.2 |
| <u>Day 2</u> | | | | | | |
| 0 H | | 8.0 | 8.4 | 8.4 | 8.2 | 7.1 |
| 24 H | | 8.1 | 7.9 | 8.0 | 8.0 | 8.1 |
| <u>Day 3</u> | | | | | | |
| 0 H | | 8.3 | 8.1 | 7.8 | 7.5 | 6.2 |
| 24 H | | 7.8 | 8.4 | 8.4 | 8.4 | 8.4 |
| <u>Day 4</u> | | | | | | |
| 0 H | | 8.3 | 7.8 | 7.6 | 7.5 | 6.5 |
| 24 H | | 8.3 | 8.2 | 8.4 | 8.3 | 8.3 |
| <u>Day 5</u> | | | | | | |
| 0 H | | 8.2 | 8.2 | 8.2 | 8.2 | 8.0 |
| 24 H | | 8.3 | 8.3 | 8.2 | 8.2 | 8.1 |
| <u>Day 6</u> | | | | | | |
| 0 H | | 8.5 | 8.3 | 8.1 | 8.4 | 8.0 |
| 24 H | | 8.2 | 8.2 | 8.1 | 8.2 | 8.1 |
| <u>Day 7</u> | | | | | | |
| 24 H | | 8.1 | 8.1 | 8.0 | 8.0 | 7.9 |

TABLE A22-1. (CONTINUED) - pH (STANDARD UNITS)

| | <u>Test Concentrations (Percent Groundwater by Volume)</u> | | | | |
|--------------|--|------|------|------|------|
| | 0 | 18 | 32 | 56 | 100 |
| <u>Day 0</u> | | | | | |
| 0 H | 7.49 | 6.88 | 6.85 | 6.87 | 6.95 |
| <u>Day 1</u> | | | | | |
| 0 H | 6.93 | 6.60 | 6.76 | 6.89 | 6.96 |
| 24 H | 6.80 | 6.63 | 6.84 | 7.06 | 7.24 |
| <u>Day 2</u> | | | | | |
| 0 H | 6.36 | 6.50 | 6.60 | 6.88 | 7.25 |
| 24 H | 7.39 | 7.60 | 7.72 | 7.87 | 8.00 |
| <u>Day 3</u> | | | | | |
| 0 H | 6.35 | 6.82 | 6.94 | 7.12 | 7.60 |
| 24 H | 6.35 | 7.04 | 7.13 | 7.29 | 7.38 |
| <u>Day 4</u> | | | | | |
| 0 H | 6.29 | 6.52 | 6.91 | 6.92 | 7.32 |
| 24 H | 6.23 | 6.57 | 6.92 | 7.16 | 7.66 |
| <u>Day 5</u> | | | | | |
| 0 H | 6.54 | 7.16 | 7.20 | 7.29 | 7.18 |
| 24 H | 7.13 | 7.36 | 7.43 | 7.54 | 7.69 |
| <u>Day 6</u> | | | | | |
| 0 H | 6.43 | 6.79 | 6.91 | 7.13 | 7.57 |
| 24 H | 6.97 | 7.56 | 7.55 | 7.71 | 7.93 |
| <u>Day 7</u> | | | | | |
| 24 H | 6.90 | 7.40 | 7.46 | 7.60 | 7.80 |

TABLE A22-1. (CONTINUED) - CONDUCTIVITY (μ MHOS/CM)

| | <u>Test Concentrations (Percent Groundwater by Volume)</u> | |
|--------------|--|------|
| | 0 | 100 |
| <u>Day 0</u> | | |
| 0 H | 180 | 1180 |
| <u>Day 1</u> | | |
| 0 H | 160 | 1400 |
| <u>Day 2</u> | | |
| 0 H | 80 | 1600 |
| <u>Day 3</u> | | |
| 0 H | 90 | 1500 |
| <u>Day 4</u> | | |
| 0 H | 90 | 1700 |
| <u>Day 5</u> | | |
| 0 H | 140 | 1350 |
| <u>Day 6</u> | | |
| 0 H | 140 | 1700 |
| <u>Day 7</u> | | |
| 24 H | 160 | 1700 |

TABLE A22-1. (CONTINUED) - ALKALINITY (MG/L AS CaCO₃)

| | <u>Test Concentrations (Percent Groundwater by Volume)</u> | |
|--------------|--|-----|
| | 0 | 100 |
| <u>Day 0</u> | | |
| 0 H | 50 | 100 |
| <u>Day 1</u> | | |
| 0 H | 65 | 100 |
| <u>Day 2</u> | | |
| 0 H | 25 | 120 |
| <u>Day 3</u> | | |
| 0 H | 25 | 110 |
| <u>Day 4</u> | | |
| 0 H | 25 | 125 |
| <u>Day 5</u> | | |
| 0 H | 55 | 125 |
| <u>Day 6</u> | | |
| 0 H | 60 | 125 |
| <u>Day 7</u> | | |
| 24 H | 50 | 105 |

TABLE A22-1. (CONTINUED) - HARDNESS (MG/L AS CaCO₃)

| | <u>Test Concentrations (Percent Groundwater by Volume)</u> | |
|--------------|--|-----|
| | 0 | 100 |
| <u>Day 0</u> | | |
| 0 H | 50 | 190 |
| <u>Day 1</u> | | |
| 0 H | 60 | 170 |
| <u>Day 2</u> | | |
| 0 H | 36 | 128 |
| <u>Day 3</u> | | |
| 0 H | 24 | 148 |
| <u>Day 4</u> | | |
| 0 H | 24 | 134 |
| <u>Day 5</u> | | |
| 0 H | 50 | 154 |
| <u>Day 6</u> | | |
| 0 H | 54 | 170 |
| <u>Day 7</u> | | |
| 24 H | 50 | 168 |

TABLE A22-2. CLADOCERAN BEACH POINT BUFFERED GROUNDWATER
TOXICITY TEST DATA (TEST NO. 2) - SURVIVAL AFTER 48
HOURS OF EXPOSURE

| Concentration (% Groundwater by Volume) | Number Tested | No. Alive at 48 Hours | Percent Alive |
|---|------------------|--------------------------|------------------|
| UMD/WREC Control | 10 | 10 | 100 |
| 10 | 10 | 10 | 100 |
| 18 | 10 | 10 | 100 |
| 32 | 10 | 10 | 100 |
| 56 | 10 | 10 | 100 |
| 100 | 10 | 10 | 100 |

TABLE A22-3. CLADOCERAN BEACH POINT BUFFERED GROUNDWATER
TOXICITY DATA (TEST NO.2) - SURVIVAL OF ADULTS,
NUMBER OF YOUNG PRODUCED PER BROOD, TOTAL NUMBER OF
YOUNG, AND MEAN NUMBER OF YOUNG PRODUCED PER BROOD
AFTER 7 DAYS OF EXPOSURE

| Conc (% by Vol) | Rep | Brood No. 1 | Brood No. 2 | Brood No. 3 | Total Young | Mean Young Per Brood |
|-----------------------|-----|----------------|----------------|----------------|----------------|-------------------------|
| UMD/WREC Control | 1 | 4 | 10 | 22 | 36 | 12.0 |
| | 2 | 4 | 8 | 15 | 27 | 9.0 |
| | 3 | 4 | 8 | 15 | 27 | 9.0 |
| | 4 | 3 | 7 | 10 | 20 | 6.7 |
| | 5 | 6 | 7 | 14 | 27 | 9.0 |
| | 6 | 4 | 6 | 12 | 22 | 7.3 |
| | 7 | 4 | 7 | 16 | 27 | 9.0 |
| | 8 | 5 | 8 | 10 | 23 | 7.7 |
| | 9 | 6 | 8 | 12 | 26 | 8.7 |
| | 10 | 11 | 12 | 13 | 36 | 12.0 |
| 18 | 1 | 3 | 0 | 5 | 8 | 4.0 |
| | 2 | 5 | 0 | 0 | 5 | 5.0 |
| | 3 | 5 | 0 | 4 | 9 | 4.5 |
| | 4 | 4 | 0 | 8 | 12 | 6.0 |
| | 5 | 4 | 0 | 0 | 4 | 4.0 |
| | 6 | 5 | 0 | 6 | 11 | 5.5 |
| | 7 | 4 | 0 | 0 | 4 | 4.0 |
| | 8 | 1 | 5 | 0 | 6 | 3.0 |
| | 9 | 2 | 1 | 5 | 8 | 2.7 |
| | 10 | 4 | 0 | 4 | 8 | 4.0 |
| 32 | 1 | 4 | 1 | 0 | 5 | 2.5 |
| | 2 | 4 | 2 | DEAD | 6 | 3.0 |
| | 3 | 4 | 0 | 0 | 4 | 4.0 |
| | 4 | 4 | 0 | 4 | 8 | 4.0 |
| | 5 | 4 | 0 | 3 | 7 | 3.5 |
| | 6 | 4 | 0 | 3 | 7 | 3.5 |
| | 7 | 3 | 0 | 4 | 7 | 3.5 |
| | 8 | 4 | 0 | 3 | 7 | 3.5 |
| | 9 | 5 | 0 | 4 | 9 | 4.5 |
| | 10 | 2 | 1 | 3 | 6 | 2.0 |

TABLE A22-3. (CONTINUED)

| Conc (% by Vol) | Rep | Brood No. 1 | Brood No. 2 | Brood No. 3 | Total Young | Mean Young Per Brood |
|-----------------------|-----|----------------|----------------|----------------|----------------|-------------------------|
| 56 | 1 | 4 | 4 | 0 | 8 | 4.0 |
| | 2 | 2 | 4 | 0 | 6 | 3.0 |
| | 3 | 0 | 1 | 0 | 1 | 1.0 |
| | 4 | 4 | 6 | 0 | 10 | 5.0 |
| | 5 | 4 | 1 | 0 | 5 | 2.5 |
| | 6 | 2 | 5 | 0 | 7 | 3.5 |
| | 7 | 4 | DEAD | | 4 | 4.0 |
| | 8 | 2 | 4 | 0 | 6 | 3.0 |
| | 9 | 4 | 6 | DEAD | 10 | 5.0 |
| | 10 | 3 | 3 | DEAD | 6 | 3.0 |
| 100 | 1 | 0 | 3 | DEAD | 3 | 3.0 |
| | 2 | 0 | 3 | 0 | 3 | 3.0 |
| | 3 | 0 | 3 | DEAD | 3 | 3.0 |
| | 4 | 1 | 0 | DEAD | 1 | 1.0 |
| | 5 | 0 | 4 | 0 | 4 | 4.0 |
| | 6 | 0 | 4 | DEAD | 4 | 4.0 |
| | 7 | 0 | 3 | DEAD | 3 | 3.0 |
| | 8 | 4 | 0 | DEAD | 4 | 4.0 |
| | 9 | 0 | 4 | DEAD | 4 | 4.0 |
| | 10 | 0 | 4 | DEAD | 4 | 4.0 |

TABLE A22-4. CLADOCERAN BEACH POINT BUFFERED GROUNDWATER
TOXICITY TEST STATISTICAL ANALYSIS (TEST NO. 2) -
ADULT SURVIVAL AFTER 7 DAYS OF EXPOSURE

Data Transformation:

None

Fisher's Exact Test:

| | |
|----------------------------|---|
| Calculated test statistic: | See Table A22-5 |
| Alpha value: | 0.05 |
| Critical value: | 6 |
| Conclusion: | Reject the null hypothesis that all groups are equal. |

TABLE A22-5. CLADOCERAN BEACH POINT BUFFERED GROUNDWATER
TOXICITY TEST STATISTICAL ANALYSIS (TEST NO. 2) -
RESULTS OF FISHER'S EXACT TEST ON ADULT SURVIVAL
AFTER 7 DAYS OF EXPOSURE

| Conc (% by Vol) | Number Alive | Number Dead | b Value | Significance |
|-----------------------|-----------------|----------------|------------|--------------|
| UMD/WREC Control | 10 | 0 | | |
| 18 | 10 | 0 | 10 | |
| 32 | 9 | 1 | 9 | |
| 56 | 7 | 3 | 7 | |
| 100 | 2 | 8 | 2 | * |

* Significantly different at $\alpha = 0.05$ (Fisher's critical
value = 6).

TABLE A22-6. CLADOCERAN BEACH POINT BUFFERED GROUNDWATER
TOXICITY TEST STATISTICAL ANALYSIS (TEST NO. 2) -
NEONATE PRODUCTION AFTER 7 DAYS OF EXPOSURE^a

Data Transformation:

None

Chi-Square Test for Normality:

| | |
|----------------------------|---|
| Calculated test statistic: | 3.18 |
| Alpha value: | 0.01 |
| Critical value: | 13.28 |
| Conclusion: | Fail to reject the null hypothesis that the data are normally distributed |

Bartlett's Test for Homogeneity of Variances:

| | |
|----------------------------|--|
| Calculated test statistic: | 13.90 |
| Alpha value: | 0.01 |
| Critical value: | 11.34 |
| Conclusion: | Reject the null hypothesis that the variances are homogenous |

Steel's Many-One Rank Test:

| | |
|-----------------------------|--|
| Calculated test statistics: | See Table A22-7 |
| Alpha value: | 0.05 |
| Critical value: | 77.0 |
| Conclusion: | Reject the null hypothesis that all groups are equal |

^a The 100% buffered Beach Point groundwater treatment was not included in the statistical analyses because all concentrations above the NOEC for survival are excluded from the hypothesis test for reproductive effects.

TABLE A22-7. CLADOCERAN BEACH POINT BUFFERED GROUNDWATER
TOXICITY STATISTICAL ANALYSIS (TEST NO. 2) -
RESULTS OF STEEL'S MANY-ONE RANK TEST ON MEAN
NEONATE PRODUCTION AFTER 7 DAYS OF EXPOSURE

| Conc (% by Vol) | No. of Reps | Mean No. Neonates Produced | Rank Sum | Significance |
|-----------------------|-------------------|----------------------------------|-------------|--------------|
| UMD/WREC Control | 10 | 27.1 | | |
| 18 | 10 | 7.5 | 55.0 | * |
| 32 | 10 | 6.6 | 55.0 | * |
| 56 | 10 | 6.3 | 55.0 | * |

* Significantly different at $\alpha = 0.05$ (Steel's critical
value = 77.0).

APPENDIX 23

CLADOCERAN ACUTE AND 7-DAY SURVIVAL AND REPRODUCTION TEST CONDUCTED ON BEACH POINT RAW (pH \approx 4) GROUNDWATER (WELL CC-33B) (TEST NO. 3)

| | |
|---------------------------|--|
| Test Method: | EPA/600/4-89/001 (Weber et al., 1989) |
| Type of Test: | Static renewal (every 24 h) |
| Date: | August 24-31, 1993 |
| Investigator: | S.D. Turley |
| Laboratory: | UMD/WREC |
| Groundwater: | |
| Source: | APG-EA Beach Point Well CC-33B |
| Chemical Characteristics: | See Appendix 48 |
| Dilution Water: | |
| Source: | 20% Perrier: 80% RO water |
| Chemical Characteristics: | See Table A23-1 |
| Test Organism: | |
| Scientific Name: | <u>Ceriodaphnia dubia</u> |
| Age at Start of Test: | <4 h |
| Source: | UMD/WREC culture |
| Experimental Chambers: | |
| Material: | 50 mL glass beaker |
| Test Solution Volume: | 25 mL |
| No. Organisms/Replicate: | 1 |
| No. Organisms/Treatment: | 10 |
| Loading: | 1 organism/beaker |
| Lighting: | Fluorescent; 60-85 foot candles |
| Aeration: | Prior to each renewal |

Endpoints:

Mortality of adults; number of
neonates produced in 3 broods

Water Quality:

Table A23-1

Results:

Mortality:

48-h Exposure:

The data for the 48-h LC50 were obtained from the 7-d study. All organisms exposed to 100% raw groundwater died during the 48-h exposure (Table A23-2). The organisms in all other treatments lived. The 48-h LC50, which was determined by the moving average angle method, is as follows:

48-h LC50 = 64.7% raw groundwater by volume (95% confidence
limits = 57.17-75.48).

7-d Exposure:

All organisms exposed to 100% raw groundwater died during the 7-d exposure (Table A23-3). Survival was not affected by exposure to concentrations up to 56% raw groundwater by volume. The 7-d LC50, which was determined by the moving average angle method, is as follows:

7-d LC50 = 64.7% raw groundwater by volume (95% confidence
limits = 57.17-75.48).

Neonate Production:

Raw groundwater significantly ($\alpha = 0.05$) reduced neonate production relative to the controls down to 18% raw groundwater by volume (see Tables A23-3, A23-4, and A23-5). The NOEC and LOEC for the daphnids, based on reduced neonate production, are as follows:

NOEC = 10% raw groundwater by volume.

LOEC = 18% raw groundwater by volume.

TABLE A23-1. SUMMARY OF THE BEACH POINT RAW GROUNDWATER BIOASSAY
WATER QUALITY DATA FOR THE CLADOCERAN 7-DAY TEST
(TEST NO. 3) - DISSOLVED OXYGEN (MG/L)

| | <u>Test Concentrations (Percent Groundwater by Volume)</u> | | | | | |
|--------------|--|-----|-----|-----|-----|-----|
| | 0 | 10 | 18 | 32 | 56 | 100 |
| <u>Day 0</u> | | | | | | |
| 0 H | 8.2 | 8.0 | 7.8 | 7.5 | 7.1 | 5.4 |
| <u>Day 1</u> | | | | | | |
| 0 H | 8.1 | 8.1 | 8.0 | 7.7 | 7.3 | |
| 24 H | 8.0 | 7.8 | 7.7 | 7.7 | 7.8 | |
| <u>Day 2</u> | | | | | | |
| 0 H | 8.1 | 8.1 | 8.1 | 8.0 | 7.9 | |
| 24 H | 7.5 | 8.0 | 8.3 | 8.1 | 8.3 | |
| <u>Day 3</u> | | | | | | |
| 0 H | 8.2 | 8.0 | 7.8 | 7.9 | 7.5 | |
| 24 H | 7.1 | 7.1 | 7.0 | 7.8 | 7.7 | |
| <u>Day 4</u> | | | | | | |
| 0 H | 8.1 | 8.0 | 7.9 | 7.8 | 7.8 | |
| 24 H | 7.7 | 7.4 | 7.8 | 7.7 | 7.6 | |
| <u>Day 5</u> | | | | | | |
| 0 H | 8.2 | 7.9 | 7.9 | 7.9 | 7.5 | |
| 24 H | 7.3 | 7.1 | 7.5 | 7.5 | 7.5 | |
| <u>Day 6</u> | | | | | | |
| 0 H | 8.4 | 7.9 | 7.8 | 7.7 | 7.2 | |
| 24 H | 7.6 | 7.8 | 7.9 | 8.0 | 8.0 | |
| <u>Day 7</u> | | | | | | |
| 24 H | 8.2 | 8.0 | 8.2 | 8.1 | 8.2 | |

TABLE A23-1. (CONTINUED) - pH (STANDARD UNITS)

| | <u>Test Concentrations (Percent Groundwater by Volume)</u> | | | | | |
|--------------|--|------|------|------|------|------|
| | 0 | 10 | 18 | 32 | 56 | 100 |
| <u>Day 0</u> | | | | | | |
| 0 H | 7.64 | 7.20 | 6.88 | 6.47 | 5.89 | 3.92 |
| <u>Day 1</u> | | | | | | |
| 0 H | 7.69 | 7.30 | 6.91 | 6.53 | 5.99 | |
| 24 H | 7.97 | 8.05 | 8.17 | 8.18 | 8.08 | |
| <u>Day 2</u> | | | | | | |
| 0 H | 7.83 | 6.91 | 6.78 | 6.53 | 6.65 | |
| 24 H | 7.46 | 7.84 | 7.99 | 7.94 | 7.81 | |
| <u>Day 3</u> | | | | | | |
| 0 H | 7.75 | 7.34 | 7.11 | 6.79 | 6.33 | |
| 24 H | 8.09 | 7.94 | 7.95 | 7.92 | 7.83 | |
| <u>Day 4</u> | | | | | | |
| 0 H | 7.84 | 7.86 | 7.51 | 6.82 | 6.20 | |
| 24 H | 7.50 | 7.62 | 7.67 | 7.64 | 7.52 | |
| <u>Day 5</u> | | | | | | |
| 0 H | 7.57 | 7.50 | 7.12 | 6.70 | 5.87 | |
| 24 H | 7.54 | 7.68 | 7.69 | 7.66 | 7.53 | |
| <u>Day 6</u> | | | | | | |
| 0 H | 7.31 | 7.10 | 6.82 | 6.52 | 5.84 | |
| 24 H | 7.94 | 7.85 | 7.84 | 7.83 | 7.61 | |
| <u>Day 7</u> | | | | | | |
| 24 H | 8.03 | 8.05 | 8.04 | 7.96 | 7.86 | |

TABLE A23-1. (CONTINUED) - CONDUCTIVITY (μ MHOS/CM)

| | <u>Test Concentrations (Percent Groundwater by Volume)</u> | |
|--------------|--|------|
| | 0 | 100 |
| <u>Day 0</u> | | |
| 0 H | 150 | 1750 |
| <u>Day 1</u> | | |
| 0 H | 140 | |
| <u>Day 2</u> | | |
| 0 H | 160 | |
| <u>Day 3</u> | | |
| 0 H | 150 | |
| <u>Day 4</u> | | |
| 0 H | 150 | |
| <u>Day 5</u> | | |
| 0 H | 150 | |
| <u>Day 6</u> | | |
| 0 H | 150 | |
| <u>Day 7</u> | | |
| 24 H | 150 | |

TABLE A23-1. (CONTINUED) - ALKALINITY (MG/L AS CaCO₃)

| <u>Test Concentrations (Percent Groundwater by Volume)</u> | | |
|--|----------|------------|
| | <u>0</u> | <u>100</u> |
| <u>Day 0</u> | | |
| 0 H | 40 | a |
| <u>Day 1</u> | | |
| 0 H | 40 | a |
| <u>Day 2</u> | | |
| 0 H | 60 | |
| <u>Day 3</u> | | |
| 0 H | 45 | |
| <u>Day 4</u> | | |
| 0 H | 55 | |
| <u>Day 5</u> | | |
| 0 H | 50 | |
| <u>Day 6</u> | | |
| 0 H | 60 | |
| <u>Day 7</u> | | |
| 24 H | 55 | |

^a Could not obtain measurement.

TABLE A23-1. (CONTINUED) - HARDNESS (MG/L AS CaCO₃)

| | <u>Test Concentrations (Percent Groundwater by Volume)</u> | |
|--------------|--|------------|
| | <u>0</u> | <u>100</u> |
| <u>Day 0</u> | | |
| 0 H | 54 | a |
| <u>Day 1</u> | | |
| 0 H | 50 | a |
| <u>Day 2</u> | | |
| 0 H | 62 | |
| <u>Day 3</u> | | |
| 0 H | 50 | |
| <u>Day 4</u> | | |
| 0 H | 56 | |
| <u>Day 5</u> | | |
| 0 H | 52 | |
| <u>Day 6</u> | | |
| 0 H | 50 | |
| <u>Day 7</u> | | |
| 24 H | 62 | |

^a Could not obtain measurement.

TABLE A23-2. CLADOCERAN BEACH POINT RAW GROUNDWATER TOXICITY
TEST DATA (TEST NO. 3) - SURVIVAL AFTER 48 HOURS OF
EXPOSURE

| Concentration (% Groundwater by Volume) | Number Tested | No. Alive at 48 Hours | Percent Alive |
|---|------------------|--------------------------|------------------|
| UMD/WREC Control | 10 | 10 | 100 |
| 10 | 10 | 10 | 100 |
| 18 | 10 | 10 | 100 |
| 32 | 10 | 10 | 100 |
| 56 | 10 | 10 | 100 |
| 100 | 10 | 0 | 0 |

TABLE A23-3. CLADOCERAN BEACH POINT RAW GROUNDWATER TOXICITY DATA (TEST NO. 3) - SURVIVAL OF ADULTS, NUMBER OF YOUNG PRODUCED PER BROOD, TOTAL NUMBER OF YOUNG, AND MEAN NUMBER OF YOUNG PER BROOD AFTER 7 DAYS OF EXPOSURE

| Conc (% by Vol) | Rep | Brood No. 1 | Brood No. 2 | Brood No. 3 | Total Young | Mean Young Per Brood |
|----------------------------|-----|----------------|----------------|----------------|----------------|-------------------------|
| UMD/WREC Control | 1 | 3 | 10 | 14 | 27 | 9.0 |
| | 2 | 5 | 8 | 12 | 25 | 8.3 |
| | 3 | 5 | 9 | 12 | 26 | 8.7 |
| | 4 | 3 | 9 | 14 | 26 | 8.7 |
| | 5 | 4 | 10 | 15 | 29 | 9.7 |
| | 6 | 4 | 9 | 14 | 27 | 9.0 |
| | 7 | 4 | 8 | 12 | 24 | 8.0 |
| | 8 | 4 | 8 | 13 | 25 | 8.3 |
| | 9 | 4 | 9 | 14 | 27 | 9.0 |
| | 10 | 3 | 10 | 13 | 26 | 8.7 |
| APG-EA Diluent Water | 1 | 3 | 7 | 13 | 23 | 7.7 |
| | 2 | 3 | 8 | 14 | 25 | 8.3 |
| | 3 | 3 | 6 | 14 | 23 | 7.7 |
| | 4 | 4 | 8 | 15 | 27 | 9.0 |
| | 5 | 3 | 7 | 14 | 24 | 8.0 |
| | 6 | 3 | 6 | 15 | 24 | 8.0 |
| | 7 | 4 | 6 | 14 | 24 | 8.0 |
| | 8 | 3 | 7 | 16 | 26 | 8.7 |
| | 9 | 3 | 7 | 16 | 26 | 8.7 |
| | 10 | 3 | 8 | 14 | 25 | 8.3 |
| 10 | 1 | 4 | 9 | 13 | 26 | 8.7 |
| | 2 | 3 | 9 | 13 | 25 | 8.3 |
| | 3 | 3 | 8 | 14 | 25 | 8.3 |
| | 4 | 3 | 9 | 13 | 25 | 8.3 |
| | 5 | 4 | 10 | 9 | 23 | 7.7 |
| | 6 | 3 | 10 | 16 | 29 | 9.7 |
| | 7 | 3 | 9 | 12 | 24 | 8.0 |
| | 8 | 3 | 9 | 14 | 26 | 8.7 |
| | 9 | 3 | 9 | 12 | 24 | 8.0 |
| | 10 | 3 | 8 | 13 | 24 | 8.0 |

TABLE A23-3. (CONTINUED)

| Conc (% by Vol) | Rep | Brood No. 1 | Brood No. 2 | Brood No. 3 | Total Young | Mean Young Per Brood |
|-----------------------|-----|----------------|----------------|----------------|----------------|-------------------------|
| 18 | 1 | 3 | 9 | 11 | 23 | 7.7 |
| | 2 | 3 | 9 | 12 | 24 | 8.0 |
| | 3 | 4 | 9 | 9 | 22 | 7.3 |
| | 4 | 3 | 8 | 11 | 22 | 7.3 |
| | 5 | 4 | 9 | 14 | 27 | 9.0 |
| | 6 | 3 | 7 | 9 | 19 | 6.3 |
| | 7 | 3 | 11 | 9 | 23 | 7.7 |
| | 8 | 3 | 11 | 12 | 26 | 8.7 |
| | 9 | 5 | 8 | 7 | 20 | 6.6 |
| | 10 | 3 | 7 | 9 | 19 | 6.3 |
| 32 | 1 | 3 | 8 | 11 | 22 | 7.3 |
| | 2 | 3 | 5 | 9 | 17 | 5.7 |
| | 3 | 1 | 1 | 2 | 4 | 1.3 |
| | 4 | 2 | 6 | 11 | 19 | 6.3 |
| | 5 | 2 | 7 | 10 | 19 | 6.3 |
| | 6 | 3 | 9 | 9 | 21 | 7.0 |
| | 7 | 2 | 5 | 15 | 22 | 7.3 |
| | 8 | 3 | 8 | 9 | 20 | 6.7 |
| | 9 | 2 | 6 | 13 | 21 | 7.0 |
| | 10 | 4 | 8 | 11 | 23 | 7.7 |
| 56 | 1 | 2 | 6 | 0 | 8 | 4.0 |
| | 2 | 2 | 4 | 11 | 17 | 5.7 |
| | 3 | 1 | 1 | 2 | 4 | 1.3 |
| | 4 | 1 | 6 | 4 | 11 | 3.7 |
| | 5 | 1 | 2 | 3 | 6 | 2.0 |
| | 6 | 5 | 4 | 0 | 9 | 4.5 |
| | 7 | 2 | 3 | 3 | 8 | 2.7 |
| | 8 | 2 | 3 | 3 | 8 | 2.7 |
| | 9 | 5 | 2 | 4 | 11 | 3.7 |
| | 10 | 1 | 0 | 0 | 1 | 1.0 |

TABLE A23-3. (CONTINUED)

| Conc (% by Vol) | Rep | Brood No. 1 | Brood No. 2 | Brood No. 3 | Total Young | Mean Young Per Brood |
|-----------------------|-----|----------------|----------------|----------------|----------------|-------------------------|
| 100 | 1 | DEAD | | | | |
| | 2 | DEAD | | | | |
| | 3 | DEAD | | | | |
| | 4 | DEAD | | | | |
| | 5 | DEAD | | | | |
| | 6 | DEAD | | | | |
| | 7 | DEAD | | | | |
| | 8 | DEAD | | | | |
| | 9 | DEAD | | | | |
| | 10 | DEAD | | | | |

TABLE A23-4. CLADOCERAN BEACH POINT RAW GROUNDWATER TOXICITY
TEST STATISTICAL ANALYSIS (TEST NO. 3) - NEONATE
PRODUCTION AFTER 7 DAYS OF EXPOSURE^a

Data Transformation:

None

Chi-square Test for Normality:

| | |
|----------------------------|--|
| Calculated test statistic: | 1.19 |
| Alpha value: | 0.01 |
| Critical value: | 13.28 |
| Conclusion | Fail to reject the null hypothesis that the data are normally distribute |

Bartlett's Test for Homogeneity of Variances:

| | |
|----------------------------|--|
| Calculated test statistic: | 20.83 |
| Alpha value: | 0.01 |
| Critical value: | 15.09 |
| Conclusion | Reject the null hypothesis that the variances are homogenous |

Steel's Many-One Rank Test:

| | |
|----------------------------|--|
| Calculated test statistic: | See Table A23-5 |
| Alpha value: | 0.05 |
| Critical value: | 75.0 |
| Conclusion: | Reject the null hypothesis that all groups are equal |

^a The 100% raw Beach Point groundwater treatment was not included in the statistical analyses because all organisms died during the test.

TABLE A23-5. CLADOCERAN BEACH POINT RAW GROUNDWATER TOXICITY
TEST STATISTICAL ANALYSIS (TEST NO. 3) - RESULTS OF
STEEL'S MANY-ONE RANK TEST ON MEAN NEONATE
PRODUCTION AFTER 7 DAYS OF EXPOSURE

| Conc (% by Vol) | No. of Reps | Mean No. Neonates Produced | Rank Sum | Significance |
|----------------------------|-------------------|----------------------------------|-------------|--------------|
| UMD/WREC Control | 10 | 26.2 | | |
| APG-EA Diluent Water | 10 | 24.9 | 79.0 | |
| 10 | 10 | 25.1 | 81.0 | |
| 18 | 10 | 22.5 | 67.5 | * |
| 32 | 10 | 20.1 | 57.0 | * |
| 56 | 10 | 8.3 | 55.0 | * |

* Significantly different at $\alpha = 0.05$ (Steel's critical
value = 75.0).

APPENDIX 24

CLADOCERAN ACUTE AND 7-DAY SURVIVAL AND REPRODUCTION TEST CONDUCTED ON BEACH POINT BUFFERED (pH \approx 7) GROUNDWATER (WELL CC-33B) (TEST NO. 3)

| | |
|---------------------------|--|
| Test Method: | EPA/600/4-89/001 (Weber et al., 1989) |
| Type of Test: | Static renewal (every 24 h) |
| Date: | August 24-31, 1993 |
| Investigator: | S. D. Turley |
| Laboratory: | UMD/WREC |
| Groundwater: | |
| Source: | APG-EA Beach Point Well CC-33B |
| Chemical Characteristics: | See Appendix 48 |
| Dilution Water: | |
| Source: | 20% Perrier: 80% RO water |
| Chemical Characteristics: | See Table A24-1 |
| Test Organism: | |
| Scientific Name: | <u>Ceriodaphnia dubia</u> |
| Age at Start of Test: | <4 h |
| Source: | UMD/WREC culture |
| Experimental Chambers: | |
| Material: | 50 mL glass beaker |
| Test Solution Volume: | 25 mL |
| No. Organisms/Replicate: | 1 |
| No. Organisms/Treatment: | 10 |
| Loading: | 1 organism/beaker |
| Lighting: | Fluorescent; 60-85 foot candles |
| Aeration: | Prior to each renewal |
| pH Buffer: | 10 N NaOH |

Endpoints:

Mortality of adults; number of neonates produced in 3 broods

Water Quality:

Table A24-1

Results:

Mortality:

48-h Exposure:

The data for the 48-h LC50 were obtained from the 7-d study. Forty percent of the organisms died in the 100% buffered groundwater concentration (Table A24-2); the organisms in all other treatments lived. Because <50% mortality occurred at the highest concentration studied, a 48-h LC50 could not be calculated.

7-d Exposure:

Significant ($\alpha = 0.05$) mortality occurred to the daphnids exposed to 100% buffered groundwater by volume; survival was not affected by exposure to concentrations up to 56% buffered groundwater by volume (see Tables A24-3, A24-4, and A24-5). The 7-d LC50 for adult mortality could not be determined by the moving average angle method because 50% mortality occurred at the highest test concentration.

Neonate Production:

Buffered groundwater significantly ($\alpha = 0.05$) reduced neonate production relative to the controls down to 18% buffered groundwater by volume (see Tables A24-3, A24-6, and A24-7). The NOEC and LOEC for the daphnids, based on reduced neonate production, are as follows:

NOEC = 10% raw groundwater by volume.

LOEC = 18% raw groundwater by volume.

TABLE A24-1. SUMMARY OF THE BEACH POINT BUFFERED GROUNDWATER
BIOASSAY WATER QUALITY DATA FOR THE CLADOCERAN 7-
DAY TEST (TEST NO. 3) - DISSOLVED OXYGEN (MG/L)

| | | <u>Test Concentrations (Percent Groundwater by Volume)</u> | | | | | |
|--------------|--|--|-----|-----|-----|-----|-----|
| | | 0 | 10 | 18 | 32 | 56 | 100 |
| <u>Day 0</u> | | | | | | | |
| 0 H | | 8.2 | 8.0 | 8.0 | 7.9 | 7.8 | 5.2 |
| <u>Day 1</u> | | | | | | | |
| 0 H | | 8.1 | 8.1 | 8.1 | 8.0 | 8.0 | 5.9 |
| 24 H | | 8.0 | 7.4 | 7.6 | 7.4 | 7.5 | 7.6 |
| <u>Day 2</u> | | | | | | | |
| 0 H | | 8.1 | 8.1 | 8.1 | 8.0 | 7.9 | 7.6 |
| 24 H | | 7.5 | 7.8 | 7.9 | 8.0 | 7.9 | 7.1 |
| <u>Day 3</u> | | | | | | | |
| 0 H | | 8.2 | 8.0 | 8.0 | 7.9 | 7.8 | 7.6 |
| 24 H | | 7.1 | 7.2 | 7.1 | 7.5 | 7.4 | 7.3 |
| <u>Day 4</u> | | | | | | | |
| 0 H | | 7.7 | 7.9 | 7.9 | 7.8 | 7.7 | 7.5 |
| 24 H | | 7.7 | 7.6 | 7.5 | 7.3 | 7.5 | 7.4 |
| <u>Day 5</u> | | | | | | | |
| 0 H | | 8.2 | 7.8 | 7.9 | 7.9 | 7.8 | 7.6 |
| 24 H | | 7.3 | 7.0 | 7.1 | 7.0 | 7.1 | 7.1 |
| <u>Day 6</u> | | | | | | | |
| 0 H | | 8.4 | 7.8 | 7.7 | 7.6 | 7.2 | 5.0 |
| 24 H | | 7.6 | 7.7 | 7.6 | 7.6 | 7.7 | 7.7 |
| <u>Day 7</u> | | | | | | | |
| 24 H | | 8.2 | 8.0 | 8.0 | 8.0 | 7.8 | 8.0 |

TABLE A24-1. (CONTINUED) - pH (STANDARD UNITS)

| | <u>Test Concentrations (Percent Groundwater by Volume)</u> | | | | | |
|--------------|--|------|------|------|------|------|
| | 0 | 10 | 18 | 32 | 56 | 100 |
| <u>Day 0</u> | | | | | | |
| 0 H | 7.64 | 7.53 | 7.63 | 7.68 | 7.64 | 7.68 |
| <u>Day 1</u> | | | | | | |
| 0 H | 7.69 | 7.59 | 7.69 | 7.79 | 7.71 | 7.78 |
| 24 H | 7.97 | 8.01 | 8.01 | 8.06 | 8.07 | 8.22 |
| <u>Day 2</u> | | | | | | |
| 0 H | 7.83 | 7.50 | 7.58 | 7.56 | 7.53 | 7.69 |
| 24 H | 7.46 | 7.71 | 7.86 | 8.01 | 8.07 | 8.01 |
| <u>Day 3</u> | | | | | | |
| 0 H | 7.75 | 7.80 | 7.92 | 7.94 | 7.90 | 7.87 |
| 24 H | 8.09 | 7.98 | 8.02 | 8.08 | 8.17 | 8.21 |
| <u>Day 4</u> | | | | | | |
| 0 H | 7.84 | 7.57 | 7.73 | 7.81 | 7.80 | 7.71 |
| 24 H | 7.50 | 7.72 | 7.80 | 7.93 | 8.05 | 8.14 |
| <u>Day 5</u> | | | | | | |
| 0 H | 7.57 | 7.63 | 7.78 | 7.81 | 7.86 | 7.67 |
| 24 H | 7.54 | 7.68 | 7.82 | 7.91 | 8.02 | 8.14 |
| <u>Day 6</u> | | | | | | |
| 0 H | 7.31 | 7.54 | 7.60 | 7.79 | 7.85 | 7.75 |
| 24 H | 7.94 | 7.84 | 7.90 | 7.97 | 8.07 | 8.21 |
| <u>Day 7</u> | | | | | | |
| 24 H | 8.03 | 7.35 | 7.62 | 7.73 | 7.86 | 8.01 |

TABLE A24-1. (CONTINUED) - CONDUCTIVITY (μ MHOS/CM)

| <u>Test Concentrations (Percent Groundwater by Volume)</u> | | |
|--|----------|------------|
| | <u>0</u> | <u>100</u> |
| <u>Day 0</u> | | |
| 0 H | 150 | 2000 |
| <u>Day 1</u> | | |
| 0 H | 140 | 1900 |
| <u>Day 2</u> | | |
| 0 H | 160 | 1950 |
| <u>Day 3</u> | | |
| 0 H | 150 | 1900 |
| <u>Day 4</u> | | |
| 0 H | 150 | 1900 |
| <u>Day 5</u> | | |
| 0 H | 150 | 1850 |
| <u>Day 6</u> | | |
| 0 H | 150 | 1925 |
| <u>Day 7</u> | | |
| 24 H | 150 | 1825 |

TABLE A24-1. (CONTINUED) - ALKALINITY (MG/L AS CaCO₃)

| | <u>Test Concentrations (Percent Groundwater by Volume)</u> | |
|--------------|--|-----|
| | 0 | 100 |
| <u>Day 0</u> | | |
| 0 H | 40 | 95 |
| <u>Day 1</u> | | |
| 0 H | 40 | 100 |
| <u>Day 2</u> | | |
| 0 H | 60 | 95 |
| <u>Day 3</u> | | |
| 0 H | 45 | 100 |
| <u>Day 4</u> | | |
| 0 H | 55 | 95 |
| <u>Day 5</u> | | |
| 0 H | 50 | 100 |
| <u>Day 6</u> | | |
| 0 H | 60 | 110 |
| <u>Day 7</u> | | |
| 24 H | 55 | 110 |

TABLE A24-1. (CONTINUED) - HARDNESS (MG/L AS CaCO₃)

| <u>Test Concentrations (Percent Groundwater by Volume)</u> | | |
|--|----------|------------|
| | <u>0</u> | <u>100</u> |
| <u>Day 0</u> | | |
| 0 H | 54 | 220 |
| <u>Day 1</u> | | |
| 0 H | 50 | 230 |
| <u>Day 2</u> | | |
| 0 H | 62 | 220 |
| <u>Day 3</u> | | |
| 0 H | 50 | 240 |
| <u>Day 4</u> | | |
| 0 H | 56 | 228 |
| <u>Day 5</u> | | |
| 0 H | 52 | 232 |
| <u>Day 6</u> | | |
| 0 H | 50 | 164 |
| <u>Day 7</u> | | |
| 24 H | 62 | 164 |

TABLE A24-2. CLADOCERAN BEACH POINT RAW GROUNDWATER TOXICITY
TEST DATA (TEST NO. 3) - SURVIVAL AFTER 48 HOURS OF
EXPOSURE

| Concentration (% Groundwater by Volume) | Number Tested | No. Alive at 48 Hours | Percent Alive |
|---|------------------|--------------------------|------------------|
| UMD/WREC Control | 10 | 10 | 100 |
| 10 | 10 | 10 | 100 |
| 18 | 10 | 10 | 100 |
| 32 | 10 | 10 | 100 |
| 56 | 10 | 10 | 100 |
| 100 | 10 | 6 | 60 |

TABLE A24-3. CLADOCERAN BEACH POINT BUFFERED GROUNDWATER
TOXICITY DATA (TEST NO.3) - SURVIVAL OF ADULTS,
NUMBER OF YOUNG PRODUCED PER BROOD, TOTAL NUMBER OF
YOUNG, AND MEAN NUMBER OF YOUNG PRODUCED PER BROOD
AFTER 7 DAYS OF EXPOSURE

| Conc (% by Vol) | Rep | Brood No. 1 | Brood No. 2 | Brood No. 3 | Total Young | Mean Young Per Brood |
|-----------------------|-----|----------------|----------------|----------------|----------------|-------------------------|
| UMD/WREC Control | 1 | 3 | 10 | 14 | 27 | 9.0 |
| | 2 | 5 | 8 | 12 | 25 | 8.3 |
| | 3 | 5 | 9 | 12 | 26 | 8.7 |
| | 4 | 3 | 9 | 14 | 26 | 8.7 |
| | 5 | 4 | 10 | 15 | 29 | 9.7 |
| | 6 | 4 | 9 | 14 | 27 | 9.0 |
| | 7 | 4 | 8 | 12 | 24 | 8.0 |
| | 8 | 4 | 8 | 13 | 25 | 8.3 |
| | 9 | 4 | 9 | 14 | 27 | 9.0 |
| | 10 | 3 | 10 | 13 | 26 | 8.7 |
| 10 | 1 | 6 | 5 | 14 | 25 | 8.3 |
| | 2 | 4 | 6 | 15 | 25 | 8.3 |
| | 3 | 8 | 5 | 16 | 29 | 9.7 |
| | 4 | 5 | 6 | 13 | 24 | 8.0 |
| | 5 | 3 | 7 | 15 | 25 | 8.3 |
| | 6 | 8 | 5 | 16 | 29 | 9.7 |
| | 7 | 3 | 6 | 15 | 24 | 8.0 |
| | 8 | 4 | 7 | 14 | 25 | 8.3 |
| | 9 | 3 | 7 | 16 | 26 | 8.7 |
| | 10 | 6 | 5 | 13 | 24 | 8.0 |
| 18 | 1 | 6 | 4 | 0 | 10 | 5.0 |
| | 2 | 6 | 2 | 0 | 8 | 4.0 |
| | 3 | 5 | 0 | 0 | 5 | 5.0 |
| | 4 | 5 | 0 | 0 | 5 | 5.0 |
| | 5 | 5 | 0 | 0 | 5 | 5.0 |
| | 6 | 4 | 0 | 0 | 4 | 4.0 |
| | 7 | 5 | 0 | 0 | 5 | 5.0 |
| | 8 | 5 | 0 | 0 | 5 | 5.0 |
| | 9 | 6 | 4 | 0 | 10 | 5.0 |
| | 10 | 5 | 0 | 0 | 5 | 5.0 |

TABLE A24-3. (CONTINUED)

| Conc (% by Vol) | Rep | Brood No. 1 | Brood No. 2 | Brood No. 3 | Total Young | Mean Young Per Brood |
|-----------------------|-----|----------------|----------------|----------------|----------------|-------------------------|
| 32 | 1 | 5 | 0 | 0 | 5 | 5.0 |
| | 2 | 4 | 0 | 0 | 4 | 4.0 |
| | 3 | 4 | 0 | 0 | 4 | 4.0 |
| | 4 | 4 | 0 | 0 | 4 | 4.0 |
| | 5 | 3 | 0 | 0 | 3 | 3.0 |
| | 6 | 4 | 0 | 0 | 0 | 4.0 |
| | 7 | 3 | 0 | 0 | 3 | 3.0 |
| | 8 | 4 | 0 | DEAD | 4 | 4.0 |
| | 9 | 5 | 0 | 0 | 5 | 5.0 |
| | 10 | 3 | 0 | 0 | 3 | 3.0 |
| 56 | 1 | 2 | 0 | 0 | 2 | 2.0 |
| | 2 | 6 | 4 | 0 | 10 | 5.0 |
| | 3 | 6 | 0 | 0 | 6 | 6.0 |
| | 4 | 3 | 0 | 0 | 3 | 3.0 |
| | 5 | 3 | 2 | 0 | 5 | 2.5 |
| | 6 | 2 | 0 | 0 | 2 | 2.0 |
| | 7 | 5 | 0 | 0 | 5 | 5.0 |
| | 8 | 3 | 0 | DEAD | 3 | 3.0 |
| | 9 | 2 | 0 | 0 | 2 | 2.0 |
| | 10 | 1 | 0 | 0 | 1 | 1.0 |
| 100 | 1 | 2 | 0 | 0 | 2 | 2.0 |
| | 2 | DEAD | | | | |
| | 3 | DEAD | | | | |
| | 4 | 3 | 0 | 0 | 3 | 3.0 |
| | 5 | DEAD | | | | |
| | 6 | 0 | 0 | 0 | 0 | 0.0 |
| | 7 | 4 | 0 | 0 | 4 | 4.0 |
| | 8 | 0 | 0 | 0 | 0 | 0.0 |
| | 9 | DEAD | | | | |
| | 10 | DEAD | | | | |

TABLE A24-4. CLADOCERAN BEACH POINT BUFFERED GROUNDWATER
TOXICITY TEST STATISTICAL ANALYSIS (TEST NO. 3) -
ADULT SURVIVAL AFTER 7 DAYS OF EXPOSURE

Data Transformation:

None

Fisher's Exact Test:

| | |
|----------------------------|---|
| Calculated test statistic: | See Table A24-5 |
| Alpha value: | 0.05 |
| Critical value: | 6 |
| Conclusion: | Reject the null hypothesis that all groups are equal. |

TABLE A24-5. CLADOCERAN BEACH POINT BUFFERED GROUNDWATER
TOXICITY TEST STATISTICAL ANALYSIS (TEST NO. 3) -
RESULTS OF FISHER'S EXACT TEST ON ADULT SURVIVAL
AFTER 7 DAYS OF EXPOSURE

| Conc (% by Vol) | Number Alive | Number Dead | b Value | Significance |
|-----------------------|-----------------|----------------|------------|--------------|
| UMD/WREC Control | 10 | 0 | | |
| 10 | 10 | 0 | 10 | |
| 18 | 10 | 0 | 10 | |
| 32 | 10 | 1 | 9 | |
| 56 | 10 | 1 | 9 | |
| 100 | 5 | 5 | 5 | * |

* Significantly different at $\alpha = 0.05$ (Fisher's critical value = 6).

TABLE A24-6. CLADOCERAN BEACH POINT BUFFERED GROUNDWATER
TOXICITY TEST STATISTICAL ANALYSIS (TEST NO. 3) -
NEONATE PRODUCTION AFTER 7 DAYS OF EXPOSURE^a

Data Transformation:

None

Chi-Square Test for Normality:

| | |
|----------------------------|---|
| Calculated test statistic: | 11.50 |
| Alpha value: | 0.01 |
| Critical value: | 13.28 |
| Conclusion: | Fail to reject the null hypothesis that the data are normally distributed |

Bartlett's Test for Homogeneity of Variances:

| | |
|----------------------------|--|
| Calculated test statistic: | 13.77 |
| Alpha value: | 0.01 |
| Critical value: | 13.28 |
| Conclusion: | Reject the null hypothesis that the variances are homogenous |

Steel's Many-One Rank Test:

| | |
|-----------------------------|--|
| Calculated test statistics: | See Table A24-7 |
| Alpha value: | 0.05 |
| Critical value: | 76.0 |
| Conclusion: | Reject the null hypothesis that all groups are equal |

^a The 100% buffered Beach Point groundwater treatment was not included in the statistical analyses because all concentrations above the NOEC for survival are excluded from the hypothesis test for reproductive effects.

TABLE A24-7. CLADOCERAN BEACH POINT BUFFERED GROUNDWATER TOXICITY STATISTICAL ANALYSIS (TEST NO. 3) - RESULTS OF STEEL'S MANY-ONE RANK TEST ON MEAN NEONATE PRODUCTION AFTER 7 DAYS OF EXPOSURE

| Conc (% by Vol) | No. of Reps | Mean No. Neonates Produced | Rank Sum | Significance |
|-----------------------|-------------------|----------------------------------|-------------|--------------|
| UMD/WREC Control | 10 | 26.2 | | |
| 10 | 10 | 24.9 | 77.0 | |
| 18 | 10 | 6.2 | 55.0 | * |
| 32 | 9 | 3.9 | 55.0 | * |
| 56 | 9 | 3.9 | 55.0 | * |

* Significantly different at $\alpha = 0.05$ (Steel's critical value = 76.0).

APPENDIX 25

CLADOCERAN ACUTE AND 7-DAY SURVIVAL AND REPRODUCTION TEST CONDUCTED ON BEACH POINT RAW (pH \approx 4) GROUNDWATER (WELL CC-33B) (TEST NO. 4)

| | |
|---------------------------|--|
| Test Method: | EPA/600/4-89/001 (Weber et al., 1989) |
| Type of Test: | Static renewal (every 24 h) |
| Date: | November 30- December 7, 1993 |
| Investigator: | S.D. Turley |
| Laboratory: | UMD/WREC |
| Groundwater: | |
| Source: | APG-EA Beach Point Well CC-33B |
| Chemical Characteristics: | See Appendix 48 |
| Dilution Water: | |
| Source: | 20% Perrier: 80% RO water |
| Chemical Characteristics: | See Table A25-1 |
| Test Organism: | |
| Scientific Name: | <u>Ceriodaphnia dubia</u> |
| Age at Start of Test: | <4 h |
| Source: | UMD/WREC culture |
| Experimental Chambers: | |
| Material: | 50 mL glass beaker |
| Test Solution Volume: | 25 mL |
| No. Organisms/Replicate: | 1 |
| No. Organisms/Treatment: | 10 |
| Loading: | 1 organism/beaker |
| Lighting: | Fluorescent; 60-85 foot candles |
| Aeration: | Prior to each renewal |
| Endpoints: | Mortality of adults; number of |

neonates produced in 3 broods

Water Quality:

Table A25-1

Results:

Mortality:

48-h Exposure:

The data for the 48-h LC50 were obtained from the 7-d study. All organisms exposed to 100% raw groundwater died during the 48-h exposure (Table A25-2). Twenty percent died at the next dilution of 56% groundwater by volume. No organisms died in the remaining treatments. The 48-h LC50, which was determined by the moving average angle method, is as follows:

48-h LC50 = 62.8% raw groundwater by volume (95% confidence limits = 54.99-73.69).

7-d Exposure:

All organisms exposed to 100% raw groundwater died during the 7-d exposure (Table A25-3). Survival was significantly ($\alpha = 0.05$) reduced by exposure to concentrations of 32% and 56% raw groundwater by volume (Tables A25-4 and A25-5). The 7-d LC50, which was determined by the moving average angle method, is as follows:

7-d LC50 = 42.3% raw groundwater by volume (95% confidence limits = 33.51-53.47).

Neonate Production:

Raw groundwater significantly ($\alpha = 0.05$) reduced neonate production relative to the controls down to 18% raw groundwater by volume (see Tables A25-3, A25-6, and A25-7). The NOEC and LOEC for the daphnids, based on reduced neonate production, are as follows:

NOEC = 10% raw groundwater by volume.
LOEC = 18% raw groundwater by volume.

TABLE A25-1. SUMMARY OF THE BEACH POINT RAW GROUNDWATER BIOASSAY
WATER QUALITY DATA FOR THE CLADOCERAN 7-DAY TEST
(TEST NO. 4) - DISSOLVED OXYGEN (MG/L)

| | | <u>Test Concentrations (Percent Groundwater by Volume)</u> | | | | | |
|--------------|--|--|-----|-----|-----|-----|-----|
| | | 0 | 10 | 18 | 32 | 56 | 100 |
| <u>Day 0</u> | | | | | | | |
| 0 H | | 8.4 | 8.6 | 8.6 | 8.2 | 8.6 | 7.7 |
| <u>Day 1</u> | | | | | | | |
| 0 H | | 8.7 | 8.4 | 8.4 | 8.2 | 8.1 | 7.4 |
| 24 H | | 8.8 | 9.0 | 9.0 | 9.1 | 8.9 | 8.8 |
| <u>Day 2</u> | | | | | | | |
| 0 H | | 8.5 | 8.3 | 8.3 | 8.7 | 8.7 | |
| 24 H | | 8.6 | 8.7 | 8.7 | 8.8 | 8.6 | |
| <u>Day 3</u> | | | | | | | |
| 0 H | | 8.3 | 7.9 | 8.3 | 8.2 | 8.5 | |
| 24 H | | 8.5 | 8.3 | 8.8 | 8.8 | 8.7 | |
| <u>Day 4</u> | | | | | | | |
| 0 H | | 8.5 | 8.4 | 8.4 | 8.3 | 8.5 | |
| 24 H | | 8.6 | 8.5 | 8.7 | 8.6 | 8.6 | |
| <u>Day 5</u> | | | | | | | |
| 0 H | | 8.1 | 8.2 | 8.1 | 8.4 | 8.2 | |
| 24 H | | 8.5 | 8.7 | 8.5 | 8.7 | 8.4 | |
| <u>Day 6</u> | | | | | | | |
| 0 H | | 8.6 | 8.5 | 8.5 | 8.4 | 8.4 | |
| 24 H | | 8.1 | 8.4 | 8.3 | 8.6 | 8.5 | |
| <u>Day 7</u> | | | | | | | |
| 24 H | | 8.0 | 8.3 | 8.2 | 8.3 | 8.3 | |

TABLE A25-1 (CONTINUED) - pH (STANDARD UNITS)

| | <u>Test Concentrations (Percent Groundwater by Volume)</u> | | | | | |
|--------------|--|------|------|------|------|------|
| | 0 | 10 | 18 | 32 | 56 | 100 |
| <u>Day 0</u> | | | | | | |
| 0 H | 6.50 | 6.56 | 6.50 | 6.37 | 6.26 | 4.01 |
| <u>Day 1</u> | | | | | | |
| 0 H | 7.65 | 7.61 | 7.42 | 7.08 | 6.70 | 3.97 |
| 24 H | 7.84 | 7.95 | 7.98 | 8.04 | 7.91 | |
| <u>Day 2</u> | | | | | | |
| 0 H | 7.70 | 7.83 | 7.37 | 6.72 | 6.40 | 3.97 |
| 24 H | 7.68 | 8.06 | 8.16 | 8.20 | 8.15 | |
| <u>Day 3</u> | | | | | | |
| 0 H | 7.51 | 7.43 | 7.35 | 7.07 | 6.57 | 3.91 |
| 24 H | 7.78 | 8.05 | 8.19 | 8.24 | 8.14 | |
| <u>Day 4</u> | | | | | | |
| 0 H | 7.66 | 7.73 | 7.62 | 7.14 | 6.76 | 3.99 |
| 24 H | 7.77 | 8.03 | 8.08 | 8.05 | 8.02 | |
| <u>Day 5</u> | | | | | | |
| 0 H | 7.87 | 7.88 | 7.78 | 7.52 | 6.99 | 3.85 |
| 24 H | 7.84 | 7.88 | 7.86 | 7.84 | 7.78 | |
| <u>Day 6</u> | | | | | | |
| 0 H | 7.71 | 7.48 | 7.35 | 7.33 | 6.80 | 3.88 |
| 24 H | 8.02 | 8.32 | 8.41 | 8.47 | 8.40 | |
| <u>Day 7</u> | | | | | | |
| 24 H | 8.09 | 8.01 | 7.92 | 7.84 | 7.76 | |

TABLE A25-1. (CONTINUED) - CONDUCTIVITY (μ MHOS/CM)

| | <u>Test Concentrations (Percent Groundwater by Volume)</u> | |
|--------------|--|------|
| | 0 | 100 |
| <u>Day 0</u> | | |
| 0 H | 235 | 1750 |
| <u>Day 1</u> | | |
| 0 H | 220 | 1700 |
| <u>Day 2</u> | | |
| 0 H | 230 | 1800 |
| <u>Day 3</u> | | |
| 0 H | 220 | 1700 |
| <u>Day 4</u> | | |
| 0 H | 220 | 1750 |
| <u>Day 5</u> | | |
| 0 H | 200 | 1700 |
| <u>Day 6</u> | | |
| 0 H | 220 | 1750 |
| <u>Day 7</u> | | |
| 24 H | 220 | 1700 |

TABLE A25-1. (CONTINUED) - ALKALINITY (MG/L AS CaCO₃)

| | <u>Test Concentrations (Percent Groundwater by Volume)</u> | |
|--------------|--|--------------|
| | <u>0</u> | <u>100</u> |
| <u>Day 0</u> | | |
| 0 H | 60 | ^a |
| <u>Day 1</u> | | |
| 0 H | 55 | ^a |
| <u>Day 2</u> | | |
| 0 H | 50 | |
| <u>Day 3</u> | | |
| 0 H | 50 | |
| <u>Day 4</u> | | |
| 0 H | 55 | |
| <u>Day 5</u> | | |
| 0 H | 50 | |
| <u>Day 6</u> | | |
| 0 H | 50 | |
| <u>Day 7</u> | | |
| 24 H | 55 | |

^a Could not obtain measurement.

TABLE A25-1. (CONTINUED) - HARDNESS (MG/L AS CaCO₃)

| <u>Test Concentrations (Percent Groundwater by Volume)</u> | | |
|--|----------|------------|
| | <u>0</u> | <u>100</u> |
| <hr/> | | |
| <u>Day 0</u> | | |
| 0 H | 80 | a |
| <u>Day 1</u> | | |
| 0 H | 70 | a |
| <u>Day 2</u> | | |
| 0 H | 64 | |
| <u>Day 3</u> | | |
| 0 H | 70 | |
| <u>Day 4</u> | | |
| 0 H | 60 | |
| <u>Day 5</u> | | |
| 0 H | 60 | |
| <u>Day 6</u> | | |
| 0 H | 80 | |
| <u>Day 7</u> | | |
| 24 H | 70 | |

^a Could not obtain measurement.

TABLE A25-2. CLADOCERAN BEACH POINT RAW GROUNDWATER TOXICITY
TEST DATA (TEST NO. 4) - SURVIVAL AFTER 48 HOURS OF
EXPOSURE

| Concentration (% Groundwater by Volume) | Number Tested | No. Alive at 48 Hours | Percent Alive |
|---|------------------|--------------------------|------------------|
| UMD/WREC Control | 10 | 10 | 100 |
| 10 | 10 | 10 | 100 |
| 18 | 10 | 10 | 100 |
| 32 | 10 | 10 | 100 |
| 56 | 10 | 8 | 80 |
| 100 | 10 | 0 | 0 |

TABLE A25-3. CLADOCERAN BEACH POINT RAW GROUNDWATER TOXICITY DATA (TEST NO. 4) - SURVIVAL OF ADULTS, NUMBER OF YOUNG PRODUCED PER BROOD, TOTAL NUMBER OF YOUNG, AND MEAN NUMBER OF YOUNG PER BROOD AFTER 7 DAYS OF EXPOSURE

| Conc (% by Vol) | Rep | Brood No. 1 | Brood No. 2 | Brood No. 3 | Total Young | Mean Young Per Brood |
|----------------------------|-----|----------------|----------------|----------------|----------------|-------------------------|
| UMD/WREC Control | 1 | 6 | 11 | 20 | 37 | 12.3 |
| | 2 | 5 | 10 | 15 | 30 | 10.0 |
| | 3 | 5 | 9 | 13 | 27 | 9.0 |
| | 4 | 4 | 9 | 14 | 27 | 9.0 |
| | 5 | 4 | 10 | 13 | 27 | 9.0 |
| | 6 | 5 | 11 | 12 | 28 | 9.3 |
| | 7 | 4 | 15 | 14 | 33 | 11.0 |
| | 8 | 4 | 10 | 17 | 31 | 10.3 |
| | 9 | 4 | 11 | 19 | 34 | 11.3 |
| | 10 | 5 | 12 | 11 | 28 | 9.3 |
| APG-EA Diluent Water | 1 | 5 | 9 | 15 | 29 | 9.7 |
| | 2 | 3 | 7 | 19 | 29 | 9.7 |
| | 3 | 3 | 8 | 16 | 27 | 9.0 |
| | 4 | 5 | 9 | 14 | 28 | 9.3 |
| | 5 | 4 | 10 | 17 | 31 | 10.3 |
| | 6 | 4 | 11 | 17 | 32 | 10.7 |
| | 7 | 5 | 8 | 15 | 28 | 9.3 |
| | 8 | 6 | 9 | 12 | 27 | 9.0 |
| | 9 | 4 | 10 | 16 | 30 | 10.0 |
| | 10 | 4 | 14 | 14 | 32 | 10.7 |
| 10 | 1 | 4 | 8 | 18 | 30 | 10.0 |
| | 2 | 4 | 9 | 19 | 32 | 10.7 |
| | 3 | 4 | 9 | 17 | 30 | 10.0 |
| | 4 | 4 | 8 | 17 | 29 | 9.7 |
| | 5 | 4 | 8 | 17 | 29 | 9.7 |
| | 6 | 4 | 8 | 18 | 30 | 10.0 |
| | 7 | 5 | 10 | 18 | 33 | 11.0 |
| | 8 | 4 | 8 | 17 | 29 | 9.7 |
| | 9 | 3 | 9 | 19 | 31 | 10.3 |
| | 10 | 4 | 9 | 18 | 31 | 10.3 |

TABLE A25-3. (CONTINUED)

| Conc (% by Vol) | Rep | Brood No. 1 | Brood No. 2 | Brood No. 3 | Total Young | Mean Young Per Brood |
|-----------------------|-----|----------------|----------------|----------------|----------------|-------------------------|
| 18 | 1 | 0 | 10 | 12 | 22 | 11.0 |
| | 2 | 0 | 7 | 9 | 16 | 8.0 |
| | 3 | 4 | 8 | 10 | 22 | 7.3 |
| | 4 | 0 | 8 | 8 | 16 | 8.0 |
| | 5 | 4 | 7 | 8 | 19 | 6.3 |
| | 6 | 4 | 8 | 9 | 21 | 7.0 |
| | 7 | 4 | 6 | 8 | 18 | 6.0 |
| | 8 | 4 | 7 | 7 | 18 | 6.0 |
| | 9 | 5 | 7 | 8 | 20 | 6.7 |
| | 10 | 3 | 7 | 11 | 21 | 7.0 |
| 32 | 1 | 4 | DEAD | 0 | 4 | 4.0 |
| | 2 | 0 | 0 | DEAD | 0 | 0.0 |
| | 3 | 3 | DEAD | 0 | 3 | 3.0 |
| | 4 | 2 | 0 | 0 | 2 | 2.0 |
| | 5 | 3 | 1 | 0 | 4 | 2.0 |
| | 6 | 4 | 2 | 0 | 6 | 3.0 |
| | 7 | 3 | DEAD | 0 | 3 | 3.0 |
| | 8 | 0 | 0 | 7 | 7 | 7.0 |
| | 9 | 3 | DEAD | 0 | 3 | 3.0 |
| | 10 | 3 | 2 | 0 | 5 | 2.5 |
| 56 | 1 | 0 | 0 | 0 | 0 | 0.0 |
| | 2 | 3 | 0 | 0 | 3 | 3.0 |
| | 3 | 1 | DEAD | 0 | 1 | 1.0 |
| | 4 | 0 | 0 | 0 | 0 | 0.0 |
| | 5 | 0 | DEAD | 0 | 0 | 0.0 |
| | 6 | 0 | DEAD | 0 | 0 | 0.0 |
| | 7 | 0 | DEAD | 0 | 0 | 0.0 |
| | 8 | 0 | 0 | 0 | 0 | 0.0 |
| | 9 | 0 | 0 | 0 | 0 | 0.0 |
| | 10 | DEAD | 0 | 0 | 0 | 0.0 |

TABLE A25-3. (CONTINUED)

| Conc (% by Vol) | Rep | Brood No. 1 | Brood No. 2 | Brood No. 3 | Total Young | Mean Young Per Brood |
|-----------------------|-----|----------------|----------------|----------------|----------------|-------------------------|
| 100 | 1 | DEAD | | | | |
| | 2 | DEAD | | | | |
| | 3 | DEAD | | | | |
| | 4 | DEAD | | | | |
| | 5 | DEAD | | | | |
| | 6 | DEAD | | | | |
| | 7 | DEAD | | | | |
| | 8 | DEAD | | | | |
| | 9 | DEAD | | | | |
| | 10 | DEAD | | | | |

TABLE A25-4. CLADOCERAN BEACH POINT RAW GROUNDWATER TOXICITY
TEST STATISTICAL ANALYSIS (TEST NO. 4) - ADULT
SURVIVAL AFTER 7 DAYS OF EXPOSURE

Data Transformation:

None

Fisher's Exact Test:

| | |
|----------------------------|---|
| Calculated test statistic: | See Table A25-5 |
| Alpha value: | 0.05 |
| Critical value: | 6 |
| Conclusion: | Reject the null hypothesis that all groups are equal. |

TABLE A25-5. CLADOCERAN BEACH POINT RAW GROUNDWATER TOXICITY
TEST STATISTICAL ANALYSIS (TEST NO. 4) - RESULTS
OF FISHER'S EXACT TEST ON ADULT SURVIVAL AFTER 7
DAYS OF EXPOSURE

| Conc (% by | Number Alive | Number Dead | b Value | Significance |
|----------------------------|-----------------|----------------|------------|--------------|
| UMD/WREC Control | 10 | 0 | | |
| APG-EA Diluent Water | 10 | 0 | 10 | |
| 10 | 10 | 0 | 10 | |
| 18 | 10 | 0 | 10 | |
| 32 | 5 | 5 | 5 | * |
| 56 | 5 | 5 | 5 | * |

* Significantly different at $\alpha = 0.05$ (Fisher's critical
value = 6).

TABLE A25-6. CLADOCERAN BEACH POINT RAW GROUNDWATER TOXICITY
TEST STATISTICAL ANALYSIS (TEST NO. 4) - NEONATE
PRODUCTION AFTER 7 DAYS OF EXPOSURE^a

Data Transformation:

None

Chi-square Test for Normality:

| | |
|----------------------------|---|
| Calculated test statistic: | 8.36 |
| Alpha value: | 0.01 |
| Critical value: | 13.28 |
| Conclusion | Fail to reject the null hypothesis that the data are normally distributed |

Bartlett's Test for Homogeneity of Variances:

| | |
|----------------------------|--|
| Calculated test statistic: | 7.30 |
| Alpha value: | 0.01 |
| Critical value: | 11.34 |
| Conclusion | Fail to reject the null hypothesis that the variances are homogenous |

ANOVA:

| | |
|----------------------------|--|
| Calculated test statistic: | 48.70 |
| Alpha value: | 0.05 |
| Critical value: | 2.92 |
| Conclusion: | Reject the null hypothesis that all groups are equal |

Dunnett's Test:

| | |
|----------------------------|--|
| Calculated test statistic: | See Table A25-7 |
| Alpha value: | 0.05 |
| Critical value: | 2.15 |
| Conclusion: | Reject the null hypothesis that all groups are equal |

^a The 32%, 56% and 100% concentrations of Beach Point raw groundwater were not included in the statistical analyses because significant mortality occurred at these concentrations during the test.

TABLE A25-7. CLADOCERAN BEACH POINT RAW GROUNDWATER TOXICITY
TEST STATISTICAL ANALYSIS (TEST NO. 4) - RESULTS OF
DUNNETT'S TEST ON MEAN NEONATE PRODUCTION AFTER 7
DAYS OF EXPOSURE

| Conc (% by Vol) | No. of Reps | Mean No. Neonates Produced | T-Stat | Significance |
|----------------------------|-------------------|----------------------------------|--------|--------------|
| UMD/WREC Control | 10 | 30.2 | | |
| APG-EA Diluent Water | 10 | 29.7 | 0.456 | |
| 10 | 10 | 30.4 | -0.182 | |
| 18 | 10 | 19.3 | 9.945 | * |

* Significantly different at alpha = 0.05 (Dunnett's critical
value = 2.15).

APPENDIX 26

CLADOCERAN ACUTE AND 7-DAY SURVIVAL AND REPRODUCTION TEST CONDUCTED ON BEACH POINT BUFFERED (pH \approx 7) GROUNDWATER (WELL CC-33B) (TEST NO. 4)

Test Method: EPA/600/4-89/001
(Weber et al., 1989)

Type of Test: Static renewal (every 24 h)

Date: November 30 - December 7, 1993

Investigator: S. D. Turley

Laboratory: UMD/WREC

Groundwater:

Source: APG-EA Beach Point Well CC-33B

Chemical Characteristics: See Appendix 48

Dilution Water:

Source: 20% Perrier: 80% RO water

Chemical Characteristics: See Table A26-1

Test Organism:

Scientific Name: Ceriodaphnia dubia

Age at Start of Test: <4 h

Source: UMD/WREC culture

Experimental Chambers:

Material: 50 mL glass beaker

Test Solution Volume: 25 mL

No. Organisms/Replicate: 1

No. Organisms/Treatment: 10

Loading: 1 organism/beaker

Lighting: Fluorescent; 60-85 foot
candles

Aeration: Prior to each renewal

pH Buffer: 10 N NaOH

Endpoints:

Mortality of adults; number of neonates produced in 3 broods

Water Quality:

Table A26-1

Results:

Mortality:

48-h Exposure:

The data for the 48-h LC50 were obtained from the 7-d study. Fifty percent of the organisms died in the 100% buffered groundwater concentration (Table A26-2); twenty percent died at both 56 and 32% groundwater by volume. No mortality occurred in the remaining treatments. Because 50% mortality occurred at the highest concentration studied, a 48-h LC50 could not be calculated.

7-d Exposure:

All organisms exposed to 100% buffered groundwater died during the 7-d exposure (Table A26-3). Survival was significantly ($\alpha = 0.05$) reduced by exposure to concentrations of 32% and 56% buffered groundwater by volume (Tables A26-4 and A26-5). The 7-d LC50, which was determined by the moving average angle method, is as follows:

7-d LC50 = 39.8% buffered groundwater by volume (95% confidence limits = 29.74-51.98).

Neonate Production:

Buffered groundwater significantly ($\alpha = 0.05$) reduced neonate production relative to the controls down to 18% buffered groundwater by volume (see Tables A26-3, A26-6, and A26-7). The NOEC and LOEC for the daphnids, based on reduced neonate production, are as follows:

NOEC = 10% raw groundwater by volume.

LOEC = 18% raw groundwater by volume.

TABLE A26-1. SUMMARY OF THE BEACH POINT BUFFERED GROUNDWATER
BIOASSAY WATER QUALITY DATA FOR THE CLADOCERAN 7-
DAY TEST (TEST NO. 4) - DISSOLVED OXYGEN (MG/L)

| | <u>Test Concentrations (Percent Groundwater by Volume)</u> | | | | | |
|--------------|--|-----|-----|-----|-----|-----|
| | 0 | 10 | 18 | 32 | 56 | 100 |
| <u>Day 0</u> | | | | | | |
| 0 H | 8.4 | 8.5 | 8.6 | 8.2 | 8.2 | 7.6 |
| <u>Day 1</u> | | | | | | |
| 0 H | 8.7 | 8.3 | 8.3 | 8.4 | 8.2 | 7.8 |
| 24 H | 8.8 | 8.9 | 9.0 | 8.8 | 8.6 | 8.8 |
| <u>Day 2</u> | | | | | | |
| 0 H | 8.5 | 8.4 | 8.3 | 8.2 | 8.1 | 7.5 |
| 24 H | 8.6 | 8.7 | 8.7 | 8.7 | 8.7 | 8.8 |
| <u>Day 3</u> | | | | | | |
| 0 H | 8.3 | 8.6 | 8.7 | 8.4 | 8.5 | 7.5 |
| 24 H | 8.5 | 8.8 | 8.9 | 8.8 | 8.8 | 8.8 |
| <u>Day 4</u> | | | | | | |
| 0 H | 8.5 | 8.7 | 8.5 | 8.3 | 8.2 | 7.7 |
| 24 H | 8.5 | 8.6 | 8.7 | 8.4 | 8.2 | 8.6 |
| <u>Day 5</u> | | | | | | |
| 0 H | 8.1 | 8.5 | 8.6 | 8.3 | 8.2 | 7.8 |
| 24 H | 8.5 | 8.6 | 8.4 | 8.6 | 8.4 | 8.6 |
| <u>Day 6</u> | | | | | | |
| 0 H | 8.6 | 8.3 | 8.4 | 8.0 | 8.4 | 8.5 |
| 24 H | 8.1 | 8.5 | 8.4 | 8.5 | 8.4 | |
| <u>Day 7</u> | | | | | | |
| 24 H | 8.0 | 8.4 | 8.3 | 8.4 | 8.2 | |

TABLE A26-1. (CONTINUED) - pH (STANDARD UNITS)

| | | <u>Test Concentrations (Percent Groundwater by Volume)</u> | | | | | |
|--------------|--|--|------|------|------|------|------|
| | | 0 | 10 | 18 | 32 | 56 | 100 |
| <u>Day 0</u> | | | | | | | |
| 0 H | | 6.50 | 6.71 | 6.69 | 6.60 | 6.85 | 7.08 |
| <u>Day 1</u> | | | | | | | |
| 0 H | | 7.65 | 7.65 | 7.80 | 7.82 | 7.85 | 7.59 |
| 24 H | | 7.84 | 8.12 | 8.28 | 8.34 | 8.30 | 8.24 |
| <u>Day 2</u> | | | | | | | |
| 0 H | | 7.70 | 7.40 | 7.64 | 7.81 | 7.79 | 7.47 |
| 24 H | | 7.68 | 8.33 | 8.53 | 8.58 | 8.61 | 8.51 |
| <u>Day 3</u> | | | | | | | |
| 0 H | | 7.51 | 7.59 | 7.86 | 8.06 | 8.08 | 7.69 |
| 24 H | | 7.78 | 8.43 | 8.46 | 8.48 | 8.48 | 8.44 |
| <u>Day 4</u> | | | | | | | |
| 0 H | | 7.66 | 7.76 | 7.90 | 7.96 | 8.03 | 7.70 |
| 24 H | | 7.77 | 8.40 | 8.42 | 8.54 | 8.69 | 8.63 |
| <u>Day 5</u> | | | | | | | |
| 0 H | | 7.87 | 7.88 | 7.99 | 8.01 | 8.06 | 7.11 |
| 24 H | | 7.34 | 8.16 | 8.26 | 8.37 | 8.60 | 8.68 |
| <u>Day 6</u> | | | | | | | |
| 0 H | | 7.71 | 7.56 | 7.66 | 7.69 | 7.70 | 7.39 |
| 24 H | | 8.02 | 8.70 | 8.83 | 8.83 | 8.87 | |
| <u>Day 7</u> | | | | | | | |
| 24 H | | 8.09 | 7.89 | 7.87 | 7.93 | 8.03 | |

TABLE A26-1. (CONTINUED) - CONDUCTIVITY (μ MHOS/CM)

| | <u>Test Concentrations (Percent Groundwater by Volume)</u> | |
|--------------|--|------|
| | 0 | 100 |
| <u>Day 0</u> | | |
| 0 H | 235 | 1750 |
| <u>Day 1</u> | | |
| 0 H | 220 | 1650 |
| <u>Day 2</u> | | |
| 0 H | 230 | 1700 |
| <u>Day 3</u> | | |
| 0 H | 220 | 1700 |
| <u>Day 4</u> | | |
| 0 H | 220 | 1750 |
| <u>Day 5</u> | | |
| 0 H | 200 | 1700 |
| <u>Day 6</u> | | |
| 0 H | 220 | 1750 |
| <u>Day 7</u> | | |
| 24 H | 210 | 1800 |

TABLE A26-1. (CONTINUED) - ALKALINITY (MG/L AS CaCO₃)

| | <u>Test Concentrations (Percent Groundwater by Volume)</u> | |
|--------------|--|-----|
| | 0 | 100 |
| <u>Day 0</u> | | |
| 0 H | 60 | 55 |
| <u>Day 1</u> | | |
| 0 H | 55 | 50 |
| <u>Day 2</u> | | |
| 0 H | 50 | 55 |
| <u>Day 3</u> | | |
| 0 H | 50 | 60 |
| <u>Day 4</u> | | |
| 0 H | 55 | 55 |
| <u>Day 5</u> | | |
| 0 H | 50 | 50 |
| <u>Day 6</u> | | |
| 0 H | 50 | 60 |
| <u>Day 7</u> | | |
| 24 H | 55 | 60 |

TABLE A26-1. (CONTINUED) - HARDNESS (MG/L AS CaCO₃)

| | <u>Test Concentrations (Percent Groundwater by Volume)</u> | |
|--------------|--|-----|
| | 0 | 100 |
| <u>Day 0</u> | | |
| 0 H | 80 | 210 |
| <u>Day 1</u> | | |
| 0 H | 70 | 200 |
| <u>Day 2</u> | | |
| 0 H | 64 | 200 |
| <u>Day 3</u> | | |
| 0 H | 70 | 210 |
| <u>Day 4</u> | | |
| 0 H | 60 | 220 |
| <u>Day 5</u> | | |
| 0 H | 60 | 220 |
| <u>Day 6</u> | | |
| 0 H | 80 | 198 |
| <u>Day 7</u> | | |
| 24 H | 68 | 214 |

TABLE A26-2. CLADOCERAN BEACH POINT RAW GROUNDWATER TOXICITY
TEST DATA (TEST NO. 4) - SURVIVAL AFTER 48 HOURS OF
EXPOSURE

| Concentration (% Groundwater by Volume) | Number Tested | No. Alive at 48 Hours | Percent Alive |
|---|------------------|--------------------------|------------------|
| UMD/WREC Control | 10 | 10 | 100 |
| 10 | 10 | 10 | 100 |
| 18 | 10 | 10 | 100 |
| 32 | 10 | 8 | 80 |
| 56 | 10 | 8 | 80 |
| 100 | 10 | 5 | 50 |

TABLE A26-3. CLADOCERAN BEACH POINT BUFFERED GROUNDWATER
TOXICITY DATA (TEST NO.4) - SURVIVAL OF ADULTS,
NUMBER OF YOUNG PRODUCED PER BROOD, TOTAL NUMBER OF
YOUNG, AND MEAN NUMBER OF YOUNG PRODUCED PER BROOD
AFTER 7 DAYS OF EXPOSURE

| Conc (% by Vol) | Rep | Brood No. 1 | Brood No. 2 | Brood No. 3 | Total Young | Mean Young Per Brood |
|-----------------------|-----|----------------|----------------|----------------|----------------|-------------------------|
| UMD/WREC Control | 1 | 6 | 11 | 20 | 37 | 12.3 |
| | 2 | 5 | 10 | 15 | 30 | 10.0 |
| | 3 | 5 | 9 | 13 | 27 | 9.0 |
| | 4 | 4 | 9 | 14 | 27 | 9.0 |
| | 5 | 4 | 10 | 13 | 27 | 9.0 |
| | 6 | 5 | 11 | 12 | 28 | 9.3 |
| | 7 | 4 | 15 | 14 | 33 | 11.0 |
| | 8 | 4 | 10 | 17 | 31 | 10.3 |
| | 9 | 4 | 11 | 19 | 34 | 11.3 |
| | 10 | 5 | 12 | 11 | 28 | 9.3 |
| 10 | 1 | 4 | 8 | 18 | 30 | 10.0 |
| | 2 | 3 | 7 | 19 | 29 | 9.7 |
| | 3 | 3 | 11 | 18 | 32 | 10.7 |
| | 4 | 4 | 9 | 18 | 31 | 10.3 |
| | 5 | 5 | 10 | 19 | 34 | 11.3 |
| | 6 | 4 | 8 | 17 | 29 | 9.7 |
| | 7 | 4 | 8 | 18 | 30 | 10.0 |
| | 8 | 4 | 8 | 20 | 32 | 10.7 |
| | 9 | 4 | 6 | 17 | 27 | 9.0 |
| | 10 | 4 | 8 | 20 | 32 | 10.7 |
| 18 | 1 | 0 | 0 | 9 | 9 | 9.0 |
| | 2 | 4 | 7 | 9 | 20 | 6.7 |
| | 3 | 0 | 7 | 9 | 16 | 8.0 |
| | 4 | 4 | 8 | 10 | 22 | 7.3 |
| | 5 | 3 | 14 | 0 | 17 | 8.5 |
| | 6 | 4 | 12 | 0 | 16 | 8.0 |
| | 7 | 3 | 8 | 9 | 20 | 6.7 |
| | 8 | 0 | 0 | 6 | 6 | 6.0 |
| | 9 | 3 | 8 | 9 | 20 | 6.7 |
| | 10 | 3 | 10 | 12 | 25 | 8.3 |

TABLE A26-3. (CONTINUED)

| Conc (% by Vol) | Rep | Brood No. 1 | Brood No. 2 | Brood No. 3 | Total Young | Mean Young Per Brood |
|-----------------------|-----|----------------|----------------|----------------|----------------|-------------------------|
| 32 | 1 | 3 | 5 | 7 | 15 | 5.0 |
| | 2 | 0 | 0 | 3 | 3 | 3.0 |
| | 3 | 3 | 0 | 6 | 9 | 4.5 |
| | 4 | 3 | 0 | 5 | 8 | 4.0 |
| | 5 | 3 | 3 | DEAD | 6 | 3.0 |
| | 6 | DEAD | 0 | 0 | 0 | 0.0 |
| | 7 | DEAD | 0 | 0 | 0 | 0.0 |
| | 8 | 4 | 0 | DEAD | 4 | 4.0 |
| | 9 | DEAD | 0 | 0 | 0 | 0.0 |
| | 10 | 0 | 5 | 2 | 7 | 3.5 |
| 56 | 1 | DEAD | 0 | 0 | 0 | 0.0 |
| | 2 | DEAD | 0 | 0 | 0 | 0.0 |
| | 3 | 0 | DEAD | 0 | 0 | 0.0 |
| | 4 | DEAD | 0 | 0 | 0 | 0.0 |
| | 5 | 0 | DEAD | 0 | 0 | 0.0 |
| | 6 | DEAD | 0 | 0 | 0 | 0.0 |
| | 7 | 0 | 0 | 0 | 0 | 0.0 |
| | 8 | 0 | 0 | 0 | 0 | 0.0 |
| | 9 | 3 | 4 | 0 | 7 | 3.5 |
| | 10 | 0 | DEAD | 0 | 0 | 0.0 |
| 100 | 1 | DEAD | | | | |
| | 2 | DEAD | | | | |
| | 3 | DEAD | | | | |
| | 4 | 0 | DEAD | | | |
| | 5 | DEAD | | | | |
| | 6 | DEAD | | | | |
| | 7 | DEAD | | | | |
| | 8 | 0 | DEAD | | | |
| | 9 | DEAD | | | | |
| | 10 | DEAD | | | | |

TABLE A26-4. CLADOCERAN BEACH POINT BUFFERED GROUNDWATER
TOXICITY TEST STATISTICAL ANALYSIS (TEST NO. 4) -
ADULT SURVIVAL AFTER 7 DAYS OF EXPOSURE

Data Transformation:

None

Fisher's Exact Test:

| | |
|----------------------------|---|
| Calculated test statistic: | See Table A26-5 |
| Alpha value: | 0.05 |
| Critical value: | 6 |
| Conclusion: | Reject the null hypothesis that all groups are equal. |

TABLE A26-5. CLADOCERAN BEACH POINT BUFFERED GROUNDWATER
TOXICITY TEST STATISTICAL ANALYSIS (TEST NO. 4) -
RESULTS OF FISHER'S EXACT TEST ON ADULT SURVIVAL
AFTER 7 DAYS OF EXPOSURE

| Conc (% by Vol) | Number Alive | Number Dead | b Value | Significance |
|-----------------------|-----------------|----------------|------------|--------------|
| UMD/WREC Control | 10 | 0 | | |
| 10 | 10 | 0 | 10 | |
| 18 | 10 | 0 | 10 | |
| 32 | 5 | 5 | 5 | * |
| 56 | 3 | 7 | 3 | * |

* Significantly different at $\alpha = 0.05$ (Fisher's critical
value = 6).

TABLE A26-6. CLADOCERAN BEACH POINT BUFFERED GROUNDWATER
TOXICITY TEST STATISTICAL ANALYSIS (TEST NO. 4) -
NEONATE PRODUCTION AFTER 7 DAYS OF EXPOSURE^a

Data Transformation:

None

Chi-Square Test for Normality:

| | |
|----------------------------|---|
| Calculated test statistic: | 0.10 |
| Alpha value: | 0.01 |
| Critical value: | 13.28 |
| Conclusion: | Fail to reject the null hypothesis that the data are normally distributed |

Bartlett's Test for Homogeneity of Variances:

| | |
|----------------------------|--|
| Calculated test statistic: | 8.75 |
| Alpha value: | 0.01 |
| Critical value: | 13.28 |
| Conclusion: | Fail to reject the null hypothesis that the variances are homogenous |

ANOVA:

| | |
|----------------------------|--|
| Calculated test statistic: | 35.49 |
| Alpha value: | 0.05 |
| Critical value: | 3.35 |
| Conclusion: | Reject the null hypothesis that all groups are equal |

Dunnett's Test:

| | |
|-----------------------------|--|
| Calculated test statistics: | See Table A26-7 |
| Alpha value: | 0.05 |
| Critical value: | 2.01 |
| Conclusion: | Reject the null hypothesis that all groups are equal |

^a The 32%, 56% and 100% buffered Beach Point groundwater treatments were not included in the statistical analyses because all concentrations above the NOEC for survival are excluded from the hypothesis test for reproductive effects.

TABLE A26-7. CLADOCERAN BEACH POINT BUFFERED GROUNDWATER
TOXICITY STATISTICAL ANALYSIS (TEST NO. 4) -
RESULTS OF DUNNETT'S TEST ON MEAN NEONATE
PRODUCTION AFTER 7 DAYS OF EXPOSURE

| Conc (% by Vol) | No. of Reps | Mean No. Neonates Produced | T-Stat | Significance |
|-----------------------|-------------------|----------------------------------|--------|--------------|
| UMD/WREC Control | 10 | 30.2 | | |
| 10 | 10 | 30.6 | -0.219 | |
| 18 | 10 | 17.1 | 7.184 | * |

* Significantly different at $\alpha = 0.05$ (Dunnett's critical value = 2.01).

APPENDIX 27

FATHEAD MINNOW ACUTE AND 7-DAY SURVIVAL AND GROWTH TEST CONDUCTED ON BEACH POINT RAW (pH \approx 4) GROUNDWATER (WELL CC-33B) (TEST NO. 1)

| | |
|---------------------------|---|
| Test Method: | EPA/600/4-89/001 (Weber et al., 1989) |
| Type of Test: | Static renewal (every 24 h) |
| Date: | April 7-14, 1993 |
| Investigator: | S.D. Turley |
| Laboratory: | UMD/WREC |
| Groundwater: | |
| Source: | APG-EA Beach Point Well CC-33B |
| Chemical Characteristics: | See Appendix 48 |
| Dilution Water: | |
| Source: | 20% Perrier: 80% RO water |
| Chemical Characteristics: | See Table A27-1 |
| Test Organism: | |
| Scientific Name: | <u>Pimephales promelas</u> |
| Dry Weight: | 0.69 mg (mean weight of controls at end of test) |
| Age at Start of Test: | <24 h |
| Source: | UMD/WREC culture |
| Experimental Chambers: | |
| Material: | 600 mL glass beaker |
| Test Solution Volume: | 400 mL |
| No. Organisms/Replicate: | 10 |
| No. Organisms/Treatment: | 40 |
| Loading: | <0.5 g/L |
| Lighting: | Fluorescent; 60-85 foot candles |

| | |
|----------------|-----------------------|
| Aeration: | Prior to each renewal |
| Endpoints: | Mortality; growth |
| Water Quality: | Table A27-1 |

Results:

Mortality:

96-h Exposure:

The data for the 96-h LC50 were obtained from the 7-d study. All organisms exposed to 100% raw groundwater died during the 96-h exposure (Table A27-2). One organism died at the next dilution of 56% groundwater by volume. No organisms died in the remaining treatments. The 96-h LC50, which was determined by the moving average angle method, is as follows:

96-h LC50 = 63.9% raw groundwater by volume (95% confidence limits = 60.34-68.07).

7-d Exposure:

All organisms exposed to 100% raw groundwater died during the 7-d exposure (Table A27-3). Three organisms died at 56% groundwater by volume; one death occurred at 32% groundwater by volume. No organisms died in the remaining treatments. The 7-d LC50, which was determined by the moving average angle method, is as follows:

7-d LC50 = 61.7% raw groundwater by volume (95% confidence limits = 58.06-65.93).

Growth:

A significant ($\alpha = 0.05$) reduction in growth occurred in fathead minnow larvae exposed for 7 d to both 100% and 56% raw groundwater by volume (see Tables A27-3 A27-4, and A27-5). The NOEC and LOEC for the larval fish, based on reduced growth, are as follows:

NOEC = 32% raw groundwater by volume.

LOEC = 56% raw groundwater by volume.

TABLE A27-1. SUMMARY OF THE BEACH POINT RAW GROUNDWATER BIOASSAY
TEST WATER QUALITY DATA FOR THE FATHEAD MINNOW 7-
DAY TEST (TEST NO. 1) - DISSOLVED OXYGEN (MG/L)

| <u>Test Concentrations (Percent Groundwater by Volume)</u> | | | | | | |
|--|-----|-----|-----|-----|-----|-----|
| | 0 | 10 | 18 | 32 | 56 | 100 |
| <u>Day 0</u> | | | | | | |
| 0 H | 8.8 | 8.7 | 8.6 | 8.3 | 8.3 | 8.2 |
| <u>Day 1</u> | | | | | | |
| 0 H | 8.7 | 8.6 | 8.5 | 8.5 | 8.5 | 5.8 |
| 24 H | 8.1 | 8.0 | 7.9 | 8.0 | 8.0 | 8.2 |
| <u>Day 2</u> | | | | | | |
| 0 H | 8.1 | 7.6 | 7.6 | 7.5 | 7.5 | |
| 24 H | 6.5 | 6.6 | 7.0 | 7.1 | 7.2 | |
| <u>Day 3</u> | | | | | | |
| 0 H | 8.5 | 8.4 | 8.3 | 8.3 | 8.3 | |
| 24 H | 6.4 | 6.5 | 6.9 | 7.0 | 7.1 | |
| <u>Day 4</u> | | | | | | |
| 0 H | 8.6 | 8.4 | 8.4 | 8.2 | 8.2 | |
| 24 H | 6.3 | 6.5 | 6.2 | 6.3 | 6.6 | |
| <u>Day 5</u> | | | | | | |
| 0 H | 8.2 | 8.1 | 8.0 | 8.0 | 7.5 | |
| 24 H | 6.4 | 6.4 | 6.2 | 6.1 | 6.3 | |
| <u>Day 6</u> | | | | | | |
| 0 H | 8.0 | 8.1 | 8.0 | 8.0 | 7.8 | |
| 24 H | 6.1 | 6.2 | 6.0 | 6.0 | 5.9 | |
| <u>Day 7</u> | | | | | | |
| 24 H 6.7 | 6.7 | 6.6 | 6.6 | 6.6 | 6.6 | |

TABLE A27-1. (CONTINUED) - pH (STANDARD UNITS)

| | | <u>Test Concentrations (Percent Groundwater by Volume)</u> | | | | | |
|--------------|--|--|------|------|------|------|------|
| | | 0 | 10 | 18 | 32 | 56 | 100 |
| <u>Day 0</u> | | | | | | | |
| 0 H | | 6.71 | 6.69 | 6.43 | 6.20 | 5.93 | 3.95 |
| <u>Day 1</u> | | | | | | | |
| 0 H | | 6.46 | 6.45 | 6.46 | 6.42 | 5.64 | 3.74 |
| 24 H | | 6.63 | 6.73 | 6.80 | 6.84 | 6.84 | 4.80 |
| <u>Day 2</u> | | | | | | | |
| 0 H | | 7.30 | 7.11 | 6.80 | 6.51 | 6.02 | |
| 24 H | | 6.42 | 6.54 | 6.63 | 6.70 | 6.68 | |
| <u>Day 3</u> | | | | | | | |
| 0 H | | 6.93 | 6.91 | 6.74 | 6.44 | 6.07 | |
| 24 H | | 6.51 | 6.57 | 6.69 | 6.78 | 6.71 | |
| <u>Day 4</u> | | | | | | | |
| 0 H | | 7.25 | 7.10 | 7.05 | 6.68 | 6.19 | |
| 24 H | | 6.58 | 6.48 | 6.55 | 6.59 | 6.67 | |
| <u>Day 5</u> | | | | | | | |
| 0 H | | 6.64 | 6.65 | 6.56 | 6.65 | 6.36 | |
| 24 H | | 6.68 | 6.45 | 6.48 | 6.57 | 6.62 | |
| <u>Day 6</u> | | | | | | | |
| 0 H | | 6.70 | 6.70 | 6.57 | 6.69 | 6.41 | |
| 24 H | | 7.20 | 6.86 | 6.94 | 7.04 | 7.10 | |
| <u>Day 7</u> | | | | | | | |
| 24 H | | 6.34 | 6.43 | 6.53 | 6.61 | 6.65 | |

TABLE A27-1. (CONTINUED) - CONDUCTIVITY (μ MHOS/CM)

| | <u>Test Concentrations (Percent Groundwater by Volume)</u> | |
|--------------|--|------|
| | 0 | 100 |
| <u>Day 0</u> | | |
| 0 H | 140 | 1500 |
| <u>Day 1</u> | | |
| 0 H | 155 | 1600 |
| <u>Day 2</u> | | |
| 0 H | 155 | |
| <u>Day 3</u> | | |
| 0 H | 155 | |
| <u>Day 4</u> | | |
| 0 H | 155 | |
| <u>Day 5</u> | | |
| 0 H | 155 | |
| <u>Day 6</u> | | |
| 0 H | 155 | |
| <u>Day 7</u> | | |
| 24 H | 152 | |

TABLE A27-1. (CONTINUED) - ALKALINITY (MG/L AS CaCO₃)

| <u>Test Concentrations (Percent Groundwater by Volume)</u> | | |
|--|----------|--------------|
| | <u>0</u> | <u>100</u> |
| <u>Day 0</u> | | |
| 0 H | 50 | ^a |
| <u>Day 1</u> | | |
| 0 H | 50 | ^a |
| <u>Day 2</u> | | |
| 0 H | 50 | |
| <u>Day 3</u> | | |
| 0 H | 45 | |
| <u>Day 4</u> | | |
| 0 H | 50 | |
| <u>Day 5</u> | | |
| 0 H | 50 | |
| <u>Day 6</u> | | |
| 0 H | 50 | |
| <u>Day 7</u> | | |
| 24 H | 49 | |

^a Could not obtain measurement.

TABLE A27-1. (CONTINUED) - HARDNESS (MG/L AS CaCO₃)

| <u>Test Concentrations (Percent Groundwater by Volume)</u> | | |
|--|----------|------------|
| | <u>0</u> | <u>100</u> |
| <u>Day 0</u> | | |
| 0 H | 48 | a |
| <u>Day 1</u> | | |
| 0 H | 54 | a |
| <u>Day 2</u> | | |
| 0 H | 54 | |
| <u>Day 3</u> | | |
| 0 H | 54 | |
| <u>Day 4</u> | | |
| 0 H | 54 | |
| <u>Day 5</u> | | |
| 0 H | 46 | |
| <u>Day 6</u> | | |
| 0 H | 46 | |
| <u>Day 7</u> | | |
| 24 H | 50 | |

^a Could not obtain measurement.

TABLE A27-2. FATHEAD MINNOW BEACH POINT RAW GROUNDWATER TOXICITY
TEST DATA (TEST NO. 1) - SURVIVAL AFTER 96 HOURS OF
EXPOSURE

| Concentration (% Groundwater by Volume) | Rep | Number Tested | No. Alive at 96 Hours | Percent Alive |
|---|-----|------------------|--------------------------|------------------|
| UMD/WREC Control | A | 10 | 10 | 100 |
| | B | 10 | 10 | 100 |
| | C | 10 | 10 | 100 |
| | D | 10 | 10 | 100 |
| 10 | A | 10 | 10 | 100 |
| | B | 10 | 10 | 100 |
| | C | 10 | 10 | 100 |
| | D | 10 | 10 | 100 |
| 18 | A | 10 | 10 | 100 |
| | B | 10 | 10 | 100 |
| | C | 10 | 10 | 100 |
| | D | 10 | 10 | 100 |
| 32 | A | 10 | 10 | 100 |
| | B | 10 | 10 | 100 |
| | C | 10 | 10 | 100 |
| | D | 10 | 10 | 100 |
| 56 | A | 10 | 9 | 90 |
| | B | 10 | 10 | 100 |
| | C | 10 | 10 | 100 |
| | D | 10 | 10 | 100 |
| 100 | A | 10 | 0 | 0 |
| | B | 10 | 0 | 0 |
| | C | 10 | 0 | 0 |
| | D | 10 | 0 | 0 |

TABLE A27-3. FATHEAD MINNOW BEACH POINT RAW GROUNDWATER TOXICITY
TEST DATA (TEST NO. 1) - LARVAL SURVIVAL AND DRY
WEIGHT AFTER 7 DAYS OF EXPOSURE

| Conc (% by Vol) | Rep | Number Larvae Alive | Percent Survival | Dry Weight (mg) | Mean Dry Weight (mg) |
|----------------------------|-----|---------------------------|---------------------|--------------------|----------------------------|
| UMD/WREC Control | 1 | 10 | 100 | 0.76 | 0.69 |
| | 2 | 10 | 100 | 0.61 | |
| | 3 | 10 | 100 | 0.70 | |
| | 4 | 10 | 100 | 0.67 | |
| APG-EA Diluent Water | 1 | 10 | 100 | 0.71 | 0.67 |
| | 2 | 10 | 100 | 0.61 | |
| | 3 | 10 | 100 | 0.70 | |
| | 4 | 10 | 100 | 0.66 | |
| 10 | 1 | 10 | 100 | 0.71 | 0.68 |
| | 2 | 10 | 100 | 0.60 | |
| | 3 | 10 | 100 | 0.70 | |
| | 4 | 10 | 100 | 0.71 | |
| 18 | 1 | 10 | 100 | 0.65 | 0.67 |
| | 2 | 10 | 100 | 0.71 | |
| | 3 | 10 | 100 | 0.56 | |
| | 4 | 10 | 100 | 0.75 | |
| 32 | 1 | 10 | 100 | 0.67 | 0.62 |
| | 2 | 9 | 90 | 0.66 | |
| | 3 | 10 | 100 | 0.61 | |
| | 4 | 10 | 100 | 0.56 | |
| 56 | 1 | 9 | 90 | 0.41 | 0.40 |
| | 2 | 10 | 100 | 0.41 | |
| | 3 | 9 | 90 | 0.35 | |
| | 4 | 9 | 90 | 0.42 | |
| 100 | 1 | 0 | 0 | | |
| | 2 | 0 | 0 | | |
| | 3 | 0 | 0 | | |
| | 4 | 0 | 0 | | |

TABLE A27-4. FATHEAD MINNOW BEACH POINT RAW GROUNDWATER TOXICITY
TEST STATISTICAL ANALYSIS (TEST NO. 1) - DRY WEIGHT
OF LARVAE AFTER 7 DAYS OF EXPOSURE

Data Transformation:

None

Chi-Square Test for Normality:

| | |
|----------------------------|---|
| Calculated test statistic: | 6.40 |
| Alpha value: | 0.01 |
| Critical value: | 13.28 |
| Conclusion: | Fail to reject the null hypothesis that the data are normally distributed |

Bartlett's Test for Homogeneity of Variances:

| | |
|----------------------------|--|
| Calculated test statistic: | 2.57 |
| Alpha value: | 0.01 |
| Critical value: | 15.09 |
| Conclusion: | Fail to reject the null hypothesis that the variances are homogenous |

ANOVA:

| | |
|----------------------------|--|
| Calculated test statistic: | 16.67 |
| Alpha value: | 0.05 |
| Critical value: | 2.41 |
| Conclusion: | Reject the null hypothesis that all groups are equal |

Dunnett's Test:

| | |
|----------------------------|--|
| Calculated test statistic: | See Table A27-5 |
| Alpha value: | 0.05 |
| Critical value: | 2.41 |
| Conclusion: | Reject the null hypothesis that all groups are equal |

TABLE A27-5. FATHEAD MINNOW BEACH POINT RAW GROUNDWATER TOXICITY TEST STATISTICAL ANALYSIS (TEST NO. 1) - RESULTS OF DUNNETT'S TEST ON DRY WEIGHT OF LARVAE AFTER 7 DAYS OF EXPOSURE

| Conc (% by Vol) | No. of Reps | Mean Dry Weight (mg) | T Statistic | Significance |
|----------------------------|-------------------|----------------------------|-------------|--------------|
| UMD/WREC Control | 4 | 0.69 | | |
| APG-EA Diluent Water | 4 | 0.67 | 0.39 | |
| 10 | 4 | 0.68 | 0.13 | |
| 18 | 4 | 0.67 | 0.45 | |
| 32 | 4 | 0.63 | 1.55 | |
| 56 | 4 | 0.40 | 7.41 | * |

* Significantly different at $\alpha = 0.05$ (Dunnett's critical value = 2.41).

APPENDIX 28

FATHEAD MINNOW ACUTE AND 7-DAY SURVIVAL AND GROWTH
TEST CONDUCTED ON BEACH POINT BUFFERED (pH \approx 7)
GROUNDWATER (WELL CC-33B)
(TEST NO. 1)

| | |
|---------------------------|---|
| Test Method: | EPA/600/4-89/001 (Weber et al., 1989) |
| Type of Test: | Static renewal (every 24 h) |
| Date: | April 7-14, 1993 |
| Investigator: | S.D. Turley |
| Laboratory: | UMD/WREC |
| Groundwater: | |
| Source: | APG-EA Beach Point Well CC-33B |
| Chemical Characteristics: | See Appendix 48 |
| Dilution Water: | |
| Source: | 20% Perrier: 80% RO water |
| Chemical Characteristics: | See Table A28-1 |
| Test Organism: | |
| Scientific Name: | <u>Pimephales promelas</u> |
| Dry Weight: | 0.69 mg (mean weight of controls at end of test) |
| Age at Start of Test: | <24 h |
| Source: | UMD/WREC culture |
| Experimental Chambers: | |
| Material: | 600 mL glass beaker |
| Test Solution Volume: | 400 mL |
| No. Organisms/Replicate: | 10 |
| No. Organisms/Treatment: | 40 |
| Loading: | <0.5 g/L |
| Lighting: | Fluorescent; 60-85 foot candles |
| Aeration: | Prior to renewals |

| | |
|----------------|-------------------|
| pH Buffer: | 10 N NaOH |
| Endpoints: | Mortality; growth |
| Water Quality: | Table A28-1 |

Results:

Mortality:

96-h Exposure:

The data for the 96-h LC50 were obtained from the 7-d study. The buffered groundwater did not affect larval survival. The data are summarized in Table A28-2.

7-d Exposure:

The survival of fathead minnow larvae was not affected after 7 days of exposure to Beach Point buffered groundwater (Table A28-3).

Growth:

A significant ($\alpha = 0.05$) reduction in growth occurred in fathead minnow larvae exposed to 100% buffered groundwater (see Tables A28-3, A28-4, and A28-5). The NOEC and LOEC for the larval fish, based on reduced growth, are as follows:

NOEC = 56% raw groundwater by volume.
LOEC = 100% raw groundwater by volume.

TABLE A28-1. SUMMARY OF THE BEACH POINT BUFFERED GROUNDWATER
BIOASSAY TEST WATER QUALITY DATA FOR THE FATHEAD
MINNOW 7-DAY TEST (TEST NO. 1) - DISSOLVED OXYGEN
(MG/L)

| | | <u>Test Concentrations (Percent Groundwater by Volume)</u> | | | |
|--------------|--|--|-----|-----|-----|
| | | 0 | 32 | 56 | 100 |
| <u>Day 0</u> | | | | | |
| 0 H | | 8.8 | 8.1 | 8.1 | 8.2 |
| <u>Day 1</u> | | | | | |
| 0 H | | 8.7 | 8.2 | 8.2 | 5.6 |
| 24 H | | 8.1 | 8.0 | 8.0 | 8.1 |
| <u>Day 2</u> | | | | | |
| 0 H | | 8.1 | 8.2 | 7.3 | 6.6 |
| 24 H | | 6.5 | 6.8 | 6.8 | 7.0 |
| <u>Day 3</u> | | | | | |
| 0 H | | 8.5 | 8.2 | 8.2 | 7.5 |
| 24 H | | 6.4 | 6.7 | 6.7 | 6.9 |
| <u>Day 4</u> | | | | | |
| 0 H | | 8.6 | 7.9 | 7.9 | 6.2 |
| 24 H | | 6.3 | 6.0 | 6.0 | 6.7 |
| <u>Day 5</u> | | | | | |
| 0 H | | 8.2 | 7.5 | 7.5 | 5.1 |
| 24 H | | 6.4 | 6.1 | 6.1 | 6.5 |
| <u>Day 6</u> | | | | | |
| 0 H | | 8.0 | 8.1 | 8.1 | 6.1 |
| 24 H | | 6.1 | 6.0 | 5.9 | 6.0 |
| <u>Day 7</u> | | | | | |
| 24 H | | 6.6 | 6.6 | 6.6 | 6.5 |

TABLE A28-1. (CONTINUED) - pH (STANDARD UNITS)

| | <u>Test Concentrations (Percent Groundwater by Volume)</u> | | | |
|--------------|--|------|------|------|
| | 0 | 32 | 56 | 100 |
| <u>Day 0</u> | | | | |
| 0 H | 6.71 | 6.47 | 6.61 | 6.98 |
| <u>Day 1</u> | | | | |
| 0 H | 6.46 | 6.37 | 6.51 | 6.69 |
| 24 H | 6.63 | 6.65 | 6.87 | 7.06 |
| <u>Day 2</u> | | | | |
| 0 H | 7.30 | 6.91 | 6.97 | 7.05 |
| 24 H | 6.42 | 6.75 | 7.02 | 7.14 |
| <u>Day 3</u> | | | | |
| 0 H | 6.93 | 6.79 | 6.84 | 6.92 |
| 24 H | 6.51 | 6.86 | 7.10 | 7.17 |
| <u>Day 4</u> | | | | |
| 0 H | 7.25 | 6.91 | 7.00 | 7.06 |
| 24 H | 6.58 | 6.91 | 7.03 | 7.00 |
| <u>Day 5</u> | | | | |
| 0 H | 6.64 | 6.89 | 6.95 | 7.00 |
| 24 H | 6.68 | 6.61 | 6.69 | 6.90 |
| <u>Day 6</u> | | | | |
| 0 H | 6.70 | 6.91 | 6.93 | 6.95 |
| 24 H | 7.20 | 7.14 | 7.18 | 7.20 |
| <u>Day 7</u> | | | | |
| 24 H | 6.34 | 6.86 | 6.95 | 7.08 |

TABLE A28-1. (CONTINUED) - CONDUCTIVITY (μ MHOS/CM)

| <u>Test Concentrations (Percent Groundwater by Volume)</u> | | |
|--|----------|------------|
| | <u>0</u> | <u>100</u> |
| <u>Day 0</u> | | |
| 0 H | 140 | 1700 |
| <u>Day 1</u> | | |
| 0 H | 155 | 1700 |
| <u>Day 2</u> | | |
| 0 H | 155 | 1700 |
| <u>Day 3</u> | | |
| 0 H | 155 | 1700 |
| <u>Day 4</u> | | |
| 0 H | 155 | 1750 |
| <u>Day 5</u> | | |
| 0 H | 155 | 1800 |
| <u>Day 6</u> | | |
| 0 H | 155 | 1800 |
| <u>Day 7</u> | | |
| 24 H | 152 | 1736 |

TABLE A28-1. (CONTINUED) - ALKALINITY (MG/L AS CaCO₃)

| | <u>Test Concentrations (Percent Groundwater by Volume)</u> | |
|--------------|--|-----|
| | 0 | 100 |
| <u>Day 0</u> | | |
| 0 H | 50 | 80 |
| <u>Day 1</u> | | |
| 0 H | 50 | 80 |
| <u>Day 2</u> | | |
| 0 H | 50 | 85 |
| <u>Day 3</u> | | |
| 0 H | 45 | 80 |
| <u>Day 4</u> | | |
| 0 H | 50 | 100 |
| <u>Day 5</u> | | |
| 0 H | 50 | 85 |
| <u>Day 6</u> | | |
| 0 H | 50 | 85 |
| <u>Day 7</u> | | |
| 24 H | 49 | 85 |

TABLE A28-1. (CONTINUED) - HARDNESS (MG/L AS CaCO₃)

| <u>Test Concentrations (Percent Groundwater by Volume)</u> | | |
|--|----------|------------|
| | <u>0</u> | <u>100</u> |
| <u>Day 0</u> | | |
| 0 H | 48 | 280 |
| <u>Day 1</u> | | |
| 0 H | 54 | 170 |
| <u>Day 2</u> | | |
| 0 H | 54 | 136 |
| <u>Day 3</u> | | |
| 0 H | 54 | 148 |
| <u>Day 4</u> | | |
| 0 H | 54 | 138 |
| <u>Day 5</u> | | |
| 0 H | 46 | 160 |
| <u>Day 6</u> | | |
| 0 H | 46 | 160 |
| <u>Day 7</u> | | |
| 24 H | 50 | 170 |

TABLE A28-2. FATHEAD MINNOW BEACH POINT BUFFERED GROUNDWATER TOXICITY TEST DATA (TEST NO. 1) - SURVIVAL AFTER 96 HOURS OF EXPOSURE

| Concentration (% Groundwater by Volume) | Rep | Number Tested | No. Alive at 96 Hours | Percent Alive |
|---|-----|------------------|--------------------------|------------------|
| UMD/WREC Control | A | 10 | 10 | 100 |
| | B | 10 | 10 | 100 |
| | C | 10 | 10 | 100 |
| | D | 10 | 10 | 100 |
| 10 | A | ^a | | |
| | B | ^a | | |
| | C | ^a | | |
| | D | ^a | | |
| 18 | A | ^a | | |
| | B | ^a | | |
| | C | ^a | | |
| | D | ^a | | |
| 32 | A | 10 | 10 | 100 |
| | B | 10 | 10 | 100 |
| | C | 10 | 10 | 100 |
| | D | 10 | 10 | 100 |
| 56 | A | 10 | 9 | 90 |
| | B | 10 | 10 | 100 |
| | C | 10 | 10 | 100 |
| | D | 10 | 10 | 100 |
| 100 | A | 10 | 10 | 100 |
| | B | 10 | 10 | 100 |
| | C | 10 | 10 | 100 |
| | D | 10 | 10 | 100 |

^a Larvae were not tested at this treatment.

TABLE A28-3. FATHEAD MINNOW BEACH POINT BUFFERED GROUNDWATER
TOXICITY TEST DATA (TEST NO. 1) - LARVAL SURVIVAL
AND DRY WEIGHT AFTER 7 DAYS OF EXPOSURE

| Conc (% by Vol) | Rep | Number Larvae Alive | Percent Survival | Dry Weight (mg) | Mean Dry Weight (mg) |
|-----------------------|-----|---------------------------|---------------------|--------------------|----------------------------|
| UMD/WREC Control | 1 | 10 | 100 | 0.76 | 0.69 |
| | 2 | 10 | 100 | 0.61 | |
| | 3 | 10 | 100 | 0.70 | |
| | 4 | 10 | 100 | 0.67 | |
| 32 | 1 | 10 | 100 | 0.68 | 0.65 |
| | 2 | 10 | 100 | 0.62 | |
| | 3 | 10 | 100 | 0.61 | |
| | 4 | 10 | 100 | 0.68 | |
| 56 | 1 | 10 | 100 | 0.76 | 0.67 |
| | 2 | 10 | 100 | 0.70 | |
| | 3 | 10 | 100 | 0.62 | |
| | 4 | 10 | 100 | 0.59 | |
| 100 | 1 | 10 | 100 | 0.50 | 0.54 |
| | 2 | 10 | 100 | 0.55 | |
| | 3 | 10 | 100 | 0.57 | |
| | 4 | 10 | 100 | 0.52 | |

TABLE A28-4. FATHEAD MINNOW BEACH POINT BUFFERED GROUNDWATER
TOXICITY TEST STATISTICAL ANALYSIS (TEST NO. 1) -
DRY WEIGHT OF LARVAE AFTER 7 DAYS OF EXPOSURE

Data Transformation:

None

Chi-Square Test for Normality:

| | |
|----------------------------|---|
| Calculated test statistic: | 3.84 |
| Alpha value: | 0.01 |
| Critical value: | 13.28 |
| Conclusion: | Fail to reject the null hypothesis that the data are normally distributed |

Bartlett's Test for Homogeneity of Variances:

| | |
|----------------------------|--|
| Calculated test statistic: | 2.75 |
| Alpha value: | 0.01 |
| Critical value: | 11.34 |
| Conclusion: | Fail to reject the null hypothesis that the variances are homogenous |

ANOVA:

| | |
|----------------------------|--|
| Calculated test statistic: | 6.00 |
| Alpha value: | 0.05 |
| Critical Value: | 3.49 |
| Conclusion: | Reject the null hypothesis that all groups are equal |

Dunnett's Test:

| | |
|----------------------------|--|
| Calculated test statistic: | See Table A28-5 |
| Alpha value: | 0.05 |
| Critical value: | 2.29 |
| Conclusion: | Reject the null hypothesis that all groups are equal |

TABLE A28-5. FATHEAD MINNOW BEACH POINT BUFFERED GROUNDWATER TOXICITY TEST STATISTICAL ANALYSIS (TEST NO. 1) - RESULTS OF DUNNETT'S TEST ON DRY WEIGHT OF LARVAE AFTER 7 DAYS OF EXPOSURE

| Conc (% by Vol) | No. of Reps | Mean Dry Weight (mg) | T Statistic | Significance |
|-----------------------|-------------------|----------------------------|-------------|--------------|
| UMD/WREC Control | 4 | 0.69 | | |
| 32 | 4 | 0.65 | 0.994 | |
| 56 | 4 | 0.67 | 0.458 | |
| 100 | 4 | 0.54 | 3.873 | * |

* Significantly different at alpha = 0.05 (Dunnett's critical value = 2.29).

APPENDIX 29

FATHEAD MINNOW ACUTE AND 7-DAY SURVIVAL AND GROWTH TEST CONDUCTED ON BEACH POINT RAW (pH \approx 4) GROUNDWATER (WELL CC-33B) (TEST NO. 2)

| | |
|---------------------------|---|
| Test Method: | EPA/600/4-89/001 (Weber et al., 1989) |
| Type of Test: | Static renewal (every 24 h) |
| Date: | June 2-9, 1993 |
| Investigator: | S.D. Turley |
| Laboratory: | UMD/WREC |
| Groundwater: | |
| Source: | APG-EA Beach Point Well CC-33B |
| Chemical Characteristics: | See Appendix 48 |
| Dilution Water: | |
| Source: | 20% Perrier: 80% RO water |
| Chemical Characteristics: | See Table A29-1 |
| Test Organism: | |
| Scientific Name: | <u>Pimephales promelas</u> |
| Dry Weight: | 0.55 mg (mean weight of controls at end of test) |
| Age at Start of Test: | <24 h |
| Source: | Chesapeake Cultures Hayes, VA |
| Experimental Chambers: | |
| Material: | 600 mL glass beaker |
| Test Solution Volume: | 400 mL |
| No. Organisms/Replicate: | 10 |
| No. Organisms/Treatment: | 40 |
| Loading: | <0.5 g/L |
| Lighting: | Fluorescent; 60-85 foot candles |

| | |
|----------------|-----------------------|
| Aeration: | Prior to each renewal |
| Endpoints: | Mortality; growth |
| Water Quality: | Table A29-1 |

Results:

Mortality:

96-h Exposure:

The data for the 96-h LC50 were obtained from the 7-d study. All organisms exposed to the 100% and 56% raw groundwater by volume treatment died during the 96-h exposure (Table A29-2). All but two larvae died at the next dilution of 32% groundwater by volume. No organisms died in the remaining treatments. The 96-h LC50, which was determined by the moving average angle method, is as follows:

96-h LC50 = 21.6% raw groundwater by volume (95% confidence limits = 19.27-24.01).

7-d Exposure:

Significant ($\alpha = 0.05$) mortality occurred in fathead minnow larvae exposed to 32, 56, and 100% raw groundwater for 7 d (see Tables A29-3, A29-4, and A29-5). The 7-d LC50, which was determined by the moving average angle method, is as follows:

7-d LC50 = 21.6% raw groundwater by volume (95% confidence limits = 19.27-24.01).

Growth:

A significant ($\alpha = 0.05$) reduction in growth occurred in fathead minnow larvae exposed for 7 d to 18% raw groundwater by volume (see Tables A29-3, A29-6, and A29-7). The NOEC and LOEC for the larval fish, based on reduced growth, are as follows:

NOEC = 10% raw groundwater by volume.

LOEC = 18% raw groundwater by volume.

TABLE A29-1. SUMMARY OF THE BEACH POINT RAW GROUNDWATER BIOASSAY
TEST WATER QUALITY DATA FOR THE FATHEAD MINNOW 7-
DAY TEST (TEST NO. 2) - DISSOLVED OXYGEN (MG/L)

| | <u>Test Concentrations (Percent Groundwater by Volume)</u> | | | | | |
|--------------|--|-----|-----|-----|-----|-----|
| | 0 | 10 | 18 | 32 | 56 | 100 |
| <u>Day 0</u> | | | | | | |
| 0 H | 8.3 | 8.1 | 8.1 | 8.0 | 7.9 | 6.6 |
| <u>Day 1</u> | | | | | | |
| 0 H | 8.0 | 8.0 | 7.7 | 7.4 | 7.5 | 7.1 |
| 24 H | 7.2 | 7.2 | 7.4 | 7.5 | 7.6 | 7.7 |
| <u>Day 2</u> | | | | | | |
| 0 H | 8.3 | 8.0 | 7.9 | 7.6 | 7.4 | |
| 24 H | 7.1 | 7.3 | 7.5 | 7.8 | | |
| <u>Day 3</u> | | | | | | |
| 0 H | 8.1 | 8.0 | 7.9 | 7.9 | | |
| 24 H | 7.0 | 6.8 | 6.7 | 6.8 | | |
| <u>Day 4</u> | | | | | | |
| 0 H | 8.4 | 8.2 | 8.2 | 8.2 | | |
| 24 H | 6.7 | 6.7 | 6.7 | 6.7 | | |
| <u>Day 5</u> | | | | | | |
| 0 H | 8.2 | 8.1 | 8.1 | 8.0 | | |
| 24 H | 7.3 | 7.5 | 7.3 | 7.3 | | |
| <u>Day 6</u> | | | | | | |
| 0 H | 7.8 | 7.8 | 7.7 | 8.0 | | |
| 24 H | 7.2 | 7.3 | 7.1 | 7.2 | | |
| <u>Day 7</u> | | | | | | |
| 24 H | 7.3 | 7.2 | 7.3 | 7.0 | | |

TABLE A29-1. (CONTINUED) - pH (STANDARD UNITS)

| | <u>Test Concentrations (Percent Groundwater by Volume)</u> | | | | | |
|--------------|--|------|------|------|------|------|
| | 0 | 10 | 18 | 32 | 56 | 100 |
| <u>Day 0</u> | | | | | | |
| 0 H | 6.32 | 6.31 | 6.34 | 6.36 | 6.31 | 3.97 |
| <u>Day 1</u> | | | | | | |
| 0 H | 6.50 | 6.75 | 6.76 | 6.74 | 6.57 | 3.93 |
| 24 H | 6.83 | 6.88 | 6.95 | 6.98 | 6.98 | 4.68 |
| <u>Day 2</u> | | | | | | |
| 0 H | 6.92 | 6.99 | 6.95 | 6.89 | 6.58 | |
| 24 H | 6.76 | 6.79 | 6.85 | 6.89 | | |
| <u>Day 3</u> | | | | | | |
| 0 H | 7.02 | 6.99 | 6.92 | 6.86 | | |
| 24 H | 6.82 | 6.87 | 6.87 | 6.94 | | |
| <u>Day 4</u> | | | | | | |
| 0 H | 6.90 | 6.96 | 6.92 | 6.84 | | |
| 24 H | 6.90 | 6.81 | 6.86 | 6.82 | | |
| <u>Day 5</u> | | | | | | |
| 0 H | 6.38 | 6.50 | 6.46 | 6.41 | | |
| 24 H | 6.56 | 6.68 | 6.77 | 6.81 | | |
| <u>Day 6</u> | | | | | | |
| 0 H | 7.25 | 7.29 | 7.18 | 6.79 | | |
| 24 H | 6.68 | 6.72 | 6.80 | 6.88 | | |
| <u>Day 7</u> | | | | | | |
| 24 H | 6.71 | 6.75 | 6.75 | 6.77 | | |

TABLE A29-1. (CONTINUED) - CONDUCTIVITY (μ MHOS/CM)

| <u>Test Concentrations (Percent Groundwater by Volume)</u> | | |
|--|----------|------------|
| | <u>0</u> | <u>100</u> |
| <u>Day 0</u> | | |
| 0 H | 150 | 1350 |
| <u>Day 1</u> | | |
| 0 H | 150 | 1600 |
| <u>Day 2</u> | | |
| 0 H | 185 | |
| <u>Day 3</u> | | |
| 0 H | 140 | |
| <u>Day 4</u> | | |
| 0 H | 140 | |
| <u>Day 5</u> | | |
| 0 H | 130 | |
| <u>Day 6</u> | | |
| 0 H | 140 | |
| <u>Day 7</u> | | |
| 24 H | 155 | |

TABLE A29-1. (CONTINUED) - ALKALINITY (MG/L AS CaCO₃)

| <u>Test Concentrations (Percent Groundwater by Volume)</u> | | |
|--|----------|------------|
| | <u>0</u> | <u>100</u> |
| <u>Day 0</u> | | |
| 0 H | 55 | a |
| <u>Day 1</u> | | |
| 0 H | 55 | a |
| <u>Day 2</u> | | |
| 0 H | 55 | |
| <u>Day 3</u> | | |
| 0 H | 55 | |
| <u>Day 4</u> | | |
| 0 H | 55 | |
| <u>Day 5</u> | | |
| 0 H | 50 | |
| <u>Day 6</u> | | |
| 0 H | 50 | |
| <u>Day 7</u> | | |
| 24 H | 50 | |

^a Could not obtain measurement.

TABLE A29-1. (CONTINUED) - HARDNESS (MG/L AS CaCO₃)

| <u>Test Concentrations (Percent Groundwater by Volume)</u> | | |
|--|----------|------------|
| | <u>0</u> | <u>100</u> |
| <u>Day 0</u> | | |
| 0 H | 50 | a |
| <u>Day 1</u> | | |
| 0 H | 54 | a |
| <u>Day 2</u> | | |
| 0 H | 60 | |
| <u>Day 3</u> | | |
| 0 H | 44 | |
| <u>Day 4</u> | | |
| 0 H | 54 | |
| <u>Day 5</u> | | |
| 0 H | 40 | |
| <u>Day 6</u> | | |
| 0 H | 44 | |
| <u>Day 7</u> | | |
| 24 H | 46 | |

a Could not obtain measurement.

TABLE A29-2. FATHEAD MINNOW BEACH POINT RAW GROUNDWATER TOXICITY
TEST DATA (TEST NO. 2) - SURVIVAL AFTER 96 HOURS OF
EXPOSURE

| Concentration (% Groundwater by Volume) | Rep | Number Tested | No. Alive at 96 Hours | Percent Alive |
|---|-----|------------------|--------------------------|------------------|
| UMD/WREC Control | A | 10 | 10 | 100 |
| | B | 10 | 10 | 100 |
| | C | 10 | 10 | 100 |
| | D | 10 | 10 | 100 |
| 10 | A | 10 | 10 | 100 |
| | B | 10 | 10 | 100 |
| | C | 10 | 9 | 90 |
| | D | 10 | 10 | 100 |
| 18 | A | 10 | 10 | 100 |
| | B | 10 | 8 | 80 |
| | C | 10 | 8 | 80 |
| | D | 10 | 8 | 80 |
| 32 | A | 10 | 1 | 10 |
| | B | 10 | 0 | 0 |
| | C | 10 | 1 | 10 |
| | D | 10 | 0 | 0 |
| 56 | A | 10 | 0 | 0 |
| | B | 10 | 0 | 0 |
| | C | 10 | 0 | 0 |
| | D | 10 | 0 | 0 |
| 100 | A | 10 | 0 | 0 |
| | B | 10 | 0 | 0 |
| | C | 10 | 0 | 0 |
| | D | 10 | 0 | 0 |

TABLE A29-3. FATHEAD MINNOW BEACH POINT RAW GROUNDWATER TOXICITY
TEST DATA (TEST NO. 2) - LARVAL SURVIVAL AND DRY
WEIGHT AFTER 7 DAYS OF EXPOSURE

| Conc (% by Vol) | Rep | Number Larvae Alive | Percent Survival | Dry Weight (mg) | Mean Dry Weight (mg) |
|----------------------------|-----|---------------------------|---------------------|--------------------|----------------------------|
| UMD/WREC Control | 1 | 10 | 100 | 0.56 | 0.55 |
| | 2 | 10 | 100 | 0.53 | |
| | 3 | 10 | 100 | 0.55 | |
| | 4 | 10 | 100 | 0.57 | |
| APG-EA Diluent Water | 1 | 10 | 100 | 0.57 | 0.59 |
| | 2 | 10 | 100 | 0.58 | |
| | 3 | 10 | 100 | 0.61 | |
| | 4 | 10 | 100 | 0.58 | |
| 10 | 1 | 10 | 100 | 0.50 | 0.52 |
| | 2 | 10 | 100 | 0.53 | |
| | 3 | 9 | 90 | 0.52 | |
| | 4 | 10 | 100 | 0.52 | |
| 18 | 1 | 10 | 100 | 0.44 | 0.42 |
| | 2 | 8 | 80 | 0.45 | |
| | 3 | 8 | 80 | 0.42 | |
| | 4 | 8 | 80 | 0.37 | |
| 32 | 1 | 0 | 0 | 0.11 | |
| | 2 | 0 | 0 | | |
| | 3 | 1 | 10 | | |
| | 4 | 0 | 0 | | |
| 56 | 1 | 0 | 0 | | |
| | 2 | 0 | 0 | | |
| | 3 | 0 | 0 | | |
| | 4 | 0 | 0 | | |
| 100 | 1 | 0 | 0 | | |
| | 2 | 0 | 0 | | |
| | 3 | 0 | 0 | | |
| | 4 | 0 | 0 | | |

TABLE A29-4. FATHEAD MINNOW BEACH POINT RAW GROUNDWATER TOXICITY
TEST STATISTICAL ANALYSIS (TEST NO. 2) - LARVAL
SURVIVAL AFTER 7 DAYS OF EXPOSURE^a

Data Transformation:

Arc sine square root

Chi-Square Test for Normality:

| | |
|----------------------------|---|
| Calculated test statistic: | 18.86 |
| Alpha value: | 0.01 |
| Critical value: | 13.28 |
| Conclusion: | Reject the null hypothesis that the data are normally distributed |

Steel's Many-One Rank Test:

| | |
|----------------------------|--|
| Calculated test statistic: | See Table A29-5 |
| Alpha value: | 0.05 |
| Critical Value: | 10.0 |
| Conclusion: | Reject the null hypothesis that all groups are equal |

^a The 56% and 100% raw Beach Point groundwater treatments were not included in the statistical analyses because all organisms died during the test.

TABLE A29-5. FATHEAD MINNOW BEACH POINT RAW GROUNDWATER TOXICITY TEST STATISTICAL ANALYSIS (TEST NO. 2) - RESULTS OF STEEL'S MANY-ONE RANK TEST ON LARVAL SURVIVAL AFTER 7 DAYS OF EXPOSURE

| Conc (% by Vol) | No. of Reps | Mean Survival (%) ^a | Rank Sum | Significance |
|----------------------------|-------------------|--------------------------------------|-------------|--------------|
| UMD/WREC Control | 4 | 100 | | |
| APG-EA Diluent Water | 4 | 100 | 18.0 | |
| 10 | 4 | 97.5 | 16.0 | |
| 18 | 4 | 85.0 | 12.0 | |
| 32 | 4 | 2.5 | 10.0 | * |

^a Values given are actual percent survival means rather than arc sine square root transformed means which were used in the statistical analysis.

* Significantly different at alpha = 0.05 (Steel's critical value = 10.0).

TABLE A29-6. FATHEAD MINNOW BEACH POINT RAW GROUNDWATER TOXICITY
TEST STATISTICAL ANALYSIS (TEST NO. 2) - DRY WEIGHT
OF LARVAE AFTER 7 DAYS OF EXPOSURE^a

Data Transformation:

None

Chi-Square Test for Normality:

| | |
|----------------------------|---|
| Calculated test statistic: | 2.61 |
| Alpha value: | 0.01 |
| Critical value: | 13.28 |
| Conclusion: | Fail to reject the null hypothesis that the data are normally distributed |

Bartlett's Test for Homogeneity of Variances:

| | |
|----------------------------|--|
| Calculated test statistic: | 3.43 |
| Alpha value: | 0.01 |
| Critical value: | 11.34 |
| Conclusion: | Fail to reject the null hypothesis that the variances are homogenous |

ANOVA:

| | |
|----------------------------|--|
| Calculated test statistic: | 40.80 |
| Alpha value: | 0.05 |
| Critical value: | 3.49 |
| Conclusion: | Reject the null hypothesis that all groups are equal |

Dunnett's Test:

| | |
|----------------------------|--|
| Calculated test statistic: | See Table A29-7 |
| Alpha value: | 0.05 |
| Critical value: | 2.29 |
| Conclusion: | Reject the null hypothesis that all groups are equal |

^a The 32% buffered Beach Point groundwater treatment was not included in the statistical analyses because all concentrations above the NOEC for survival are excluded from the hypothesis test for growth effects.

TABLE A29-7. FATHEAD MINNOW BEACH POINT RAW GROUNDWATER TOXICITY TEST STATISTICAL ANALYSIS (TEST NO. 2) - RESULTS OF DUNNETT'S TEST ON DRY WEIGHT OF LARVAE AFTER 7 DAYS OF EXPOSURE

| Conc (% by Vol) | No. of Reps | Mean Dry Weight (mg) | T Statistic | Significance |
|----------------------------|-------------------|----------------------------|-------------|--------------|
| UMD/WREC Control | 4 | 0.55 | | |
| APG-EA Diluent Water | 4 | 0.59 | -2.06 | |
| 10 | 4 | 0.52 | 2.21 | |
| 18 | 4 | 0.42 | 8.38 | * |

* Significantly different at alpha = 0.05 (Dunnett's critical value = 2.29).

APPENDIX 30

FATHEAD MINNOW ACUTE AND 7-DAY SURVIVAL AND GROWTH
TEST CONDUCTED ON BEACH POINT BUFFERED (pH \approx 7)
GROUNDWATER (WELL CC-33B)
(TEST NO. 2)

| | |
|---------------------------|---|
| Test Method: | EPA/600/4-89/001 (Weber et al., 1989) |
| Type of Test: | Static renewal (every 24 h) |
| Date: | June 2-9, 1993 |
| Investigator: | S.D. Turley |
| Laboratory: | UMD/WREC |
| Groundwater: | |
| Source: | APG-EA Beach Point Well CC-33B |
| Chemical Characteristics: | See Appendix 48 |
| Dilution Water: | |
| Source: | 20% Perrier: 80% RO water |
| Chemical Characteristics: | See Table A30-1 |
| Test Organism: | |
| Scientific Name: | <u>Pimephales promelas</u> |
| Dry Weight: | 0.55 mg (mean weight of controls at end of test) |
| Age at Start of Test: | <24 h |
| Source: | Chesapeake Cultures Hayes, VA |
| Experimental Chambers: | |
| Material: | 600 mL glass beaker |
| Test Solution Volume: | 400 mL |
| No. Organisms/Replicate: | 10 |
| No. Organisms/Treatment: | 40 |
| Loading: | <0.5 g/L |
| Lighting: | Fluorescent; 60-85 foot candles |

| | |
|----------------|-------------------|
| Aeration: | Prior to renewals |
| pH Buffer: | 10 N NaOH |
| Endpoints: | Mortality; growth |
| Water Quality: | Table A30-1 |

Results:

Mortality:

96-h Exposure:

The data for the 96-h LC50 were obtained from the 7-d study. Fifteen percent of the larvae exposed to 100% buffered groundwater died during the 96-h exposure; thus, a LC50 could not be calculated. The data are summarized in Table A30-2.

7-d Exposure:

Significant ($\alpha = 0.05$) mortality occurred in fathead minnow larvae exposed to 100% raw groundwater for 7 d (see Tables A30-3, A30-4, and A30-5). However, less than 50% mortality occurred to the fathead minnow exposed for 7 days to 100% Beach Point buffered groundwater; thus, an LC50 could not be calculated (Table A30-3). The NOEC and LOEC for the larval fish, based on mortality are as follows:

NOEC = 56% raw groundwater by volume.
LOEC = 100% raw groundwater by volume.

Growth:

Growth (dry weight) of the fathead minnow larvae was not affected by 7 days of exposure to buffered groundwater (see Tables A30-3, and A30-6).

TABLE A30-1. SUMMARY OF THE BEACH POINT BUFFERED GROUNDWATER
BIOASSAY TEST WATER QUALITY DATA FOR THE FATHEAD
MINNOW 7-DAY TEST (TEST NO. 21) - DISSOLVED OXYGEN
(MG/L)

| | | <u>Test Concentrations (Percent Groundwater by Volume)</u> | | | |
|--------------|-----|--|-----|-----|-----|
| | | 0 | 32 | 56 | 100 |
| <u>Day 0</u> | | | | | |
| 0 H | 8.3 | 7.8 | 7.6 | 6.5 | |
| <u>Day 1</u> | | | | | |
| 0 H | 8.0 | 7.3 | 6.6 | 6.2 | |
| 24 H | 7.2 | 7.1 | 7.1 | 7.1 | |
| <u>Day 2</u> | | | | | |
| 0 H | 8.3 | 8.0 | 8.0 | 6.2 | |
| 24 H | 7.1 | 7.2 | 7.3 | 7.3 | |
| <u>Day 3</u> | | | | | |
| 0 H | 8.1 | 7.6 | 7.5 | 7.5 | |
| 24 H | 7.0 | 7.3 | 7.2 | 7.2 | |
| <u>Day 4</u> | | | | | |
| 0 H | 8.4 | 8.1 | 8.1 | 8.0 | |
| 24 H | 6.7 | 7.2 | 7.1 | 7.2 | |
| <u>Day 5</u> | | | | | |
| 0 H | 8.2 | 7.9 | 7.9 | 8.0 | |
| 24 H | 7.3 | 7.0 | 7.0 | 7.1 | |
| <u>Day 6</u> | | | | | |
| 0 H | 7.8 | 7.8 | 7.8 | 7.6 | |
| 24 H | 7.2 | 7.0 | 7.1 | 7.0 | |
| <u>Day 7</u> | | | | | |
| 24 H | 7.3 | 7.1 | 7.0 | 7.0 | |

TABLE A30-1. (CONTINUED) - pH (STANDARD UNITS)

| | <u>Test Concentrations (Percent Groundwater by Volume)</u> | | | |
|--------------|--|------|------|------|
| | 0 | 32 | 56 | 100 |
| <u>Day 0</u> | | | | |
| 0 H | 6.32 | 6.75 | 6.92 | 6.96 |
| <u>Day 1</u> | | | | |
| 0 H | 6.50 | 6.94 | 7.06 | 7.13 |
| 24 H | 6.83 | 6.64 | 6.89 | 6.95 |
| <u>Day 2</u> | | | | |
| 0 H | 6.92 | 7.08 | 7.34 | 7.60 |
| 24 H | 6.76 | 6.91 | 7.09 | 7.29 |
| <u>Day 3</u> | | | | |
| 0 H | 7.07 | 7.13 | 7.23 | 7.32 |
| 24 H | 6.82 | 7.04 | 7.20 | 7.41 |
| <u>Day 4</u> | | | | |
| 0 H | 6.90 | 6.84 | 6.96 | 7.18 |
| 24 H | 6.90 | 7.03 | 7.19 | 7.38 |
| <u>Day 5</u> | | | | |
| 0 H | 6.38 | 7.26 | 7.33 | 7.57 |
| 24 H | 6.56 | 6.99 | 7.11 | 7.25 |
| <u>Day 6</u> | | | | |
| 0 H | 7.25 | 7.07 | 7.17 | 7.23 |
| 24 H | 6.68 | 7.03 | 7.17 | 7.39 |
| <u>Day 7</u> | | | | |
| 24 H | 6.71 | 7.05 | 7.13 | 7.26 |

TABLE A30-1. (CONTINUED) - CONDUCTIVITY (μ MHOS/CM)

| <u>Test Concentrations (Percent Groundwater by Volume)</u> | | |
|--|----------|------------|
| | <u>0</u> | <u>100</u> |
| <u>Day 0</u> | | |
| 0 H | 150 | 1400 |
| <u>Day 1</u> | | |
| 0 H | 150 | 1600 |
| <u>Day 2</u> | | |
| 0 H | 185 | 1500 |
| <u>Day 3</u> | | |
| 0 H | 140 | 1700 |
| <u>Day 4</u> | | |
| 0 H | 140 | 1350 |
| <u>Day 5</u> | | |
| 0 H | 130 | 1700 |
| <u>Day 6</u> | | |
| 0 H | 140 | 1550 |
| <u>Day 7</u> | | |
| 24 H | 150 | 1650 |

TABLE A30-1. (CONTINUED) - ALKALINITY (MG/L AS CaCO_3)

| | <u>Test Concentrations (Percent Groundwater by Volume)</u> | |
|--------------|--|-----|
| | 0 | 100 |
| <u>Day 0</u> | | |
| 0 H | 55 | 100 |
| <u>Day 1</u> | | |
| 0 H | 55 | 120 |
| <u>Day 2</u> | | |
| 0 H | 55 | 110 |
| <u>Day 3</u> | | |
| 0 H | 55 | 125 |
| <u>Day 4</u> | | |
| 0 H | 55 | 125 |
| <u>Day 5</u> | | |
| 0 H | 50 | 125 |
| <u>Day 6</u> | | |
| 0 H | 50 | 100 |
| <u>Day 7</u> | | |
| 24 H | 50 | 110 |

TABLE A30-1. (CONTINUED) - HARDNESS (MG/L AS CaCO₃)

| <u>Test Concentrations (Percent Groundwater by Volume)</u> | | |
|--|----------|------------|
| | <u>0</u> | <u>100</u> |
| <u>Day 0</u> | | |
| 0 H | 50 | 170 |
| <u>Day 1</u> | | |
| 0 H | 54 | 128 |
| <u>Day 2</u> | | |
| 0 H | 60 | 148 |
| <u>Day 3</u> | | |
| 0 H | 44 | 134 |
| <u>Day 4</u> | | |
| 0 H | 54 | 154 |
| <u>Day 5</u> | | |
| 0 H | 40 | 170 |
| <u>Day 6</u> | | |
| 0 H | 44 | 164 |
| <u>Day 7</u> | | |
| 24 H | 50 | 140 |

TABLE A30-2. FATHEAD MINNOW BEACH POINT BUFFERED GROUNDWATER
TOXICITY TEST DATA (TEST NO. 2) - SURVIVAL AFTER 96
HOURS OF EXPOSURE

| Concentration (% Groundwater by Volume) | Rep | Number Tested | No. Alive at 96 Hours | Percent Alive |
|---|-----|------------------|--------------------------|------------------|
| UMD/WREC Control | A | 10 | 10 | 100 |
| | B | 10 | 10 | 100 |
| | C | 10 | 10 | 100 |
| | D | 10 | 10 | 100 |
| 10 | A | a | | |
| | B | a | | |
| | C | a | | |
| | D | a | | |
| 18 | A | a | | |
| | B | a | | |
| | C | a | | |
| | D | a | | |
| 32 | A | 10 | 9 | 90 |
| | B | 10 | 10 | 100 |
| | C | 10 | 9 | 90 |
| | D | 10 | 9 | 90 |
| 56 | A | 10 | 10 | 100 |
| | B | 10 | 10 | 100 |
| | C | 10 | 10 | 100 |
| | D | 10 | 10 | 100 |
| 100 | A | 10 | 7 | 70 |
| | B | 10 | 9 | 90 |
| | C | 10 | 9 | 90 |
| | D | 10 | 9 | 90 |

^a Larvae were not tested at this treatment.

TABLE A30-3. FATHEAD MINNOW BEACH POINT BUFFERED GROUNDWATER
TOXICITY TEST DATA (TEST NO. 2) - LARVAL SURVIVAL
AND DRY WEIGHT AFTER 7 DAYS OF EXPOSURE

| Conc (% by Vol) | Rep | Number Larvae Alive | Percent Survival | Dry Weight (mg) | Mean Dry Weight (mg) |
|-----------------------|-----|---------------------------|---------------------|--------------------|----------------------------|
| UMD/WREC Control | 1 | 10 | 100 | 0.56 | 0.55 |
| | 2 | 10 | 100 | 0.53 | |
| | 3 | 10 | 100 | 0.55 | |
| | 4 | 10 | 100 | 0.57 | |
| 32 | 1 | 9 | 90 | 0.54 | 0.53 |
| | 2 | 9 | 90 | 0.55 | |
| | 3 | 10 | 100 | 0.53 | |
| | 4 | 8 | 80 | 0.50 | |
| 56 | 1 | 9 | 90 | 0.48 | 0.53 |
| | 2 | 9 | 90 | 0.55 | |
| | 3 | 10 | 100 | 0.51 | |
| | 4 | 10 | 100 | 0.57 | |
| 100 | 1 | 6 | 60 | 0.54 | 0.50 |
| | 2 | 9 | 90 | 0.48 | |
| | 3 | 8 | 80 | 0.48 | |
| | 4 | 9 | 90 | 0.51 | |

TABLE A30-4. FATHEAD MINNOW BEACH POINT BUFFERED GROUNDWATER
TOXICITY TEST STATISTICAL ANALYSIS (TEST NO. 2) -
LARVAL SURVIVAL AFTER 7 DAYS OF EXPOSURE

Data Transformation:

Arc sine square root

Chi-Square Test for Normality:

| | |
|----------------------------|---|
| Calculated test statistic: | 2.61 |
| Alpha value: | 0.01 |
| Critical value: | 13.28 |
| Conclusion: | Fail to reject the null hypothesis that the data are normally distributed |

Bartlett's Test for Homogeneity of Variances:

Data failed to meet homogeneity of variance test because at least one group had zero variance.

Steel' Many One Rank Test:

| | |
|----------------------------|--|
| Calculated test statistic: | See Table A30-5 |
| Alpha value: | 0.05 |
| Critical Value: | 10.0 |
| Conclusion: | Reject the null hypothesis that all groups are equal |

TABLE A30-5. FATHEAD MINNOW BEACH POINT BUFFERED GROUNDWATER TOXICITY TEST STATISTICAL ANALYSIS (TEST NO. 2) - RESULTS OF STEEL' MANY ONE RANK TEST ON LARVAL SURVIVAL AFTER 7 DAYS OF EXPOSURE

| Conc (% by Vol) | No. of Reps | Mean Survival (%) ^a | Rank Sum | Significance |
|-----------------------|-------------------|--------------------------------------|-------------|--------------|
| UMD/WREC Control | 4 | 100 | | |
| 32 | 4 | 90 | 12.0 | |
| 56 | 4 | 95 | 14.0 | |
| 100 | 4 | 80 | 10.0 | * |

^a Values given are actual percent survival means rather than arc sine square root transformed means which were used in the statistical analysis.

* Significantly different at alpha = 0.05 (Steel's critical value = 10.0).

TABLE A30-5. FATHEAD MINNOW BEACH POINT BUFFERED GROUNDWATER
TOXICITY TEST STATISTICAL ANALYSIS (TEST NO. 2) -
DRY WEIGHT OF LARVAE AFTER 7 DAYS OF EXPOSURE^a

Data Transformation:

None

Chi-Square Test for Normality:

| | |
|----------------------------|---|
| Calculated test statistic: | 2.06 |
| Alpha value: | 0.01 |
| Critical value: | 13.28 |
| Conclusion: | Fail to reject the null hypothesis that the data are normally distributed |

Bartlett's Test for Homogeneity of Variances:

| | |
|----------------------------|--|
| Calculated test statistic: | 2.14 |
| Alpha value: | 0.01 |
| Critical value: | 9.21 |
| Conclusion: | Fail to reject the null hypothesis that the variances are homogenous |

ANOVA:

| | |
|----------------------------|--|
| Calculated test statistic: | 1.00 |
| Alpha value: | 0.05 |
| Critical Value: | 4.26 |
| Conclusion: | Reject the null hypothesis that all groups are equal |

^a The 100% buffered Beach Point groundwater treatment was not included in the statistical analyses because all concentrations above the NOEC for survival are excluded from the hypothesis test for growth effects.

APPENDIX 31

FATHEAD MINNOW ACUTE AND 7-DAY SURVIVAL AND GROWTH
TEST CONDUCTED ON BEACH POINT RAW (pH \approx 4)
GROUNDWATER (WELL CC-33B)
(TEST NO. 3)

| | |
|---------------------------|---|
| Test Method: | EPA/600/4-89/001 (Weber et al., 1989) |
| Type of Test: | Static renewal (every 24 h) |
| Date: | August 24-31, 1993 |
| Investigator: | S.D. Turley |
| Laboratory: | UMD/WREC |
| Groundwater: | |
| Source: | APG-EA Beach Point Well CC-33B |
| Chemical Characteristics: | See Appendix 48 |
| Dilution Water: | |
| Source: | 20% Perrier: 80% RO water |
| Chemical Characteristics: | See Table A31-1 |
| Test Organism: | |
| Scientific Name: | <u>Pimephales promelas</u> |
| Dry Weight: | 0.58 mg (mean weight of controls at end of test) |
| Age at Start of Test: | <24 h |
| Source: | UMD/WREC culture |
| Experimental Chambers: | |
| Material: | 600 mL glass beaker |
| Test Solution Volume: | 400 mL |
| No. Organisms/Replicate: | 10 |
| No. Organisms/Treatment: | 40 |
| Loading: | <0.5 g/L |
| Lighting: | Fluorescent; 60-85 foot candles |

| | |
|----------------|-----------------------|
| Aeration: | Prior to each renewal |
| Endpoints: | Mortality; growth |
| Water Quality: | Table A31-1 |

Results:

Mortality:

96-h Exposure:

The data for the 96-h LC50 were obtained from the 7-d study. All organisms exposed to the 100% raw groundwater and 87.5% of the larvae in the 56% raw groundwater by volume treatment died during the 96-h exposure (Table A31-2). The 96-h LC50, which was determined by the moving average angle method, is as follows:

96-h LC50 = 44.8% raw groundwater by volume (95% confidence limits = 40.63-49.49).

7-d Exposure:

Significant ($\alpha = 0.05$) mortality occurred in fathead minnow larvae exposed to 100% and 56% groundwater by volume for 7 d (see Tables A31-3, A31-4, and A31-5). The 7-d LC50, which was determined by the moving average angle method, is as follows:

7-d LC50 = 38.7% raw groundwater by volume (95% confidence limits = 34.22-43.55).

The NOEC and LOEC for the larval fish, based on mortality, are as follows:

NOEC = 32% raw groundwater by volume.

LOEC = 56% raw groundwater by volume.

Growth:

The growth of fathead minnow larvae was not affected by a 7-d exposure to concentrations which ranged from 10% to 32% groundwater by volume (see Tables A31-3 and A31-6).

TABLE A31-1. SUMMARY OF THE BEACH POINT RAW GROUNDWATER BIOASSAY
TEST WATER QUALITY DATA FOR THE FATHEAD MINNOW 7-
DAY TEST (TEST NO. 3) - DISSOLVED OXYGEN (MG/L)

| | | <u>Test Concentrations (Percent Groundwater by Volume)</u> | | | | | |
|--------------|--|--|-----|-----|-----|-----|-----|
| | | 0 | 10 | 18 | 32 | 56 | 100 |
| <u>Day 0</u> | | | | | | | |
| 0 H | | 8.0 | 7.7 | 7.7 | 7.5 | 6.9 | 6.3 |
| <u>Day 1</u> | | | | | | | |
| 0 H | | 8.3 | 7.9 | 7.7 | 7.5 | 6.7 | |
| 24 H | | 5.5 | 5.1 | 5.8 | 5.9 | 6.0 | 7.0 |
| <u>Day 2</u> | | | | | | | |
| 0 H | | 8.1 | 8.0 | 7.9 | 7.5 | 7.2 | |
| 24 H | | 6.0 | 6.5 | 6.0 | 6.5 | 6.5 | |
| <u>Day 3</u> | | | | | | | |
| 0 H | | 8.2 | 8.0 | 7.8 | 7.4 | 7.0 | |
| 24 H | | 6.0 | 6.3 | 6.4 | 6.6 | 6.8 | |
| <u>Day 4</u> | | | | | | | |
| 0 H | | 8.1 | 8.0 | 7.7 | 7.5 | 7.3 | |
| 24 H | | 6.0 | 6.2 | 5.8 | 6.0 | 6.1 | |
| <u>Day 5</u> | | | | | | | |
| 0 H | | 8.3 | 8.1 | 7.9 | 7.7 | 7.5 | |
| 24 H | | 6.2 | 6.1 | 5.9 | 6.3 | 5.9 | |
| <u>Day 6</u> | | | | | | | |
| 0 H | | 8.4 | 8.2 | 8.1 | 7.9 | 7.1 | |
| 24 H | | 6.1 | 6.0 | 6.0 | 6.2 | 5.9 | |
| <u>Day 7</u> | | | | | | | |
| 24 H | | 6.3 | 6.1 | 6.0 | 6.1 | 6.0 | |

TABLE A31-1. (CONTINUED) - pH (STANDARD UNITS)

| | <u>Test Concentrations (Percent Groundwater by Volume)</u> | | | | | |
|--------------|--|------|------|------|------|------|
| | 0 | 10 | 18 | 32 | 56 | 100 |
| <u>Day 0</u> | | | | | | |
| 0 H | 7.99 | 7.57 | 7.21 | 6.64 | 6.03 | 3.97 |
| <u>Day 1</u> | | | | | | |
| 0 H | 7.30 | 6.95 | 6.65 | 6.31 | 5.64 | |
| 24 H | 7.09 | 7.02 | 6.96 | 6.89 | 6.66 | 4.45 |
| <u>Day 2</u> | | | | | | |
| 0 H | 8.06 | 7.39 | 6.99 | 6.52 | 5.94 | |
| 24 H | 7.53 | 7.08 | 6.87 | 6.73 | 6.47 | |
| <u>Day 3</u> | | | | | | |
| 0 H | 7.75 | 7.41 | 7.16 | 6.73 | 6.01 | |
| 24 H | 7.09 | 7.00 | 6.90 | 6.76 | 6.33 | |
| <u>Day 4</u> | | | | | | |
| 0 H | 7.80 | 7.44 | 7.17 | 6.84 | 6.09 | |
| 24 H | 7.08 | 7.02 | 7.00 | 6.93 | 6.60 | |
| <u>Day 5</u> | | | | | | |
| 0 H | 7.57 | 7.37 | 7.07 | 6.80 | 6.01 | |
| 24 H | 7.15 | 7.00 | 7.10 | 6.87 | 6.51 | |
| <u>Day 6</u> | | | | | | |
| 0 H | 7.83 | 7.20 | 6.81 | 6.41 | 5.89 | |
| 24 H | 7.29 | 7.11 | 7.15 | 6.99 | 6.60 | |
| <u>Day 7</u> | | | | | | |
| 24 H | 7.22 | 7.13 | 7.02 | 6.89 | 6.49 | |

TABLE A31-1. (CONTINUED) - CONDUCTIVITY (μ MHOS/CM)

| <u>Test Concentrations (Percent Groundwater by Volume)</u> | | | |
|--|----------|-----------|------------|
| | <u>0</u> | <u>56</u> | <u>100</u> |
| <u>Day 0</u> | | | |
| 0 H | 170 | 1130 | 1400 |
| <u>Day 1</u> | | | |
| 0 H | 160 | 1100 | |
| <u>Day 2</u> | | | |
| 0 H | 160 | 1100 | |
| <u>Day 3</u> | | | |
| 0 H | 160 | 1050 | |
| <u>Day 4</u> | | | |
| 0 H | 160 | 1000 | |
| <u>Day 5</u> | | | |
| 0 H | 170 | 1050 | |
| <u>Day 6</u> | | | |
| 0 H | 160 | 1100 | |
| <u>Day 7</u> | | | |
| 24 H | 150 | 1150 | |

TABLE A31-1. (CONTINUED) - ALKALINITY (MG/L AS CaCO₃)

| | <u>Test Concentrations (Percent Groundwater by Volume)</u> | | |
|--------------|--|----|-----|
| | 0 | 56 | 100 |
| <u>Day 0</u> | | | |
| 0 H | 50 | 25 | a |
| <u>Day 1</u> | | | |
| 0 H | 55 | 30 | a |
| <u>Day 2</u> | | | |
| 0 H | 55 | 20 | a |
| <u>Day 3</u> | | | |
| 0 H | 50 | 25 | |
| <u>Day 4</u> | | | |
| 0 H | 55 | 30 | |
| <u>Day 5</u> | | | |
| 0 H | 50 | 25 | |
| <u>Day 6</u> | | | |
| 0 H | 55 | 20 | |
| <u>Day 7</u> | | | |
| 24 H | 55 | 25 | |

^a Could not obtain measurement.

TABLE A31-1. (CONTINUED) - HARDNESS (MG/L AS CaCO₃)

| | | <u>Test Concentrations (Percent Groundwater by Volume)</u> | | |
|--------------|----|--|----|-----|
| | | 0 | 56 | 100 |
| <u>Day 0</u> | | | | |
| 0 H | 52 | | a | a |
| <u>Day 1</u> | | | | |
| 0 H | 50 | | a | |
| <u>Day 2</u> | | | | |
| 0 H | 52 | | a | |
| <u>Day 3</u> | | | | |
| 0 H | 52 | | a | |
| <u>Day 4</u> | | | | |
| 0 H | 50 | | a | |
| <u>Day 5</u> | | | | |
| 0 H | 54 | | a | |
| <u>Day 6</u> | | | | |
| 0 H | 50 | | a | |
| <u>Day 7</u> | | | | |
| 24 H | 54 | | a | |

^a Could not obtain measurement.

TABLE A31-2. FATHEAD MINNOW BEACH POINT RAW GROUNDWATER TOXICITY
TEST DATA (TEST NO. 3) - SURVIVAL AFTER 96 HOURS OF
EXPOSURE

| Concentration (% Groundwater by Volume) | Rep | Number Tested | No. Alive at 96 Hours | Percent Alive |
|---|-----|------------------|--------------------------|------------------|
| UMD/WREC Control | A | 10 | 10 | 100 |
| | B | 10 | 10 | 100 |
| | C | 10 | 10 | 100 |
| | D | 10 | 10 | 100 |
| 10 | A | 10 | 10 | 100 |
| | B | 10 | 10 | 100 |
| | C | 10 | 9 | 90 |
| | D | 10 | 10 | 100 |
| 18 | A | 10 | 9 | 90 |
| | B | 10 | 9 | 90 |
| | C | 10 | 10 | 100 |
| | D | 10 | 9 | 90 |
| 32 | A | 10 | 9 | 90 |
| | B | 10 | 9 | 90 |
| | C | 10 | 10 | 100 |
| | D | 10 | 10 | 100 |
| 56 | A | 10 | 0 | 0 |
| | B | 10 | 0 | 0 |
| | C | 10 | 3 | 30 |
| | D | 10 | 2 | 20 |
| 100 | A | 10 | 0 | 0 |
| | B | 10 | 0 | 0 |
| | C | 10 | 0 | 0 |
| | D | 10 | 0 | 0 |

TABLE A31-4. FATHEAD MINNOW BEACH POINT RAW GROUNDWATER TOXICITY
TEST DATA (TEST NO. 3) - LARVAL SURVIVAL AND DRY
WEIGHT AFTER 7 DAYS OF EXPOSURE

| Conc (% by Vol) | Rep | Number Larvae Alive | Percent Survival | Dry Weight (mg) | Mean Dry Weight (mg) |
|----------------------------|-----|---------------------------|---------------------|--------------------|----------------------------|
| UMD/WREC Control | 1 | 10 | 100 | 0.56 | 0.58 |
| | 2 | 10 | 100 | 0.57 | |
| | 3 | 10 | 100 | 0.59 | |
| | 4 | 10 | 100 | 0.60 | |
| APG-EA Diluent Water | 1 | 10 | 100 | 0.59 | 0.57 |
| | 2 | 10 | 100 | 0.56 | |
| | 3 | 9 | 90 | 0.57 | |
| | 4 | 10 | 100 | 0.57 | |
| 10 | 1 | 10 | 100 | 0.53 | 0.55 |
| | 2 | 9 | 90 | 0.57 | |
| | 3 | 9 | 90 | 0.59 | |
| | 4 | 10 | 100 | 0.52 | |
| 18 | 1 | 8 | 80 | 0.57 | 0.56 |
| | 2 | 9 | 90 | 0.58 | |
| | 3 | 10 | 100 | 0.51 | |
| | 4 | 9 | 90 | 0.56 | |
| 32 | 1 | 9 | 90 | 0.52 | 0.54 |
| | 2 | 8 | 80 | 0.54 | |
| | 3 | 7 | 70 | 0.54 | |
| | 4 | 10 | 100 | 0.55 | |
| 56 | 1 | 0 | 00 | 0.00 | 0.29 |
| | 2 | 0 | 00 | 0.00 | |
| | 3 | 3 | 30 | 0.30 | |
| | 4 | 2 | 20 | 0.28 | |
| 100 | 1 | 0 | 0 | | |
| | 2 | 0 | 0 | | |
| | 3 | 0 | 0 | | |
| | 4 | 0 | 0 | | |

TABLE A31-4. FATHEAD MINNOW BEACH POINT RAW GROUNDWATER TOXICITY
TEST STATISTICAL ANALYSIS (TEST NO. 3) - LARVAL
SURVIVAL AFTER 7 DAYS OF EXPOSURE^a

Data Transformation:

Arc sine square root

Chi-Square Test for Normality:

| | |
|----------------------------|---|
| Calculated test statistic: | 4.10 |
| Alpha value: | 0.01 |
| Critical Value: | 13.28 |
| Conclusion: | Fail to reject the null hypothesis that the data are normally distributed |

Bartlett's Test for Homogeneity of Variances:

Data failed to meet homogeneity of variance test because at least one group had zero variance.

Steel's Many-One Rank Test:

| | |
|----------------------------|--|
| Calculated test statistic: | See Table A31-5 |
| Alpha value: | 0.05 |
| Critical Value: | 10.0 |
| Conclusion: | Reject the null hypothesis that all groups are equal |

^a The 100% raw Beach Point groundwater treatment was not included in the statistical analyses because all organisms died during the test.

TABLE A31-5. FATHEAD MINNOW BEACH POINT RAW GROUNDWATER TOXICITY TEST STATISTICAL ANALYSIS (TEST NO. 3) - RESULTS OF STEEL'S MANY-ONE RANK TEST ON LARVAL SURVIVAL AFTER 7 DAYS OF EXPOSURE

| Conc (% by Vol) | No. of Reps | Mean Survival (%) ^a | Rank Sum | Significance |
|----------------------------|-------------------|--------------------------------------|-------------|--------------|
| UMD/WREC Control | 4 | 100 | | |
| APG-EA Diluent Water | 4 | 97.5 | 16.0 | |
| 10 | 4 | 95.0 | 14.0 | |
| 18 | 4 | 90.0 | 12.0 | |
| 32 | 4 | 85.0 | 12.0 | |
| 56 | 4 | 12.5 | 10.0 | * |

^a Values given are actual percent survival means rather than arc sine square root transformed means which were used in the statistical analysis.

* Significantly different at alpha = 0.05 (Steel's critical value = 10.0).

TABLE A31-6. FATHEAD MINNOW BEACH POINT RAW GROUNDWATER TOXICITY
TEST STATISTICAL ANALYSIS (TEST NO. 3) - DRY WEIGHT
OF LARVAE AFTER 7 DAYS OF EXPOSURE^a

Data Transformation:

None

Chi-Square Test for Normality:

| | |
|----------------------------|---|
| Calculated test statistic: | 4.96 |
| Alpha value: | 0.01 |
| Critical value: | 13.28 |
| Conclusion: | Fail to reject the null hypothesis that the data are normally distributed |

Bartlett's Test for Homogeneity of Variances:

| | |
|----------------------------|--|
| Calculated test statistic: | 4.41 |
| Alpha value: | 0.01 |
| Critical value: | 13.28 |
| Conclusion: | Fail to reject the null hypothesis that the variances are homogenous |

ANOVA:

| | |
|----------------------------|--|
| Calculated test statistic: | 2.20 |
| Alpha value: | 0.05 |
| Critical value: | 3.06 |
| Conclusion: | Fail to reject the null hypothesis that all groups are equal |

^a The 56% buffered Beach Point groundwater treatment was not included in the statistical analyses because all concentrations above the NOEC for survival are excluded from the hypothesis test for growth effects.

APPENDIX 32

FATHEAD MINNOW ACUTE AND 7-DAY SURVIVAL AND GROWTH TEST CONDUCTED ON BEACH POINT BUFFERED (pH ≈7) GROUNDWATER (WELL CC-33B) (TEST NO. 3)

| | |
|---------------------------|---|
| Test Method: | EPA/600/4-89/001 (Weber et al., 1989) |
| Type of Test: | Static renewal (every 24 h) |
| Date: | August 24-31, 1993 |
| Investigator: | S.D. Turley |
| Laboratory: | UMD/WREC |
| Groundwater: | |
| Source: | APG-EA Beach Point Well CC-33B |
| Chemical Characteristics: | See Appendix 48 |
| Dilution Water: | |
| Source: | 20% Perrier: 80% RO water |
| Chemical Characteristics: | See Table A32-1 |
| Test Organism: | |
| Scientific Name: | <u>Pimephales promelas</u> |
| Dry Weight: | 0.58 mg (mean weight of controls at end of test) |
| Age at Start of Test: | <24 h |
| Source: | UMD/WREC culture |
| Experimental Chambers: | |
| Material: | 600 mL glass beaker |
| Test Solution Volume: | 400 mL |
| No. Organisms/Replicate: | 10 |
| No. Organisms/Treatment: | 40 |
| Loading: | <0.5 g/L |
| Lighting: | Fluorescent; 60-85 foot candles |
| Aeration: | Prior to renewals |

| | |
|----------------|-------------------|
| pH Buffer: | 10 N NaOH |
| Endpoints: | Mortality; growth |
| Water Quality: | Table A32-1 |

Results:

Mortality:

96-h Exposure:

The data for the 96-h LC50 were obtained from the 7-d study. The buffered groundwater was not toxic to the larvae. The data are summarized in Table A32-2.

7-d Exposure:

The survival of fathead minnow larvae was not affected after 7 days of exposure to Beach Point buffered groundwater (Tables A32-3 and A32-4).

Growth:

The growth of fathead minnow larvae was not affected after 7 days of exposure to Beach Point buffered groundwater (Tables A32-3 and A32-5).

TABLE A32-1. SUMMARY OF THE BEACH POINT BUFFERED GROUNDWATER
BIOASSAY TEST WATER QUALITY DATA FOR THE FATHEAD
MINNOW 7-DAY TEST (TEST NO. 3) - DISSOLVED OXYGEN
(MG/L)

| | <u>Test Concentrations (Percent Groundwater by Volume)</u> | | | | | |
|--------------|--|-----|-----|-----|-----|-----|
| | 0 | 10 | 18 | 32 | 56 | 100 |
| <u>Day 0</u> | | | | | | |
| 0 H | 8.0 | 7.8 | 7.5 | 7.2 | 7.0 | 5.5 |
| <u>Day 1</u> | | | | | | |
| 0 H | 8.3 | 7.4 | 7.5 | 7.2 | 7.0 | 7.0 |
| 24 H | 5.5 | 5.5 | 5.6 | 5.4 | 5.5 | 5.0 |
| <u>Day 2</u> | | | | | | |
| 0 H | 8.1 | 7.8 | 7.6 | 7.3 | 7.3 | 5.3 |
| 24 H | 6.0 | 5.8 | 6.2 | 6.0 | 6.0 | 6.5 |
| <u>Day 3</u> | | | | | | |
| 0 H | 8.2 | 7.7 | 7.5 | 7.2 | 7.3 | 6.1 |
| 24 H | 6.0 | 6.0 | 5.9 | 6.6 | 6.3 | 6.5 |
| <u>Day 4</u> | | | | | | |
| 0 H | 8.1 | 7.8 | 7.6 | 7.4 | 7.5 | 6.5 |
| 24 H | 6.0 | 5.6 | 5.8 | 5.6 | 6.0 | 6.0 |
| <u>Day 5</u> | | | | | | |
| 0 H | 8.3 | 7.9 | 7.7 | 7.5 | 7.4 | 6.8 |
| 24 H | 6.2 | 5.8 | 5.9 | 5.8 | 6.1 | 6.3 |
| <u>Day 6</u> | | | | | | |
| 0 H | 8.4 | 8.1 | 7.9 | 7.6 | 7.3 | 5.0 |
| 24 H | 6.1 | 6.0 | 6.0 | 5.9 | 6.0 | 6.1 |
| <u>Day 7</u> | | | | | | |
| 24 H | 6.3 | 6.1 | 6.1 | 5.8 | 5.9 | 6.0 |

TABLE A32-1. (CONTINUED) - pH (STANDARD UNITS)

| | <u>Test Concentrations (Percent Groundwater by Volume)</u> | | | | | |
|--------------|--|------|------|------|------|------|
| | 0 | 10 | 18 | 32 | 56 | 100 |
| <u>Day 0</u> | | | | | | |
| 0 H | 7.99 | 7.53 | 7.75 | 7.83 | 7.84 | 7.69 |
| <u>Day 1</u> | | | | | | |
| 0 H | 7.30 | 7.44 | 7.63 | 7.71 | 7.67 | 7.57 |
| 24 H | 7.09 | 7.16 | 7.19 | 7.22 | 7.29 | 7.41 |
| <u>Day 2</u> | | | | | | |
| 0 H | 8.06 | 7.96 | 7.89 | 7.81 | 7.60 | 7.57 |
| 24 H | 7.53 | 7.09 | 7.20 | 7.30 | 7.38 | 7.59 |
| <u>Day 3</u> | | | | | | |
| 0 H | 7.75 | 7.85 | 8.00 | 8.07 | 8.04 | 7.75 |
| 24 H | 7.09 | 7.11 | 7.24 | 7.45 | 7.43 | 7.43 |
| <u>Day 4</u> | | | | | | |
| 0 H | 7.80 | 7.77 | 7.79 | 7.90 | 7.88 | 7.57 |
| 24 H | 7.08 | 7.21 | 7.41 | 7.56 | 7.65 | 7.79 |
| <u>Day 5</u> | | | | | | |
| 0 H | 7.57 | 7.29 | 7.40 | 7.59 | 7.69 | 7.60 |
| 24 H | 7.15 | 7.21 | 7.41 | 7.55 | 7.70 | 7.84 |
| <u>Day 6</u> | | | | | | |
| 0 H | 7.85 | 7.53 | 7.69 | 7.78 | 7.79 | 7.75 |
| 24 H | 7.29 | 7.30 | 7.45 | 7.48 | 7.53 | 7.75 |
| <u>Day 7</u> | | | | | | |
| 24 H | 7.22 | 7.25 | 7.38 | 7.52 | 7.61 | 7.71 |

TABLE A32-1. (CONTINUED) - CONDUCTIVITY (μ MHOS/CM)

| <u>Test Concentrations (Percent Groundwater by Volume)</u> | | |
|--|----------|------------|
| | <u>0</u> | <u>100</u> |
| <u>Day 0</u> | | |
| 0 H | 170 | 2000 |
| <u>Day 1</u> | | |
| 0 H | 160 | 1950 |
| <u>Day 2</u> | | |
| 0 H | 160 | 1625 |
| <u>Day 3</u> | | |
| 0 H | 160 | 1800 |
| <u>Day 4</u> | | |
| 0 H | 160 | 1900 |
| <u>Day 5</u> | | |
| 0 H | 170 | 1850 |
| <u>Day 6</u> | | |
| 0 H | 160 | 1925 |
| <u>Day 7</u> | | |
| 24 H | 150 | 1900 |

TABLE A32-1. (CONTINUED) - ALKALINITY (MG/L AS CaCO₃)

| | <u>Test Concentrations (Percent Groundwater by Volume)</u> | |
|--------------|--|-----|
| | 0 | 100 |
| <u>Day 0</u> | | |
| 0 H | 50 | 90 |
| <u>Day 1</u> | | |
| 0 H | 55 | 100 |
| <u>Day 2</u> | | |
| 0 H | 55 | 75 |
| <u>Day 3</u> | | |
| 0 H | 50 | 85 |
| <u>Day 4</u> | | |
| 0 H | 55 | 80 |
| <u>Day 5</u> | | |
| 0 H | 50 | 80 |
| <u>Day 6</u> | | |
| 0 H | 55 | 110 |
| <u>Day 7</u> | | |
| 24 H | 55 | 100 |

TABLE A32-1. (CONTINUED) - HARDNESS (MG/L AS CaCO₃)

| <u>Test Concentrations (Percent Groundwater by Volume)</u> | | |
|--|----------|------------|
| | <u>0</u> | <u>100</u> |
| <u>Day 0</u> | | |
| 0 H | 52 | 228 |
| <u>Day 1</u> | | |
| 0 H | 50 | 240 |
| <u>Day 2</u> | | |
| 0 H | 52 | 194 |
| <u>Day 3</u> | | |
| 0 H | 52 | 210 |
| <u>Day 4</u> | | |
| 0 H | 50 | 220 |
| <u>Day 5</u> | | |
| 0 H | 54 | 230 |
| <u>Day 6</u> | | |
| 0 H | 50 | 164 |
| <u>Day 7</u> | | |
| 24 H | 54 | 184 |

TABLE A32-2. FATHEAD MINNOW BEACH POINT BUFFERED GROUNDWATER
TOXICITY TEST DATA (TEST NO. 3) - SURVIVAL AFTER 96
HOURS OF EXPOSURE

| Concentration (% Groundwater by Volume) | Rep | Number Tested | No. Alive at 96 Hours | Percent Alive |
|---|-----|------------------|--------------------------|------------------|
| UMD/WREC Control | A | 10 | 10 | 100 |
| | B | 10 | 10 | 100 |
| | C | 10 | 10 | 100 |
| | D | 10 | 10 | 100 |
| 10 | A | 10 | 10 | 100 |
| | B | 10 | 10 | 100 |
| | C | 10 | 9 | 90 |
| | D | 10 | 10 | 100 |
| 18 | A | 10 | 10 | 100 |
| | B | 10 | 10 | 100 |
| | C | 10 | 10 | 100 |
| | D | 10 | 10 | 100 |
| 32 | A | 10 | 9 | 90 |
| | B | 10 | 10 | 100 |
| | C | 10 | 10 | 100 |
| | D | 10 | 10 | 100 |
| 56 | A | 10 | 9 | 90 |
| | B | 10 | 10 | 100 |
| | C | 10 | 9 | 90 |
| | D | 10 | 10 | 100 |
| 100 | A | 10 | 10 | 100 |
| | B | 10 | 9 | 90 |
| | C | 10 | 8 | 80 |
| | D | 10 | 10 | 100 |

TABLE A32-3. FATHEAD MINNOW BEACH POINT BUFFERED GROUNDWATER
TOXICITY TEST DATA (TEST NO. 3) - LARVAL SURVIVAL
AND DRY WEIGHT AFTER 7 DAYS OF EXPOSURE

| Conc (% by Vol) | Rep | Number Larvae Alive | Percent Survival | Dry Weight (mg) | Mean Dry Weight (mg) |
|-----------------------|-----|---------------------------|---------------------|--------------------|----------------------------|
| UMD/WREC Control | 1 | 10 | 100 | 0.56 | 0.58 |
| | 2 | 10 | 100 | 0.57 | |
| | 3 | 10 | 100 | 0.59 | |
| | 4 | 10 | 100 | 0.60 | |
| 10 | 1 | 10 | 100 | 0.51 | 0.58 |
| | 2 | 10 | 100 | 0.59 | |
| | 3 | 10 | 100 | 0.59 | |
| | 4 | 9 | 90 | 0.61 | |
| 18 | 1 | 9 | 90 | 0.56 | 0.57 |
| | 2 | 10 | 100 | 0.59 | |
| | 3 | 10 | 100 | 0.61 | |
| | 4 | 10 | 100 | 0.53 | |
| 32 | 1 | 9 | 90 | 0.58 | 0.58 |
| | 2 | 10 | 100 | 0.60 | |
| | 3 | 10 | 100 | 0.58 | |
| | 4 | 10 | 100 | 0.56 | |
| 56 | 1 | 9 | 90 | 0.54 | 0.56 |
| | 2 | 9 | 90 | 0.54 | |
| | 3 | 10 | 100 | 0.59 | |
| | 4 | 9 | 90 | 0.56 | |
| 100 | 1 | 10 | 100 | 0.55 | 0.55 |
| | 2 | 9 | 90 | 0.56 | |
| | 3 | 8 | 80 | 0.54 | |
| | 4 | 10 | 100 | 0.56 | |

TABLE A32-4. FATHEAD MINNOW BEACH POINT BUFFERED GROUNDWATER
TOXICITY TEST STATISTICAL ANALYSIS (TEST NO. 3) -
LARVAL SURVIVAL AFTER 7 DAYS OF EXPOSURE

Data Transformation:

Arc sine square root

Chi-Square Test for Normality:

| | |
|----------------------------|---|
| Calculated test statistic: | 11.83 |
| Alpha value: | 0.01 |
| Critical Value: | 13.28 |
| Conclusion: | Fail to reject the null hypothesis that the data are normally distributed |

Bartlett's Test for Homogeneity of Variances:

Data failed to meet homogeneity of variance test because at least one group had zero variance.

Steel's Many-One Rank Test:

| | |
|-----------------------------|--|
| Calculated test statistics: | 12.0-16.0 |
| Alpha value: | 0.05 |
| Critical Value: | 10.0 |
| Conclusion: | Fail to reject the null hypothesis that all groups are equal |

TABLE A32-5. FATHEAD MINNOW BEACH POINT BUFFERED GROUNDWATER
TOXICITY TEST STATISTICAL ANALYSIS (TEST NO. 3) -
DRY WEIGHT OF LARVAE AFTER 7 DAYS OF EXPOSURE

Data Transformation:

None

Chi-Square Test for Normality:

| | |
|----------------------------|---|
| Calculated test statistic: | 5.02 |
| Alpha value: | 0.01 |
| Critical value: | 13.28 |
| Conclusion: | Fail to reject the null hypothesis that the data are normally distributed |

Bartlett's Test for Homogeneity of Variances:

| | |
|----------------------------|--|
| Calculated test statistic: | 7.19 |
| Alpha value: | 0.01 |
| Critical value: | 15.09 |
| Conclusion: | Fail to reject the null hypothesis that the variances are homogenous |

ANOVA:

| | |
|----------------------------|--|
| Calculated test statistic: | 0.86 |
| Alpha value: | 0.05 |
| Critical Value: | 2.77 |
| Conclusion: | Fail to reject the null hypothesis that all groups are equal |

APPENDIX 33

FATHEAD MINNOW ACUTE AND 7-DAY SURVIVAL AND GROWTH TEST CONDUCTED ON BEACH POINT RAW (pH \approx 4) GROUNDWATER (WELL CC-33B) (TEST NO. 4)

| | |
|---------------------------|---|
| Test Method: | EPA/600/4-89/001 (Weber et al., 1989) |
| Type of Test: | Static renewal (every 24 h) |
| Date: | November 30 - December 7, 1993 |
| Investigator: | S.D. Turley |
| Laboratory: | UMD/WREC |
| Groundwater: | |
| Source: | APG-EA Beach Point Well CC-33B |
| Chemical Characteristics: | See Appendix 48 |
| Dilution Water: | |
| Source: | 20% Perrier: 80% RO water |
| Chemical Characteristics: | See Table A33-1 |
| Test Organism: | |
| Scientific Name: | <u>Pimephales promelas</u> |
| Dry Weight: | 0.49 mg (mean dry weight of controls at end of test) |
| Age at Start of Test: | <24 h |
| Source: | UMD/WREC culture |
| Experimental Chambers: | |
| Material: | 600 mL glass beaker |
| Test Solution Volume: | 400 mL |
| No. Organisms/Replicate: | 10 |
| No. Organisms/Treatment: | 40 |
| Loading: | <0.5 g/L |
| Lighting: | Fluorescent; 60-85 foot candles |

| | |
|----------------|-----------------------|
| Aeration: | Prior to each renewal |
| Endpoints: | Mortality; growth |
| Water Quality: | Table A33-1 |

Results:

Mortality:

96-h Exposure:

The data for the 96-h LC50 were obtained from the 7-d study. All larvae exposed to the 100% raw groundwater treatment died during the 96-h exposure (Table A33-2). The 96-h LC50, which was determined by the moving average angle method, is as follows:

96-h LC50 = 61.7% raw groundwater by volume (95% confidence limits = 58.06-65.93).

7-d Exposure:

Significant ($\alpha = 0.05$) mortality occurred to fathead minnow larvae exposed to 100% and 56% groundwater by volume for 7 d (see Tables A33-3, A33-4, and A33-5). The 7-d LC50, which was determined by the moving average angle method, is as follows:

7-d LC50 = 51.3% raw groundwater by volume (95% confidence limits = 46.27-57.57).

The NOEC and LOEC for the larval fish, based on mortality, are as follows:

NOEC = 32% raw groundwater by volume.

LOEC = 56% raw groundwater by volume.

Growth:

The growth of fathead minnow larvae was not affected by a 7-d exposure to concentrations which ranged from 10% to 32% groundwater by volume (see Tables A33-3 and A33-6).

TABLE A33-1. SUMMARY OF THE BEACH POINT RAW GROUNDWATER BIOASSAY
TEST WATER QUALITY DATA FOR THE FATHEAD MINNOW 7-
DAY TEST (TEST NO. 4) - DISSOLVED OXYGEN (MG/L)

| | <u>Test Concentrations (Percent Groundwater by Volume)</u> | | | | | |
|--------------|--|-----|-----|-----|-----|-----|
| | 0 | 10 | 18 | 32 | 56 | 100 |
| <u>Day 0</u> | | | | | | |
| 0 H | 8.4 | 8.1 | 8.2 | 7.7 | 7.6 | 7.7 |
| <u>Day 1</u> | | | | | | |
| 0 H | 8.5 | 8.2 | 8.2 | 8.1 | 8.1 | 7.6 |
| 24 H | 8.4 | 8.3 | 8.5 | 8.5 | 8.5 | 8.3 |
| <u>Day 2</u> | | | | | | |
| 0 H | 8.3 | 8.1 | 8.1 | 8.1 | 8.2 | |
| 24 H | 8.2 | 8.0 | 8.2 | 8.3 | 8.1 | |
| <u>Day 3</u> | | | | | | |
| 0 H | 8.4 | 8.3 | 8.3 | 8.0 | 8.2 | |
| 24 H | 8.3 | 8.2 | 8.1 | 8.2 | 8.1 | |
| <u>Day 4</u> | | | | | | |
| 0 H | 8.6 | 8.5 | 8.4 | 8.4 | 8.3 | |
| 24 H | 8.0 | 8.0 | 7.9 | 8.0 | 8.0 | |
| <u>Day 5</u> | | | | | | |
| 0 H | 8.7 | 8.4 | 8.3 | 8.3 | 8.3 | |
| 24 H | 8.0 | 8.1 | 7.8 | 7.8 | 7.7 | |
| <u>Day 6</u> | | | | | | |
| 0 H | 8.6 | 8.5 | 8.5 | 8.4 | 8.4 | |
| 24 H | 8.1 | 6.0 | 8.1 | 8.0 | 8.0 | |
| <u>Day 7</u> | | | | | | |
| 24 H | 8.0 | 8.1 | 7.9 | 7.9 | 7.9 | |

TABLE A33-1. (CONTINUED) - pH (STANDARD UNITS)

| | <u>Test Concentrations (Percent Groundwater by Volume)</u> | | | | | |
|--------------|--|------|------|------|------|------|
| | 0 | 10 | 18 | 32 | 56 | 100 |
| <u>Day 0</u> | | | | | | |
| 0 H | 7.59 | 7.34 | 7.00 | 6.71 | 6.22 | 4.01 |
| <u>Day 1</u> | | | | | | |
| 0 H | 7.83 | 7.59 | 7.30 | 7.01 | 6.63 | 3.97 |
| 24 H | 7.57 | 7.61 | 7.59 | 7.58 | 7.39 | 5.45 |
| <u>Day 2</u> | | | | | | |
| 0 H | 7.33 | 7.22 | 7.12 | 6.91 | 6.95 | |
| 24 H | 7.59 | 7.69 | 7.67 | 7.50 | 7.26 | |
| <u>Day 3</u> | | | | | | |
| 0 H | 7.17 | 7.07 | 7.06 | 7.09 | 6.83 | |
| 24 H | 7.55 | 7.55 | 7.65 | 7.61 | 7.54 | |
| <u>Day 4</u> | | | | | | |
| 0 H | 7.79 | 7.59 | 7.55 | 7.19 | 6.66 | |
| 24 H | 7.65 | 7.71 | 7.73 | 7.75 | 7.59 | |
| <u>Day 5</u> | | | | | | |
| 0 H | 7.66 | 7.18 | 7.08 | 6.86 | 6.31 | |
| 24 H | 7.18 | 7.27 | 7.38 | 7.31 | 7.37 | |
| <u>Day 6</u> | | | | | | |
| 0 H | 7.71 | 7.62 | 7.35 | 6.93 | 6.60 | |
| 24 H | 7.42 | 7.43 | 7.48 | 7.45 | 7.37 | |
| <u>Day 7</u> | | | | | | |
| 24 H | 7.38 | 7.40 | 7.48 | 7.50 | 7.48 | |

TABLE A33-1. (CONTINUED) - CONDUCTIVITY (μ MHOS/CM)

| | <u>Test Concentrations (Percent Groundwater by Volume)</u> | |
|--------------|--|------------|
| | <u>0</u> | <u>100</u> |
| <u>Day 0</u> | | |
| 0 H | 240 | 1750 |
| <u>Day 1</u> | | |
| 0 H | 230 | 1650 |
| <u>Day 2</u> | | |
| 0 H | 240 | 1700 |
| <u>Day 3</u> | | |
| 0 H | 240 | 1700 |
| <u>Day 4</u> | | |
| 0 H | 250 | 1750 |
| <u>Day 5</u> | | |
| 0 H | 230 | 1700 |
| <u>Day 6</u> | | |
| 0 H | 210 | 1750 |
| <u>Day 7</u> | | |
| 24 H | 200 | 1800 |

TABLE A33-1. (CONTINUED) - ALKALINITY (MG/L AS CaCO₃)

| <u>Test Concentrations (Percent Groundwater by Volume)</u> | | |
|--|----------|------------|
| | <u>0</u> | <u>100</u> |
| <u>Day 0</u> | | |
| 0 H | 70 | a |
| <u>Day 1</u> | | |
| 0 H | 65 | a |
| <u>Day 2</u> | | |
| 0 H | 65 | a |
| <u>Day 3</u> | | |
| 0 H | 65 | a |
| <u>Day 4</u> | | |
| 0 H | 60 | a |
| <u>Day 5</u> | | |
| 0 H | 55 | a |
| <u>Day 6</u> | | |
| 0 H | 50 | a |
| <u>Day 7</u> | | |
| 24 H | 55 | a |

^a Could not obtain measurement.

TABLE A33-1. (CONTINUED) - HARDNESS (MG/L AS CaCO₃)

| | <u>Test Concentrations (Percent Groundwater by Volume)</u> | |
|--------------|--|-----|
| | 0 | 100 |
| <u>Day 0</u> | | |
| 0 H | 80 | a |
| <u>Day 1</u> | | |
| 0 H | 74 | a |
| <u>Day 2</u> | | |
| 0 H | 70 | a |
| <u>Day 3</u> | | |
| 0 H | 78 | a |
| <u>Day 4</u> | | |
| 0 H | 74 | a |
| <u>Day 5</u> | | |
| 0 H | 70 | a |
| <u>Day 6</u> | | |
| 0 H | 80 | a |
| <u>Day 7</u> | | |
| 24 H | 64 | a |

^a Could not obtain measurement.

TABLE A33-2. FATHEAD MINNOW BEACH POINT RAW GROUNDWATER TOXICITY
TEST DATA (TEST NO. 4) - SURVIVAL AFTER 96 HOURS OF
EXPOSURE

| Concentration (% Groundwater by Volume) | Rep | Number Tested | No. Alive at 96 Hours | Percent Alive |
|---|-----|------------------|--------------------------|------------------|
| UMD/WREC Control | A | 10 | 10 | 100 |
| | B | 10 | 10 | 100 |
| | C | 10 | 10 | 100 |
| | D | 10 | 10 | 100 |
| 10 | A | 10 | 10 | 100 |
| | B | 10 | 10 | 100 |
| | C | 10 | 10 | 100 |
| | D | 10 | 10 | 100 |
| 18 | A | 10 | 10 | 100 |
| | B | 10 | 10 | 100 |
| | C | 10 | 10 | 100 |
| | D | 10 | 9 | 90 |
| 32 | A | 10 | 10 | 100 |
| | B | 10 | 9 | 90 |
| | C | 10 | 10 | 100 |
| | D | 10 | 9 | 90 |
| 56 | A | 10 | 9 | 90 |
| | B | 10 | 10 | 100 |
| | C | 10 | 8 | 80 |
| | D | 10 | 10 | 100 |
| 100 | A | 10 | 0 | 0 |
| | B | 10 | 0 | 0 |
| | C | 10 | 0 | 0 |
| | D | 10 | 0 | 0 |

TABLE A33-3. FATHEAD MINNOW BEACH POINT RAW GROUNDWATER TOXICITY
TEST DATA (TEST NO. 4) - LARVAL SURVIVAL AND DRY
WEIGHT AFTER 7 DAYS OF EXPOSURE

| Conc (% by Vol) | Rep | Number Larvae Alive | Percent Survival | Dry Weight (mg) | Mean Dry Weight (mg) |
|----------------------------|-----|---------------------------|---------------------|--------------------|----------------------------|
| UMD/WREC Control | 1 | 10 | 100 | 0.46 | 0.49 |
| | 2 | 10 | 100 | 0.57 | |
| | 3 | 9 | 90 | 0.50 | |
| | 4 | 10 | 100 | 0.44 | |
| APG-EA Diluent Water | 1 | 10 | 100 | 0.41 | 0.47 |
| | 2 | 10 | 100 | 0.44 | |
| | 3 | 8 | 80 | 0.54 | |
| | 4 | 10 | 100 | 0.42 | |
| 10 | 1 | 10 | 100 | 0.49 | 0.46 |
| | 2 | 10 | 100 | 0.48 | |
| | 3 | 10 | 100 | 0.46 | |
| | 4 | 10 | 100 | 0.40 | |
| 18 | 1 | 10 | 100 | 0.52 | 0.50 |
| | 2 | 10 | 100 | 0.50 | |
| | 3 | 10 | 100 | 0.48 | |
| | 4 | 9 | 90 | 0.48 | |
| 32 | 1 | 9 | 90 | 0.48 | 0.42 |
| | 2 | 9 | 90 | 0.48 | |
| | 3 | 10 | 100 | 0.35 | |
| | 4 | 7 | 70 | 0.38 | |
| 56 | 1 | 5 | 50 | 0.32 | 0.22 |
| | 2 | 7 | 70 | 0.16 | |
| | 3 | 7 | 70 | 0.16 | |
| | 4 | 4 | 40 | 0.25 | |
| 100 | 1 | 0 | 0 | | |
| | 2 | 0 | 0 | | |
| | 3 | 0 | 0 | | |
| | 4 | 0 | 0 | | |

TABLE A33-4. FATHEAD MINNOW BEACH POINT RAW GROUNDWATER TOXICITY
TEST STATISTICAL ANALYSIS (TEST NO. 4) - LARVAL
SURVIVAL AFTER 7 DAYS OF EXPOSURE^a

Data Transformation:

Arc sine square root

Chi-Square Test for Normality:

| | |
|----------------------------|---|
| Calculated test statistic: | 12.39 |
| Alpha value: | 0.01 |
| Critical Value: | 13.28 |
| Conclusion: | Fail to reject the null hypothesis that the data are normally distributed |

Bartlett's Test for Homogeneity of Variances:

Data failed to meet homogeneity of variance test because at least one group had zero variance.

Steel's Many-One Rank Test:

| | |
|----------------------------|--|
| Calculated test statistic: | See Table A33-5 |
| Alpha value: | 0.05 |
| Critical Value: | 10.0 |
| Conclusion: | Reject the null hypothesis that all groups are equal |

^a The 100% raw Beach Point groundwater treatment was not included in the statistical analyses because all organisms died during the test.

TABLE A33-5. FATHEAD MINNOW BEACH POINT RAW GROUNDWATER TOXICITY TEST STATISTICAL ANALYSIS (TEST NO. 4) - RESULTS OF STEEL'S MANY-ONE RANK TEST ON LARVAL SURVIVAL AFTER 7 DAYS OF EXPOSURE

| Conc (% by Vol) | No. of Reps | Mean Survival (%) ^a | Rank Sum | Significance |
|----------------------------|-------------------|--------------------------------------|-------------|--------------|
| UMD/WREC Control | 4 | 97.5 | | |
| APG-EA Diluent Water | 4 | 95.0 | 17.5 | |
| 10 | 4 | 100.0 | 20.0 | |
| 18 | 4 | 97.5 | 18.0 | |
| 32 | 4 | 87.5 | 13.5 | |
| 56 | 4 | 57.5 | 10.0 | * |

^a Values given are actual percent survival means rather than arc sine square root transformed means which were used in the statistical analysis.

* Significantly different at alpha = 0.05 (Steel's critical value = 10.0).

TABLE A33-6. FATHEAD MINNOW BEACH POINT RAW GROUNDWATER TOXICITY
TEST STATISTICAL ANALYSIS (TEST NO. 4) - DRY WEIGHT
OF LARVAE AFTER 7 DAYS OF EXPOSURE^a

Data Transformation:

None

Chi-Square Test for Normality:

| | |
|----------------------------|---|
| Calculated test statistic: | 8.16 |
| Alpha value: | 0.01 |
| Critical value: | 13.28 |
| Conclusion: | Fail to reject the null hypothesis that the data are normally distributed |

Bartlett's Test for Homogeneity of Variances:

| | |
|----------------------------|--|
| Calculated test statistic: | 4.77 |
| Alpha value: | 0.01 |
| Critical value: | 15.09 |
| Conclusion: | Fail to reject the null hypothesis that the variances are homogenous |

ANOVA:

| | |
|----------------------------|--|
| Calculated test statistic: | 1.33 |
| Alpha value: | 0.05 |
| Critical value: | 3.06 |
| Conclusion: | Fail to reject the null hypothesis that all groups are equal |

^a The 56% buffered Beach Point groundwater treatment was not included in the statistical analyses because all concentrations above the NOEC for survival are excluded from the hypothesis test for growth effects.

APPENDIX 34

FATHEAD MINNOW ACUTE AND 7-DAY SURVIVAL AND GROWTH TEST CONDUCTED ON BEACH POINT BUFFERED (pH \approx 7) GROUNDWATER (WELL CC-33B) (TEST NO. 4)

Test Method: EPA/600/4-89/001
(Weber et al., 1989)

Type of Test: Static renewal (every 24 h)

Date: November 30 - December 7, 1993

Investigator: S.D. Turley

Laboratory: UMD/WREC

Groundwater:

Source: APG-EA Beach Point Well CC-33B

Chemical Characteristics: See Appendix 48

Dilution Water:

Source: 20% Perrier: 80% RO water

Chemical Characteristics: See Table A34-1

Test Organism:

Scientific Name: Pimephales promelas

Dry Weight: 0.49 mg (mean dry weight of
controls at end of test)

Age at Start of Test: <24 h

Source: UMD/WREC culture

Experimental Chambers:

Material: 600 mL glass beaker

Test Solution Volume: 400 mL

No. Organisms/Replicate: 10

No. Organisms/Treatment: 40

Loading: <0.5 g/L

Lighting: Fluorescent; 60-85 foot
candles

Aeration: Prior to renewals

pH Buffer:

10 N NaOH

Endpoints:

Mortality; growth

Water Quality:

Table A34-1

Results:

Mortality:

96-h Exposure:

The data for the 96-h LC50 were obtained from the 7-d study. The buffered groundwater was not toxic to the larvae. The data are summarized in Table A34-2.

7-d Exposure:

The survival of fathead minnow larvae was not affected after 7 days of exposure to Beach Point buffered groundwater (Tables A34-3 and A34-4).

Growth:

The growth of fathead minnow larvae was not affected after 7 days of exposure to Beach Point buffered groundwater (Tables A34-3 and A34-5).

TABLE A34-1. SUMMARY OF THE BEACH POINT BUFFERED GROUNDWATER
BIOASSAY TEST WATER QUALITY DATA FOR THE FATHEAD
MINNOW 7-DAY TEST (TEST NO. 4) - DISSOLVED OXYGEN
(MG/L)

| | <u>Test Concentrations (Percent Groundwater by Volume)</u> | | | | | |
|--------------|--|-----|-----|-----|-----|-----|
| | 0 | 10 | 18 | 32 | 56 | 100 |
| <u>Day 0</u> | | | | | | |
| 0 H | 8.4 | 8.1 | 8.2 | 8.0 | 7.7 | 7.6 |
| <u>Day 1</u> | | | | | | |
| 0 H | 8.5 | 8.2 | 8.0 | 8.1 | 8.1 | 8.0 |
| 24 H | 8.4 | 8.3 | 8.3 | 8.0 | 8.1 | 7.7 |
| <u>Day 2</u> | | | | | | |
| 0 H | 8.3 | 8.1 | 8.0 | 8.0 | 7.9 | 8.3 |
| 24 H | 8.2 | 8.4 | 8.2 | 7.1 | 7.9 | 7.5 |
| <u>Day 3</u> | | | | | | |
| 0 H | 8.4 | 8.0 | 8.0 | 8.1 | 7.9 | 7.5 |
| 24 H | 8.3 | 8.0 | 8.3 | 8.4 | 8.2 | 8.3 |
| <u>Day 4</u> | | | | | | |
| 0 H | 8.6 | 8.0 | 8.1 | 8.1 | 7.7 | 7.7 |
| 24 H | 8.0 | 7.9 | 7.9 | 8.1 | 8.1 | 8.1 |
| <u>Day 5</u> | | | | | | |
| 0 H | 8.7 | 8.1 | 8.2 | 8.1 | 8.3 | 8.3 |
| 24 H | 8.0 | 7.9 | 8.0 | 8.1 | 8.0 | 7.9 |
| <u>Day 6</u> | | | | | | |
| 0 H | 8.6 | 8.3 | 8.4 | 8.0 | 8.4 | 8.5 |
| 24 H | 8.1 | 7.8 | 8.0 | 8.0 | 7.8 | 7.6 |
| <u>Day 7</u> | | | | | | |
| 24 H | 8.0 | 7.9 | 7.8 | 7.8 | 7.7 | 7.7 |

TABLE A34-1. (CONTINUED) - pH (STANDARD UNITS)

| | | <u>Test Concentrations (Percent Groundwater by Volume)</u> | | | | | |
|--------------|--|--|------|------|------|------|------|
| | | 0 | 10 | 18 | 32 | 56 | 100 |
| <u>Day 0</u> | | | | | | | |
| 0 H | | 7.59 | 7.67 | 7.77 | 7.73 | 7.54 | 7.08 |
| <u>Day 1</u> | | | | | | | |
| 0 H | | 7.57 | 7.68 | 7.80 | 7.82 | 7.89 | 7.21 |
| 24 H | | 7.57 | 7.40 | 7.35 | 7.63 | 7.64 | 7.62 |
| <u>Day 2</u> | | | | | | | |
| 0 H | | 7.33 | 7.58 | 7.68 | 7.77 | 7.23 | 7.49 |
| 24 H | | 7.59 | 7.50 | 7.44 | 7.70 | 7.60 | 7.65 |
| <u>Day 3</u> | | | | | | | |
| 0 H | | 7.17 | 7.61 | 7.77 | 7.89 | 7.82 | 7.69 |
| 24 H | | 7.55 | 7.78 | 7.84 | 7.86 | 7.90 | 7.85 |
| <u>Day 4</u> | | | | | | | |
| 0 H | | 7.79 | 7.74 | 7.98 | 8.06 | 8.14 | 7.70 |
| 24 H | | 7.82 | 7.82 | 7.88 | 7.99 | 7.96 | 7.98 |
| <u>Day 5</u> | | | | | | | |
| 0 H | | 7.39 | 7.36 | 7.49 | 7.52 | 7.46 | 7.11 |
| 24 H | | 7.18 | 7.25 | 7.36 | 7.40 | 7.43 | 7.55 |
| <u>Day 6</u> | | | | | | | |
| 0 H | | 7.71 | 7.35 | 7.69 | 7.59 | 7.55 | 7.39 |
| 24 H | | 7.56 | 7.45 | 7.49 | 7.66 | 7.65 | 7.60 |
| <u>Day 7</u> | | | | | | | |
| 24 H | | 7.38 | 7.43 | 7.63 | 7.60 | 7.74 | 7.76 |

TABLE A34-1. (CONTINUED) - CONDUCTIVITY (μ MHOS/CM)

| | <u>Test Concentrations (Percent Groundwater by Volume)</u> | |
|--------------|--|------|
| | 0 | 100 |
| <u>Day 0</u> | | |
| 0 H | 240 | 1750 |
| <u>Day 1</u> | | |
| 0 H | 230 | 1650 |
| <u>Day 2</u> | | |
| 0 H | 240 | 1700 |
| <u>Day 3</u> | | |
| 0 H | 240 | 1700 |
| <u>Day 4</u> | | |
| 0 H | 250 | 1750 |
| <u>Day 5</u> | | |
| 0 H | 230 | 1700 |
| <u>Day 6</u> | | |
| 0 H | 210 | 1750 |
| <u>Day 7</u> | | |
| 24 H | 210 | 1700 |

TABLE A34-1. (CONTINUED) - ALKALINITY (MG/L AS CaCO₃)

| <u>Test Concentrations (Percent Groundwater by Volume)</u> | | |
|--|----------|------------|
| | <u>0</u> | <u>100</u> |
| <u>Day 0</u> | | |
| 0 H | 70 | 55 |
| <u>Day 1</u> | | |
| 0 H | 65 | 50 |
| <u>Day 2</u> | | |
| 0 H | 65 | 55 |
| <u>Day 3</u> | | |
| 0 H | 65 | 60 |
| <u>Day 4</u> | | |
| 0 H | 60 | 55 |
| <u>Day 5</u> | | |
| 0 H | 55 | 50 |
| <u>Day 6</u> | | |
| 0 H | 50 | 60 |
| <u>Day 7</u> | | |
| 24 H | 55 | 65 |

TABLE A34-1. (CONTINUED) - HARDNESS (MG/L AS CaCO₃)

| | <u>Test Concentrations (Percent Groundwater by Volume)</u> | |
|--------------|--|-----|
| | 0 | 100 |
| <u>Day 0</u> | | |
| 0 H | 80 | 210 |
| <u>Day 1</u> | | |
| 0 H | 74 | 200 |
| <u>Day 2</u> | | |
| 0 H | 70 | 200 |
| <u>Day 3</u> | | |
| 0 H | 78 | 210 |
| <u>Day 4</u> | | |
| 0 H | 74 | 220 |
| <u>Day 5</u> | | |
| 0 H | 70 | 220 |
| <u>Day 6</u> | | |
| 0 H | 80 | 198 |
| <u>Day 7</u> | | |
| 24 H | 70 | 194 |

TABLE A34-2. FATHEAD MINNOW BEACH POINT BUFFERED GROUNDWATER
TOXICITY TEST DATA (TEST NO. 4) - SURVIVAL AFTER 96
HOURS OF EXPOSURE

| Concentration (% Groundwater by Volume) | Rep | Number Tested | No. Alive at 96 Hours | Percent Alive |
|---|-----|------------------|--------------------------|------------------|
| UMD/WREC Control | A | 10 | 10 | 100 |
| | B | 10 | 10 | 100 |
| | C | 10 | 10 | 100 |
| | D | 10 | 10 | 100 |
| 10 | A | 10 | 10 | 100 |
| | B | 10 | 10 | 100 |
| | C | 10 | 10 | 100 |
| | D | 10 | 10 | 100 |
| 18 | A | 10 | 7 | 70 |
| | B | 10 | 10 | 100 |
| | C | 10 | 10 | 100 |
| | D | 10 | 10 | 100 |
| 32 | A | 10 | 10 | 100 |
| | B | 10 | 10 | 100 |
| | C | 10 | 10 | 100 |
| | D | 10 | 10 | 100 |
| 56 | A | 10 | 10 | 100 |
| | B | 10 | 10 | 100 |
| | C | 10 | 10 | 100 |
| | D | 10 | 10 | 100 |
| 100 | A | 10 | 10 | 100 |
| | B | 10 | 10 | 100 |
| | C | 10 | 10 | 100 |
| | D | 10 | 9 | 90 |

TABLE A34-3. FATHEAD MINNOW BEACH POINT BUFFERED GROUNDWATER
TOXICITY TEST DATA (TEST NO. 4) - LARVAL SURVIVAL
AND DRY WEIGHT AFTER 7 DAYS OF EXPOSURE

| Conc (% by Vol) | Rep | Number Larvae Alive | Percent Survival | Dry Weight (mg) | Mean Dry Weight (mg) |
|-----------------------|-----|---------------------------|---------------------|--------------------|----------------------------|
| UMD/WREC Control | 1 | 10 | 100 | 0.46 | 0.49 |
| | 2 | 10 | 100 | 0.57 | |
| | 3 | 9 | 90 | 0.50 | |
| | 4 | 10 | 100 | 0.44 | |
| 10 | 1 | 10 | 100 | 0.41 | 0.48 |
| | 2 | 9 | 90 | 0.49 | |
| | 3 | 10 | 100 | 0.45 | |
| | 4 | 10 | 100 | 0.57 | |
| 18 | 1 | 6 | 60 | 0.27 | 0.43 |
| | 2 | 9 | 90 | 0.47 | |
| | 3 | 8 | 80 | 0.57 | |
| | 4 | 10 | 100 | 0.41 | |
| 32 | 1 | 8 | 80 | 0.44 | 0.44 |
| | 2 | 10 | 100 | 0.40 | |
| | 3 | 9 | 90 | 0.48 | |
| | 4 | 10 | 100 | 0.44 | |
| 56 | 1 | 8 | 80 | 0.52 | 0.49 |
| | 2 | 9 | 90 | 0.45 | |
| | 3 | 8 | 80 | 0.48 | |
| | 4 | 8 | 80 | 0.51 | |
| 100 | 1 | 10 | 100 | 0.51 | 0.49 |
| | 2 | 8 | 80 | 0.50 | |
| | 3 | 8 | 80 | 0.49 | |
| | 4 | 8 | 80 | 0.46 | |

TABLE A34-4. FATHEAD MINNOW BEACH POINT BUFFERED GROUNDWATER
TOXICITY TEST STATISTICAL ANALYSIS (TEST NO. 4) -
LARVAL SURVIVAL AFTER 7 DAYS OF EXPOSURE

Data Transformation:

Arc sine square root

Chi-Square Test for Normality:

| | |
|----------------------------|---|
| Calculated test statistic: | 7.60 |
| Alpha value: | 0.01 |
| Critical Value: | 13.28 |
| Conclusion: | Fail to reject the null hypothesis that the data are normally distributed |

Bartlett's Test for Homogeneity of Variances:

| | |
|----------------------------|---|
| Calculated test statistic: | 5.52 |
| Alpha value: | 0.01 |
| Critical value: | 15.09 |
| Conclusion: | Fail to reject the null hypothesis that the data are homogenous |

ANOVA:

| | |
|-----------------------------|--|
| Calculated test statistics: | 2.36 |
| Alpha value: | 0.05 |
| Critical Value: | 2.77 |
| Conclusion: | Fail to reject the null hypothesis that all groups are equal |

TABLE A34-5. FATHEAD MINNOW BEACH POINT BUFFERED GROUNDWATER
TOXICITY TEST STATISTICAL ANALYSIS (TEST NO. 4) -
DRY WEIGHT OF LARVAE AFTER 7 DAYS OF EXPOSURE

Data Transformation:

None

Chi-Square Test for Normality:

| | |
|----------------------------|---|
| Calculated test statistic: | 3.78 |
| Alpha value: | 0.01 |
| Critical value: | 13.28 |
| Conclusion: | Fail to reject the null hypothesis that the data are normally distributed |

Bartlett's Test for Homogeneity of Variances:

| | |
|----------------------------|--|
| Calculated test statistic: | 10.78 |
| Alpha value: | 0.01 |
| Critical value: | 15.09 |
| Conclusion: | Fail to reject the null hypothesis that the variances are homogenous |

ANOVA:

| | |
|----------------------------|--|
| Calculated test statistic: | 0.75 |
| Alpha value: | 0.05 |
| Critical Value: | 2.77 |
| Conclusion: | Fail to reject the null hypothesis that all groups are equal |

APPENDIX 35

JAPANESE MEDAKA 96-H ACUTE TEST CONDUCTED ON BEACH POINT
RAW (pH \approx 4) GROUNDWATER (WELL CC-33B)

| | |
|---------------------------|-----------------------------------|
| Test Method: | EPA/600/4-90/027 (Weber, 1991) |
| Type of Test: | Static renewal (every 24 h) |
| Date: | November 30 - December 4, 1993 |
| Investigators: | R. S. Herriott and S. D. Turley |
| Laboratory: | UMD/WREC |
| Groundwater: | |
| Source: | APG-EA Beach Point Well CC-33B |
| Chemical Characteristics: | See Appendix 48 |
| Dilution Water: | |
| Source: | WREC non-chlorinated well water |
| Chemical Characteristics: | See Table A35-1 |
| Test Organism: | |
| Scientific Name: | <u>Oryzias latipes</u> |
| Age at Start of Test: | 22 days old |
| Source: | Ft. Detrick BRDL culture |
| Experimental Chambers: | |
| Material: | Glass beakers |
| Size: | 600 mL |
| Volume: | 400 mL |
| No. Organisms/Replicate: | 10 |
| No. Organisms/Treatment: | 20 |
| Loading: | <0.5 g/L |
| Lighting: | Fluorescent; 60-85 foot candles |
| Aeration: | None |

Endpoint:

Mortality

Water Quality:

Table A35-1

Results:

100% Beach Point raw groundwater killed 35% of the Japanese medaka fry in 96 h (Table A35-2); thus, a 96-h LC50 could not be calculated.

TABLE A35-1. SUMMARY OF THE BEACH POINT RAW GROUNDWATER BIOASSAY
WATER QUALITY DATA FOR THE JAPANESE MEDAKA 96-H
TEST - DISSOLVED OXYGEN (MG/L)

| | | <u>Test Concentrations (Percent Groundwater by Volume)</u> | | | | | |
|--------------|--|--|-----|-----|-----|-----|-----|
| | | 0 | 10 | 18 | 32 | 56 | 100 |
| <u>Day 0</u> | | | | | | | |
| 0 H | | 8.5 | 8.2 | 8.2 | 7.9 | 8.0 | 8.2 |
| <u>Day 1</u> | | | | | | | |
| 0 H | | 8.0 | 7.9 | 8.0 | 8.0 | 7.7 | 7.5 |
| 24 H | | 6.7 | 6.7 | 7.3 | 7.0 | 7.1 | 7.5 |
| <u>Day 2</u> | | | | | | | |
| 0 H | | 8.3 | 8.0 | 8.2 | 8.0 | 8.0 | 8.2 |
| 24 H | | 8.0 | 8.0 | 7.9 | 7.8 | 7.9 | 8.1 |
| <u>Day 3</u> | | | | | | | |
| 0 H | | 8.4 | 8.3 | 8.1 | 7.9 | 7.8 | 8.0 |
| 24 H | | 8.0 | 8.1 | 8.1 | 7.8 | 8.0 | 8.1 |
| <u>Day 4</u> | | | | | | | |
| 24 H | | 8.1 | 8.1 | 7.9 | 7.7 | 8.0 | 7.7 |

TABLE A35-1. (CONTINUED) - pH (STANDARD UNITS)

| | <u>Test Concentrations (Percent Groundwater by Volume)</u> | | | | | |
|--------------|--|------|------|------|------|------|
| | 0 | 10 | 18 | 32 | 56 | 100 |
| <u>Day 0</u> | | | | | | |
| 0 H | 7.70 | 7.62 | 7.44 | 7.25 | 6.73 | 4.01 |
| <u>Day 1</u> | | | | | | |
| 0 H | 7.70 | 7.48 | 7.07 | 6.94 | 6.48 | 3.97 |
| 24 H | 7.53 | 7.56 | 7.56 | 7.50 | 7.40 | 5.40 |
| <u>Day 2</u> | | | | | | |
| 0 H | 7.66 | 7.50 | 7.00 | 6.93 | 6.61 | 3.97 |
| 24 H | 7.62 | 7.66 | 7.75 | 7.63 | 7.51 | 5.47 |
| <u>Day 3</u> | | | | | | |
| 0 H | 7.79 | 7.59 | 7.28 | 6.94 | 6.58 | 3.91 |
| 24 H | 7.83 | 7.87 | 7.96 | 7.76 | 7.58 | 5.50 |
| <u>Day 4</u> | | | | | | |
| 24 H | 8.13 | 8.02 | 7.96 | 7.85 | 7.76 | 5.41 |

TABLE A35-1. (CONTINUED) - CONDUCTIVITY (μ MHOS/CM)

| <u>Test Concentrations (Percent Groundwater by Volume)</u> | | |
|--|-----|------|
| | 0 | 100 |
| <u>Day 0</u> | | |
| 0 H | 280 | 1750 |
| <u>Day 1</u> | | |
| 0 H | 270 | 1700 |
| <u>Day 2</u> | | |
| 0 H | 270 | 1800 |
| <u>Day 3</u> | | |
| 0 H | 270 | 1700 |

TABLE A35-1. (CONTINUED) - ALKALINITY (MG/L as CaCO₃)

| | <u>Test Concentrations (Percent Groundwater by Volume)</u> | |
|--------------|--|-----|
| | 0 | 100 |
| <u>Day 0</u> | | |
| 0 H | 75 | a |
| <u>Day 1</u> | | |
| 0 H | 70 | a |
| <u>Day 2</u> | | |
| 0 H | 75 | a |
| <u>Day 3</u> | | |
| 0 H | 70 | a |

^a Could not obtain measurement.

TABLE A35-1. (CONTINUED) - HARDNESS (MG/L AS CaCO_3)

| <u>Test Concentrations (Percent Groundwater by Volume)</u> | | |
|--|----|-----|
| | 0 | 100 |
| <u>Day 0</u> | | |
| 0 H | 60 | a |
| <u>Day 1</u> | | |
| 0 H | 55 | a |
| <u>Day 2</u> | | |
| 0 H | 60 | a |
| <u>Day 3</u> | | |
| 0 H | 64 | a |

^a Could not obtain measurement.

TABLE A35-2. JAPANESE MEDAKA BEACH POINT RAW GROUNDWATER 96-H
TOXICITY TEST - PERCENT FRY SURVIVAL AFTER 96
HOURS OF EXPOSURE

| Concentration (% by Volume) | Rep | Number Tested | No. Alive at End of Test | Percent Alive |
|--------------------------------|-----|------------------|-----------------------------|------------------|
| UM/WREC Control | 1 | 10 | 10 | 100 |
| | 2 | 10 | 10 | 100 |
| APG-EA Diluent Water | 1 | 10 | 10 | 100 |
| | 2 | 10 | 10 | 100 |
| 10 | 1 | 10 | 10 | 100 |
| | 2 | 10 | 10 | 100 |
| 18 | 1 | 10 | 10 | 100 |
| | 2 | 10 | 10 | 100 |
| 32 | 1 | 10 | 10 | 100 |
| | 2 | 10 | 9 | 90 |
| 56 | 1 | 10 | 10 | 100 |
| | 2 | 10 | 10 | 100 |
| 100 | 1 | 10 | 6 | 60 |
| | 2 | 10 | 7 | 70 |

APPENDIX 36

LIST OF DATA REPORTS FOR THE AMES ASSAYS CONDUCTED
ON THE BEACH POINT HISTOPATHOLOGY EXPOSURE
TANKS AND APG-EA DILUENT WATER

Table A36-1. AMES REPORTS FOR TEST NO. 1 (SAMPLES TAKEN AT BEACH POINT ON APRIL 14, 1993)

- Lawlor, T.E. 1993. Mutagenicity test on water sample APG-EA Beach Pt. 10% groundwater in the Salmonella/mammalian-microsome reverse mutation assay (Ames test) modified for direct water samples. Final Report. HWA Study No. 15617-0-401W. Hazleton Washington, Inc., Vienna, VA.
- Lawlor, T.E. 1993. Mutagenicity test on an extract of the water sample APG-EA Beach Pt. 10% groundwater in the Salmonella/mammalian-microsome reverse mutation assay (Ames test) modified for XAD-2 resin extracts. Final Report. HWA Study No. 15617-1-401X. Hazleton Washington, Inc., Vienna, VA.
- Lawlor, T.E. 1993. Mutagenicity test on water sample APG-EA Beach Pt. 1% groundwater in the Salmonella/mammalian-microsome reverse mutation assay (Ames test) modified for direct water samples. Final Report. HWA Study No. 15616-0-401W. Hazleton Washington, Inc., Vienna, VA.
- Lawlor, T.E. 1993. Mutagenicity test on an extract of the water sample APG-EA Beach Pt. 1% groundwater in the Salmonella/mammalian-microsome reverse mutation assay (Ames test) modified for XAD-2 resin extracts. Final Report. HWA Study No. 15616-1-401X. Hazleton Washington, Inc., Vienna, VA.
- Lawlor, T.E. 1993. Mutagenicity test on water sample APG-EA diluent water in the Salmonella/mammalian-microsome reverse mutation assay (Ames test) modified for direct water samples. Final Report. HWA Study No. 15615-0-401W. Hazleton Washington, Inc., Vienna, VA.
- Lawlor, T.E. 1993. Mutagenicity test on water sample APG-EA diluent water in the Salmonella/mammalian-microsome reverse mutation assay (Ames test) modified for XAD-2 resin extracts. Final Report. HWA Study No. 15615-1-401W. Hazleton Washington, Inc., Vienna, VA.
-

Table A36-2. AMES REPORTS FOR TEST NO. 2 (SAMPLES TAKEN AT BEACH POINT ON JUNE 2, 1993)

Lawlor, T.E. 1993. Mutagenicity test on water sample APG-EA Beach Pt. 10% groundwater in the Salmonella/mammalian-microsome reverse mutation assay (Ames test) modified for direct water samples. Final Report. HWA Study No. 15676-0-401W. Hazleton Washington, Inc., Vienna, VA.

Lawlor, T.E. 1993. Mutagenicity test on an extract of the water sample APG-EA Beach Pt. 10% groundwater in the Salmonella/mammalian-microsome reverse mutation assay (Ames test) modified for XAD-2 resin extracts. Final Report. HWA Study No. 15676-1-401X. Hazleton Washington, Inc., Vienna, VA.

Lawlor, T.E. 1993. Mutagenicity test on water sample APG-EA Beach Pt. 1% groundwater in the Salmonella/mammalian-microsome reverse mutation assay (Ames test) modified for direct water samples. Final Report. HWA Study No. 15677-0-401W. Hazleton Washington, Inc., Vienna, VA.

Lawlor, T.E. 1993. Mutagenicity test on an extract of the water sample APG-EA Beach Pt. 1% groundwater in the Salmonella/mammalian-microsome reverse mutation assay (Ames test) modified for XAD-2 resin extracts. Final Report. HWA Study No. 15677-1-401X. Hazleton Washington, Inc., Vienna, VA.

Lawlor, T.E. 1993. Mutagenicity test on water sample APG-EA diluent/control water in the Salmonella/mammalian-microsome reverse mutation assay (Ames test) modified for direct water samples. Final Report. HWA Study No. 15678-0-401W. Hazleton Washington, Inc., Vienna, VA.

Lawlor, T.E. 1993. Mutagenicity test on an extract of the water sample APG-EA diluent/control water in the Salmonella/mammalian-microsome reverse mutation assay (Ames test) modified for XAD-2 resin extracts. Final Report. HWA Study No. 15678-1-401W. Hazleton Washington, Inc., Vienna, VA.

Table A36-3. AMES REPORTS FOR TEST NO. 3 (SAMPLES TAKEN AT BEACH POINT ON AUGUST 23, 1993)

- Lawlor, T.E. 1993. Mutagenicity test on water sample 100% groundwater APG-EA Beach Pt. Well 33B in the Salmonella/mammalian-microsome reverse mutation assay (Ames test) modified for direct water samples. Final Report. HWA Study No. 15817-0-401W. Hazleton Washington, Inc., Vienna, VA.
- Lawlor, T.E. 1993. Mutagenicity test on an extract of the water sample 100% groundwater APG-EA Beach Pt. Well 33B in the Salmonella/mammalian-microsome reverse mutation assay (Ames test) modified for XAD-2 resin extracts. Final Report. HWA Study No. 15817-1-401X. Hazleton Washington, Inc., Vienna, VA.
- Lawlor, T.E. 1993. Mutagenicity test on water sample 10% groundwater APG-EA Beach Pt. Well 33B in the Salmonella/mammalian-microsome reverse mutation assay (Ames test) modified for direct water samples. Final Report. HWA Study No. 15816-0-401W. Hazleton Washington, Inc., Vienna, VA.
- Lawlor, T.E. 1993. Mutagenicity test on an extract of the water sample 10% groundwater APG-EA Beach Pt. Well 33B in the Salmonella/mammalian-microsome reverse mutation assay (Ames test) modified for XAD-2 resin extracts. Final Report. HWA Study No. 15816-1-401X. Hazleton Washington, Inc., Vienna, VA.
- Lawlor, T.E. 1993. Mutagenicity test on water sample APG-EA diluent/control water in the Salmonella/mammalian-microsome reverse mutation assay (Ames test) modified for direct water samples. Final Report. HWA Study No. 15815-0-401W. Hazleton Washington, Inc., Vienna, VA.
- Lawlor, T.E. 1993. Mutagenicity test on an extract of the water sample APG-EA diluent/control water in the Salmonella/mammalian-microsome reverse mutation assay (Ames test) modified for XAD-2 resin extracts. Final Report. HWA Study No. 15815-1-401W. Hazleton Washington, Inc., Vienna, VA.
-

Table A36-4. AMES REPORTS FOR TEST NO. 4 (SAMPLES TAKEN AT BEACH POINT ON NOVEMBER 30, 1993)

Lawlor, T.E. 1994. Mutagenicity test on water sample 100% groundwater APG-EA Beach Pt. Well 33B in the Salmonella/mammalian-microsome reverse mutation assay (Ames test) modified for direct water samples. Final Report. HWA Study No. 15976-0-401W. Hazleton Washington, Inc., Vienna, VA.

Lawlor, T.E. 1994. Mutagenicity test on an extract of the water sample 100% groundwater APG-EA Beach Pt. Well 33B in the Salmonella/mammalian-microsome reverse mutation assay (Ames test) modified for XAD-2 resin extracts. Final Report. HWA Study No. 15976-1-401X. Hazleton Washington, Inc., Vienna, VA.

Lawlor, T.E. 1994. Mutagenicity test on water sample 10% groundwater APG-EA Beach Pt. Well 33B in the Salmonella/mammalian-microsome reverse mutation assay (Ames test) modified for direct water samples. Final Report. HWA Study No. 15975-0-401W. Hazleton Washington, Inc., Vienna, VA.

Lawlor, T.E. 1994. Mutagenicity test on an extract of the water sample 10% groundwater APG-EA Beach Pt. Well 33B in the Salmonella/mammalian-microsome reverse mutation assay (Ames test) modified for XAD-2 resin extracts. Final Report. HWA Study No. 15975-1-401X. Hazleton Washington, Inc., Vienna, VA.

Lawlor, T.E. 1994. Mutagenicity test on water sample APG-EA diluent/control water in the Salmonella/mammalian-microsome reverse mutation assay (Ames test) modified for direct water samples. Final Report. HWA Study No. 15974-0-401W. Hazleton Washington, Inc., Vienna, VA.

Lawlor, T.E. 1994. Mutagenicity test on an extract of the water sample APG-EA diluent/control water in the Salmonella/mammalian-microsome reverse mutation assay (Ames test) modified for XAD-2 resin extracts. Final Report. HWA Study No. 15974-1-401W. Hazleton Washington, Inc., Vienna, VA.

APPENDIX 37

LIST OF DATA REPORTS FOR THE SCE ASSAYS CONDUCTED ON THE BEACH POINT HISTOPATHOLOGY EXPOSURE TANKS AND APG-EA DILUENT WATER

Sample taken June 2, 1993

Murli, H. 1993. Mutagenicity test on an extract of the water sample APG-EA Beach Pt. 10% GW in an in vitro cytogenetic assay measuring sister chromatid exchange in Chinese hamster ovary (CHO) cells. Final Report. HWA Study No. 15676-0-438. Hazleton Washington, Inc., Vienna, VA.

Sample taken July 19, 1993

Murli, H. 1993. Mutagenicity test on an extract of the water sample 100% GW (Well 33B) APG-EA Beach Pt. in an in vitro cytogenetic assay measuring sister chromatid exchange in Chinese hamster ovary (CHO) cells. Final Report. HWA Study No. 15748-0-438. Hazleton Washington, Inc., Vienna, VA.

Sample taken August 23, 1993

Murli, H. 1993. Mutagenicity test on an extract of the water sample 0% GW (dilution/control water) in an in vitro cytogenetic assay measuring sister chromatid exchange in Chinese hamster ovary (CHO) cells. Final Report. HWA Study No. 15818-0-438. Hazleton Washington, Inc., Vienna, VA.

Sample taken November 30, 1993

Murli, H. 1994. Mutagenicity test on 100% groundwater APG-EA, Beach Pt. Well 33B in an in vitro cytogenetic assay measuring sister chromatid exchange in Chinese hamster ovary (CHO) cells. Final Report. HWA Study No. 15976-0-438. Hazleton Washington, Inc., Vienna, VA.

APPENDIX 38

FROG EMBRYO TERATOGENESIS ASSAY - Xenopus (FETAX)
CONDUCTED ON BEACH POINT RAW (pH \approx 4)
GROUNDWATER (WELL 33B)
(TEST NO. 1)

| | |
|---------------------------|---|
| Test Method: | ASTM Designation E 1439-91 ASTM (1991) |
| Type of Test: | Static renewal (every 24 h) |
| Date: | April 10-14, 1993 |
| Investigator: | S. D. Turley |
| Laboratory: | UMD/WREC |
| Groundwater: | |
| Source: | APG-EA Beach Point Well 33B |
| Chemical Characteristics: | See Appendix 48 |
| Test Medium: | |
| Source: | FETAX solution |
| pH characteristics: | See Table A38-1 |
| Test Organism: | |
| Scientific Name: | <u>Xenopus laevis</u> |
| Age at Start of Test: | Stage 8 blastula to stage 11 gastrulae |
| Source: | UMD/WREC culture |
| Experimental Chambers: | |
| Material: | Glass petri dishes |
| Test Solution Volume: | 10 mL |
| No. Organisms/Replicate: | 25 |
| No. Organisms/Treatment: | Control: 100 Groundwater: 50 |
| Loading: | n/a |
| Lighting: | Fluorescent; 60-85 foot candles |
| Aeration: | Prior to renewals |

| | |
|-------------------|-------------------------|
| Endpoints: | Mortality; malformation |
| Test Temperature: | 24 ± 0.2°C |

Results:

Mortality:

The raw groundwater did not affect embryo survival. The data are summarized in Tables A38-2 and A38-3.

Malformations:

Significant ($\alpha = 0.05$) embryo malformations occurred at all concentrations down to 18% raw groundwater by volume (see Tables A38-2, A38-4, and A38-5). Less than 50% malformation occurred at all test concentrations, thus, an EC50 could not be calculated (Table A38-2). The NOEC and LOEC for the embryos, based on increased numbers of malformations, are as follows:

NOEC = 10% raw groundwater by volume.
LOEC = 18% raw groundwater by volume.

The types of malformed embryos are given in Table A38-6.

TABLE A38-1. SUMMARY OF THE BEACH POINT RAW GROUNDWATER
BIOASSAY pH (STANDARD UNITS) DATA FOR FETAX (TEST
NO. 1)

| | <u>Test Concentrations (Percent Groundwater by Volume)</u> | | | | | |
|--------------|--|------|------|------|------|------|
| | 0 | 10 | 18 | 32 | 56 | 100 |
| <u>Day 0</u> | | | | | | |
| 0 H | 7.51 | 7.31 | 7.11 | 6.71 | 6.12 | 4.10 |
| <u>Day 1</u> | | | | | | |
| 0 H | 7.48 | 7.24 | 7.01 | 6.54 | 6.00 | 4.19 |
| <u>Day 2</u> | | | | | | |
| 0 H | 7.56 | 7.36 | 7.20 | 6.77 | 6.03 | 4.02 |
| <u>Day 3</u> | | | | | | |
| 0 H | 7.59 | 7.39 | 7.18 | 6.59 | 5.96 | 3.98 |
| <u>Day 4</u> | | | | | | |
| 24 H | 7.54 | 7.33 | 7.13 | 6.65 | 6.03 | 4.07 |

TABLE A38-2. FETAX BEACH POINT RAW GROUNDWATER TOXICITY TEST DATA (TEST NO. 1) - PERCENT EMBRYO SURVIVAL AND MALFORMATIONS AFTER 96 HOURS OF EXPOSURE^a

| Conc (% by Vol) | Rep | Number Embryos Alive | Percent Survival | Number Embryos Malformed | Percent Malformed |
|-----------------------|-----|----------------------------|---------------------|--------------------------------|----------------------|
| UMD/WREC Control | 1 | 25 | 100 | 1 | 4.0 |
| | 2 | 25 | 100 | 1 | 4.0 |
| | 3 | 24 | 96 | 0 | 0.0 |
| | 4 | 25 | 100 | 1 | 4.0 |
| 10 | 1 | 24 | 96 | 2 | 8.3 |
| | 2 | 23 | 92 | 3 | 13.0 |
| 18 | 1 | 24 | 96 | 4 | 16.7 |
| | 2 | 25 | 100 | 4 | 16.0 |
| 32 | 1 | 23 | 92 | 3 | 13.0 |
| | 2 | 24 | 96 | 4 | 16.7 |
| 56 | 1 | 24 | 96 | 4 | 16.7 |
| | 2 | 22 | 88 | 4 | 18.2 |
| 100 | 1 | 24 | 96 | 8 | 33.3 |
| | 2 | 23 | 92 | 13 | 56.5 |

^a An APG-EA diluent water control was not run during FETAX.

TABLE A38-3. FETAX BEACH POINT RAW GROUNDWATER TOXICITY TEST
STATISTICAL ANALYSIS (TEST NO. 1) - PERCENT EMBRYO
SURVIVAL AFTER 96 HOURS OF EXPOSURE

Data Transformation:

Arc sine square root

Chi-Square Test for Normality:

| | |
|----------------------------|---|
| Calculated test statistic: | 5.69 |
| Alpha value: | 0.01 |
| Critical value: | 13.28 |
| Conclusion: | Fail to reject the null hypothesis that the data are normally distributed |

Bartlett's Test for Homogeneity of Variances:

| | |
|----------------------------|--|
| Calculated test statistic: | 0.82 |
| Alpha value: | 0.01 |
| Critical value: | 15.09 |
| Conclusion: | Fail to reject the null hypothesis that the variances are homogenous |

ANOVA:

| | |
|----------------------------|--|
| Calculated test statistic: | 2.50 |
| Alpha value: | 0.05 |
| Critical value: | 3.69 |
| Conclusion: | Fail to reject the null hypothesis that all groups are equal |

TABLE A38-4. FETAX BEACH POINT RAW GROUNDWATER TOXICITY TEST
STATISTICAL ANALYSIS (TEST NO. 1) - PERCENT EMBRYO
MALFORMATIONS AFTER 96 HOURS OF EXPOSURE

Data Transformation:

Arc sine square root

Chi-Square Test for Normality:

| | |
|----------------------------|---|
| Calculated test statistic: | 5.69 |
| Alpha value: | 0.01 |
| Critical Value: | 13.28 |
| Conclusion: | Fail to reject the null hypothesis that the data are normally distributed |

Bartlett's Test for Homogeneity of Variances:

| | |
|----------------------------|--|
| Calculated test statistic: | 9.95 |
| Alpha value: | 0.01 |
| Critical value: | 15.09 |
| Conclusion: | Fail to reject the null hypothesis that the variances are homogenous |

ANOVA:

| | |
|----------------------------|--|
| Calculated test statistic: | 17.20 |
| Alpha value: | 0.05 |
| Critical value: | 3.69 |
| Conclusion: | Reject the null hypothesis that all groups are equal |

Bonferroni's T-Test:

| | |
|----------------------------|--|
| Calculated test statistic: | See Table A38-5 |
| Alpha value: | 0.05 |
| Critical value: | 2.90 |
| Conclusion: | Reject the null hypothesis that all groups are equal |

TABLE A38-5. FETAX BEACH POINT RAW GROUNDWATER TOXICITY TEST
STATISTICAL ANALYSIS (TEST NO. 1) - RESULTS OF
BONFERRONI'S T-TEST ON EMBRYO MALFORMATIONS AFTER
96 HOURS OF EXPOSURE

| Conc (% by Vol) | No. of Reps | Mean Normal Embryos (%) ^a | T Statistic | Significance |
|-----------------------|-------------------|---|-------------|--------------|
| UMD/WREC Control | 4 | 97.0 | | |
| 10 | 2 | 89.4 | 2.52 | |
| 18 | 2 | 83.7 | 3.92 | * |
| 32 | 2 | 85.1 | 3.57 | * |
| 56 | 2 | 82.5 | 4.16 | * |
| 100 | 2 | 55.1 | 9.09 | * |

^a Values given are actual percent mean normal embryos rather than arc sine square root transformed means which were used in the statistical analysis.

* Significantly different at alpha = 0.05 (Bonferroni's critical value = 2.90).

TABLE A38-6. FETAX BEACH POINT RAW GROUNDWATER TOXICITY TEST
DATA (TEST NO. 1) - TYPE AND NUMBER OF MALFORMED
EMBRYOS AFTER 96 HOURS OF EXPOSURE

| Malformation | Test Concentrations (% Groundwater by Volume) | | | | | | | | | |
|--------------|---|---|---|---|------------|---|------------|---|------------|---|
| | 0 | | | | 10 | | 18 | | 32 | |
| | <u>Rep</u> | | | | <u>Rep</u> | | <u>Rep</u> | | <u>Rep</u> | |
| | 1 | 2 | 3 | 4 | 1 | 2 | 1 | 2 | 1 | 2 |
| Severe | | | | | | | | | | |
| Gut, coiling | | | | 1 | 1 | | 2 | 2 | 3 | 1 |
| Edema: | | | | | | | | | | |
| Multiple | | | | | 2 | | 2 | 1 | 1 | 1 |
| Cardiac | | | | | | | | | | |
| Abdominal | | | | 1 | 1 | 1 | 1 | | 2 | 1 |
| Facial | | | | | | | | | | |
| Cephalic | | | | | | | | | | |
| Blisters | | | | | | | | | | |
| Tail | | | | | | | | | | |
| Notochord | | | | 1 | | | | | | 1 |
| Fin | | | | | | | | | | |
| Face | | | | | | | | | | 3 |
| Eye | | | | | | | | | | |
| Brain | | | | | | | | | | |
| Hemorrhage | | | | | | | | | | |
| Cardiac | | | | | | | | | | |
| Other | | | | | | | | | | |

APPENDIX 39

FROG EMBRYO TERATOGENESIS ASSAY - Xenopus (FETAX)
CONDUCTED ON BEACH POINT BUFFERED (pH \approx 7)
GROUNDWATER (WELL 33B)
(TEST NO. 1)

| | |
|---------------------------|---|
| Test Method: | ASTM Designation E 1439-91 ASTM (1991) |
| Type of Test: | Static renewal (every 24 h) |
| Date: | April 10-14, 1993 |
| Investigator: | S. D. Turley |
| Laboratory: | UMD/WREC |
| Groundwater: | |
| Source: | APG-EA Beach Point Well 33B |
| Chemical Characteristics: | See Appendix 48 |
| Test Medium: | |
| Source: | FETAX solution |
| pH characteristics: | See Table A39-1 |
| Test Organism: | |
| Scientific Name: | <u>Xenopus laevis</u> |
| Age at Start of Test: | Stage 8 blastula to stage 11 gastrulae |
| Source: | UMD/WREC culture |
| Experimental Chambers: | |
| Material: | Glass petri dishes |
| Test Solution Volume: | 10 mL |
| No. Organisms/Replicate: | 25 |
| No. Organisms/Treatment: | Groundwater: 50 Control: 100 |
| Loading: | n/a |
| Lighting: | Fluorescent; 60-85 foot candles |
| Aeration: | Prior to renewals |

| | |
|-------------------|--------------------------|
| pH Buffer: | 10 N NaOH |
| Endpoints: | Mortality; malformations |
| Test Temperature: | 24 ± 0.2°C |

Results:

Mortality:

The buffered groundwater did not affect embryo survival.
The data are summarized in Tables A39-2 and A39-3.

Malformations:

Significant ($\alpha = 0.05$) embryo malformations occurred at all concentrations down to 32% buffered groundwater by volume (see Tables A39-2, A39-4, and A39-5). Less than 50% malformation occurred at all test concentrations, thus, an EC50 could not be calculated (Table A39-2). The NOEC and LOEC for the embryos, based on increased numbers of malformations, are as follows:

NOEC = 18% buffered groundwater by volume.

LOEC = 32% buffered groundwater by volume.

The types of malformed embryos are given in Table A39-6.

TABLE A39-1. SUMMARY OF THE BEACH POINT BUFFERED GROUNDWATER
BIOASSAY pH (STANDARD UNITS) DATA FOR FETAX (TEST
NO. 1)

| <u>Test Concentrations (Percent Groundwater by Volume)</u> | | | | | |
|--|------|------|------|------|------|
| | 0 | 18 | 32 | 56 | 100 |
| <u>Day 0</u> | | | | | |
| 0 H | 7.51 | 7.13 | 6.99 | 7.11 | 6.92 |
| <u>Day 1</u> | | | | | |
| 0 H | 7.48 | 7.19 | 7.12 | 7.16 | 7.06 |
| <u>Day 2</u> | | | | | |
| 0 H | 7.56 | 7.18 | 7.11 | 7.13 | 7.00 |
| <u>Day 3</u> | | | | | |
| 0 H | 7.59 | 7.12 | 7.02 | 7.08 | 6.95 |
| <u>Day 4</u> | | | | | |
| 24 H | 7.54 | 7.17 | 7.06 | 7.12 | 6.98 |

TABLE A39-2. FETAX BEACH POINT BUFFERED GROUNDWATER TOXICITY
TEST DATA (TEST NO. 1) - PERCENT EMBRYO SURVIVAL
AND MALFORMATIONS AFTER 96 HOURS OF EXPOSURE

| Conc (% by Vol) | Rep | Number Embryos Alive | Percent Survival | Number Embryos Malformed | Percent Malformed |
|-----------------------|-----|----------------------------|---------------------|--------------------------------|----------------------|
| UMD/WREC Control | 1 | 25 | 100 | 1 | 4.0 |
| | 2 | 25 | 100 | 1 | 4.0 |
| | 3 | 24 | 96 | 0 | 0.0 |
| | 4 | 25 | 100 | 1 | 4.0 |
| 18 | 1 | 23 | 92 | 2 | 8.7 |
| | 2 | 25 | 100 | 2 | 8.0 |
| 32 | 1 | 23 | 92 | 2 | 8.7 |
| | 2 | 24 | 96 | 3 | 13.0 |
| 56 | 1 | 23 | 92 | 5 | 21.7 |
| | 2 | 24 | 96 | 5 | 20.8 |
| 100 | 1 | 24 | 96 | 4 | 16.7 |
| | 2 | 25 | 100 | 7 | 28.0 |

TABLE A39-3. FETAX BEACH POINT BUFFERED GROUNDWATER TOXICITY
TEST STATISTICAL ANALYSIS (TEST NO. 1) - PERCENT
EMBRYO SURVIVAL AFTER 96 HOURS OF EXPOSURE

Data Transformation:

Arc sine square root

Chi-Square Test for Normality:

| | |
|----------------------------|---|
| Calculated test statistic: | 4.08 |
| Alpha value: | 0.01 |
| Critical value: | 13.28 |
| Conclusion: | Fail to reject the null hypothesis that the data are normally distributed |

Bartlett's Test for Homogeneity of Variances:

| | |
|----------------------------|--|
| Calculated test statistic: | 1.44 |
| Alpha value: | 0.01 |
| Critical value: | 13.28 |
| Conclusion: | Fail to reject the null hypothesis that the variances are homogenous |

ANOVA:

| | |
|----------------------------|--|
| Calculated test statistic: | 1.60 |
| Alpha value: | 0.05 |
| Critical value: | 4.12 |
| Conclusion: | Fail to reject the null hypothesis that all groups are equal |

TABLE A39-4. FETAX BEACH POINT BUFFERED GROUNDWATER TOXICITY
TEST STATISTICAL ANALYSIS (TEST NO. 1) - PERCENT
EMBRYO MALFORMATIONS AFTER 96 HOURS OF EXPOSURE

Data Transformation:

Arc sine square root

Chi-Square Test for Normality:

| | |
|----------------------------|---|
| Calculated test statistic: | 4.08 |
| Alpha value: | 0.01 |
| Critical Value: | 13.28 |
| Conclusion: | Fail to reject the null hypothesis that the data are normally distributed |

Bartlett's Test for Homogeneity of Variances:

| | |
|----------------------------|--|
| Calculated test statistic: | 7.56 |
| Alpha value: | 0.01 |
| Critical value: | 13.28 |
| Conclusion: | Fail to reject the null hypothesis that the variances are homogenous |

ANOVA:

| | |
|----------------------------|--|
| Calculated test statistic: | 16.33 |
| Alpha value: | 0.05 |
| Critical value: | 4.12 |
| Conclusion: | Reject the null hypothesis that all groups are equal |

Bonferroni's T-Test:

| | |
|----------------------------|--|
| Calculated test statistic: | See Table A39-5 |
| Alpha value: | 0.05 |
| Critical value: | 2.84 |
| Conclusion: | Reject the null hypothesis that all groups are equal |

TABLE A39-5. FETAX BEACH POINT BUFFERED GROUNDWATER TOXICITY TEST STATISTICAL ANALYSIS (TEST NO. 1) - RESULTS OF BONFERRONI'S T-TEST ON EMBRYO MALFORMATIONS AFTER 96 HOURS OF EXPOSURE

| Conc (% by Vol) | No. of Reps | Mean Normal Embryos (%) ^a | T Statistic | Significance |
|-----------------------|-------------------|---|-------------|--------------|
| UMD/WREC Control | 4 | 97.0 | | |
| 18 | 2 | 91.7 | 2.47 | |
| 32 | 2 | 89.1 | 3.33 | * |
| 56 | 2 | 78.7 | 6.41 | * |
| 100 | 2 | 77.5 | 6.65 | * |

^a Values given are actual percent mean normal embryos rather than arc sine square root transformed means which were used in the statistical analysis.

* Significantly different at alpha = 0.05 (Bonferroni's critical value = 2.84).

TABLE A39-6. FETAX BEACH POINT BUFFERED GROUNDWATER TOXICITY
TEST DATA (TEST NO. 1) - TYPE AND NUMBER OF
MALFORMED EMBRYOS AFTER 96 HOURS OF EXPOSURE

| Malformation | Test Concentrations (% Groundwater by Volume) | | | | | | | | | | | |
|--------------|---|---|---|---|------------|---|------------|---|------------|---|------------|---|
| | 0 | | | | 18 | | 32 | | 56 | | 100 | |
| | <u>Rep</u> | | | | <u>Rep</u> | | <u>Rep</u> | | <u>Rep</u> | | <u>Rep</u> | |
| | 1 | 2 | 3 | 4 | 1 | 2 | 1 | 2 | 1 | 2 | 1 | 2 |
| <hr/> | | | | | | | | | | | | |
| Severe | | | | | | | | | | | | |
| Gut, coiling | | | | 1 | 1 | | 1 | | 2 | | 1 | 1 |
| Edema: | | | | | | | | | | | | |
| Multiple | | | | | 1 | 1 | 1 | 1 | 1 | 2 | 3 | |
| Cardiac | | | | | | | | | | | 1 | |
| Abdominal | | | 1 | | 1 | | | | 2 | 4 | 1 | |
| Facial | | | | | | | | | | | 1 | 1 |
| Cephalic | | | | | | | | | | | | |
| Blisters | | | | | | | | | | | | |
| Tail | | | | | | | | | | | | |
| Notochord | | 1 | | | | | | | | | 1 | |
| Fin | | | | | | | | | | | | |
| Face | | | | | | | 2 | | | | 1 | |
| Eye | | | | | | | | | | | | |
| Brain | | | | | | | | | | | | |
| Hemorrhage | | | | | | | | | | | | |
| Cardiac | | | | | | | | | | | | |
| Other | | | | | | | | | | | | |

APPENDIX 40

FROG EMBRYO TERATOGENESIS ASSAY - Xenopus (FETAX)
CONDUCTED ON BEACH POINT RAW (pH \approx 4)
GROUNDWATER (WELL 33B)
(TEST NO. 2)

| | |
|---------------------------|---|
| Test Method: | ASTM Designation E 1439-91 ASTM (1991) |
| Type of Test: | Static renewal (every 24 h) |
| Date: | June 3-7, 1993 |
| Investigator: | S. D. Turley |
| Laboratory: | UMD/WREC |
| Groundwater: | |
| Source: | APG-EA Beach Point Well 33B |
| Chemical Characteristics: | See Appendix 48 |
| Test Medium: | |
| Source: | FETAX solution |
| pH characteristics: | See Table A40-1 |
| Test Organism: | |
| Scientific Name: | <u>Xenopus laevis</u> |
| Age at Start of Test: | Stage 8 blastula to stage 11 gastrulae |
| Source: | UMD/WREC culture |
| Experimental Chambers: | |
| Material: | Glass petri dishes |
| Test Solution Volume: | 10 mL |
| No. Organisms/Replicate: | 25 |
| No. Organisms/Treatment: | Control: 100 Groundwater: 50 |
| Loading: | n/a |
| Lighting: | Fluorescent; 60-85 foot candles |
| Aeration: | Prior to renewals |

Endpoints:

Mortality; malformation

Test Temperature:

24 ± 0.2°C

Results:

Mortality:

The raw groundwater did not affect embryo survival. The data are summarized in Tables A40-2 and A40-3.

Malformations:

Significant ($\alpha = 0.05$) embryo malformations occurred at all concentrations down to 18% raw groundwater by volume (see Tables A40-2, A40-4, and A40-5). The 96-h EC50 (malformations), which was determined by the moving average angle method, NOEC and LOEC for the embryos, based on increased numbers of malformations, are as follows:

96-h EC50 = 88.7% raw groundwater by volume (95% confidence limits = 71.1- 143.1).

NOEC = 10% raw groundwater by volume.

LOEC = 18% raw groundwater by volume.

The types of malformed embryos are given in Table A40-6.

TABLE A40-1. SUMMARY OF THE BEACH POINT RAW GROUNDWATER
BIOASSAY pH (STANDARD UNITS) DATA FOR FETAX (TEST
NO. 2)

| | <u>Test Concentrations (Percent Groundwater by Volume)</u> | | | | | |
|--------------|--|------|------|------|------|------|
| | 0 | 10 | 18 | 32 | 56 | 100 |
| <u>Day 0</u> | | | | | | |
| 0 H | 7.50 | 7.48 | 7.46 | 7.37 | 7.07 | 3.99 |
| <u>Day 1</u> | | | | | | |
| 0 H | 7.65 | 7.64 | 7.61 | 7.44 | 7.30 | 3.90 |
| <u>Day 2</u> | | | | | | |
| 0 H | 7.61 | 7.57 | 7.50 | 7.30 | 6.90 | 3.92 |
| <u>Day 3</u> | | | | | | |
| 0 H | 7.53 | 7.45 | 7.40 | 7.29 | 6.97 | 3.93 |
| <u>Day 4</u> | | | | | | |
| 24 H | 7.59 | 7.38 | 7.23 | 6.86 | 6.63 | 4.17 |

TABLE A40-2. FETAX BEACH POINT RAW GROUNDWATER TOXICITY TEST
DATA (TEST NO. 2) - PERCENT EMBRYO SURVIVAL AND
MALFORMATIONS AFTER 96 HOURS OF EXPOSURE^a

| Conc (% by Vol) | Rep | Number Embryos Alive | Percent Survival | Number Embryos Malformed | Percent Malformed |
|----------------------------|-----|----------------------------|---------------------|--------------------------------|----------------------|
| UMD/WREC Control | 1 | 24 | 96 | 2 | 8.3 |
| | 2 | 23 | 92 | 1 | 4.3 |
| | 3 | 24 | 96 | 2 | 8.3 |
| | 4 | 24 | 96 | 2 | 8.3 |
| APG-EA Diluent Water | 1 | 24 | 96 | 2 | 8.3 |
| | 2 | 23 | 92 | 3 | 13.0 |
| 10 | 1 | 25 | 100 | 3 | 12.0 |
| | 2 | 23 | 92 | 2 | 8.7 |
| 18 | 1 | 25 | 100 | 3 | 12.0 |
| | 2 | 24 | 96 | 4 | 16.7 |
| 32 | 1 | 24 | 96 | 3 | 12.5 |
| | 2 | 22 | 88 | 4 | 18.1 |
| 56 | 1 | 22 | 88 | 6 | 27.3 |
| | 2 | 23 | 92 | 6 | 26.1 |
| 100 | 1 | 25 | 100 | 14 | 56.0 |
| | 2 | 24 | 96 | 13 | 54.2 |

TABLE A40-3. FETAX BEACH POINT RAW GROUNDWATER TOXICITY TEST
STATISTICAL ANALYSIS (TEST NO. 2) - PERCENT EMBRYO
SURVIVAL AFTER 96 HOURS OF EXPOSURE

Data Transformation:

Arc sine square root

Chi-Square Test for Normality:

| | |
|----------------------------|---|
| Calculated test statistic: | 7.42 |
| Alpha value: | 0.01 |
| Critical value: | 13.28 |
| Conclusion: | Fail to reject the null hypothesis that the data are normally distributed |

Bartlett's Test for Homogeneity of Variances:

| | |
|----------------------------|--|
| Calculated test statistic: | 2.03 |
| Alpha value: | 0.01 |
| Critical value: | 16.81 |
| Conclusion: | Fail to reject the null hypothesis that the variances are homogenous |

ANOVA:

| | |
|----------------------------|--|
| Calculated test statistic: | 1.33 |
| Alpha value: | 0.05 |
| Critical value: | 3.37 |
| Conclusion: | Fail to reject the null hypothesis that all groups are equal |

TABLE A40-4. FETAX BEACH POINT RAW GROUNDWATER TOXICITY TEST
STATISTICAL ANALYSIS (TEST NO. 2) - PERCENT EMBRYO
MALFORMATIONS AFTER 96 HOURS OF EXPOSURE

Data Transformation:

Arc sine square root

Chi-Square Test for Normality:

| | |
|----------------------------|---|
| Calculated test statistic: | 7.42 |
| Alpha value: | 0.01 |
| Critical Value: | 13.28 |
| Conclusion: | Fail to reject the null hypothesis that the data are normally distributed |

Bartlett's Test for Homogeneity of Variances:

| | |
|----------------------------|--|
| Calculated test statistic: | 4.04 |
| Alpha value: | 0.01 |
| Critical value: | 16.81 |
| Conclusion: | Fail to reject the null hypothesis that the variances are homogenous |

ANOVA:

| | |
|----------------------------|--|
| Calculated test statistic: | 42.00 |
| Alpha value: | 0.05 |
| Critical value: | 3.37 |
| Conclusion: | Reject the null hypothesis that all groups are equal |

Bonferroni's T-Test:

| | |
|----------------------------|--|
| Calculated test statistic: | See Table A40-5 |
| Alpha value: | 0.05 |
| Critical value: | 2.94 |
| Conclusion: | Reject the null hypothesis that all groups are equal |

TABLE A40-5. FETAX BEACH POINT RAW GROUNDWATER TOXICITY TEST
STATISTICAL ANALYSIS (TEST NO. 2) - RESULTS OF
BONFERRONI'S T-TEST ON EMBRYO MALFORMATIONS AFTER
96 HOURS OF EXPOSURE

| Conc (% by Vol) | No. of Reps | Mean Normal Embryos (%) ^a | T Statistic | Significance |
|-----------------------|-------------------|---|-------------|--------------|
| UMD/WREC Control | 4 | 93.0 | | |
| 10 | 2 | 89.5 | 1.61 | |
| 18 | 2 | 85.5 | 3.20 | * |
| 32 | 2 | 84.5 | 3.57 | * |
| 56 | 2 | 73.5 | 7.11 | * |
| 100 | 2 | 45.0 | 14.72 | * |

^a Values given are actual percent mean normal embryos rather than arc sine square root transformed means which were used in the statistical analysis.

* Significantly different at alpha = 0.05 (Bonferroni's critical value = 2.94).

TABLE A40-6. FETAX BEACH POINT RAW GROUNDWATER TOXICITY TEST
DATA (TEST NO. 2) - TYPE AND NUMBER OF MALFORMED
EMBRYOS AFTER 96 HOURS OF EXPOSURE^a

| Malformation | Test Concentrations (% Groundwater by Volume) | | | | | | | | | |
|--------------|---|---|---|---|------------|---|------------|---|------------|---|
| | 0 | | | | 10 | | 18 | | 32 | |
| | <u>Rep</u> | | | | <u>Rep</u> | | <u>Rep</u> | | <u>Rep</u> | |
| | 1 | 2 | 3 | 4 | 1 | 2 | 1 | 2 | 1 | 2 |
| Severe | | | | | | | | | 1 | 3 |
| Gut, coiling | | | | | | | 2 | 3 | 2 | 1 |
| Edema: | | | | | | | | | | |
| Multiple | 2 | 1 | 1 | | 3 | 2 | 1 | 1 | 1 | 1 |
| Cardiac | | | | | | | | | | |
| Abdominal | | | | | | | | | 1 | 1 |
| Facial | | | | | | | | | | |
| Cephalic | | | | | | | | | | |
| Blisters | | | | | | | | | | |
| Tail | | | | | | | | | | |
| Notochord | 1 | 1 | 1 | | | | | | | |
| Fin | | | | | | | | | | |
| Face | | | | | | | | | | 1 |
| Eye | | | | | | | | | | |
| Brain | | | | | | | | | | |
| Hemorrhage | | | | | | | | | | |
| Cardiac | | | | | | | | | | |
| Other | | | | | | | | | | |

^a The APG-EA diluent water controls had the following the malformations: Rep 1- 2 embryos with multiple edema. Rep 2- 1 severe and 2 multiple edema.

APPENDIX 41

FROG EMBRYO TERATOGENESIS ASSAY - Xenopus (FETAX)
CONDUCTED ON BEACH POINT BUFFERED (pH \approx 7)
GROUNDWATER (WELL 33B)
(TEST NO. 2)

| | |
|---------------------------|---|
| Test Method: | ASTM Designation E 1439-91 ASTM (1991) |
| Type of Test: | Static renewal (every 24 h) |
| Date: | June 3-7, 1993 |
| Investigator: | S. D. Turley |
| Laboratory: | UMD/WREC |
| Groundwater: | |
| Source: | APG-EA Beach Point Well 33B |
| Chemical Characteristics: | See Appendix 48 |
| Test Medium: | |
| Source: | FETAX solution |
| pH characteristics: | See Table A41-1 |
| Test Organism: | |
| Scientific Name: | <u>Xenopus laevis</u> |
| Age at Start of Test: | Stage 8 blastula to stage 11 gastrulae |
| Source: | UMD/WREC culture |
| Experimental Chambers: | |
| Material: | Glass petri dishes |
| Test Solution Volume: | 10 mL |
| No. Organisms/Replicate: | 25 |
| No. Organisms/Treatment: | Groundwater: 50 Control: 100 |
| Loading: | n/a |
| Lighting: | Fluorescent; 60-85 foot candles |
| Aeration: | Prior to renewals |

| | |
|-------------------|--------------------------|
| pH Buffer: | 10 N NaOH |
| Endpoints: | Mortality; malformations |
| Test Temperature: | 24 ± 0.2°C |

Results:

Mortality:

Less than 50% mortality occurred to the embryos exposed for 4 days to 100% Beach Point buffered groundwater; thus, an LC50 could not be calculated (Table A41-2). A significant increase in embryo mortality occurred at 56% and 100% buffered groundwater by volume (see Tables A41-2, A41-3 and A41-4).

Malformations:

Significant ($\alpha = 0.05$) embryo malformations occurred at all concentrations down to 32% buffered groundwater by volume (see Tables A41-2, A41-4, and A41-5). Less than 50% malformation occurred at all test concentrations, thus, an EC50 could not be calculated (Table A41-2). The NOEC and LOEC for the embryos, based on increased numbers of malformations, are as follows:

NOEC = 18% buffered groundwater by volume.
LOEC = 32% buffered groundwater by volume.

The types of malformed embryos are given in Table A41-7.

TABLE A41-1. SUMMARY OF THE BEACH POINT BUFFERED GROUNDWATER
BIOASSAY pH (STANDARD UNITS) DATA FOR FETAX (TEST
NO. 2)

| | <u>Test Concentrations (Percent Groundwater by Volume)</u> | | | | | |
|--------------|--|------|------|------|------|------|
| | 0 | 10 | 18 | 32 | 56 | 100 |
| <u>Day 0</u> | | | | | | |
| 0 H | 7.50 | 7.47 | 7.40 | 7.45 | 7.57 | 7.55 |
| <u>Day 1</u> | | | | | | |
| 0 H | 7.65 | 6.98 | 7.21 | 7.38 | 7.48 | 7.55 |
| <u>Day 2</u> | | | | | | |
| 0 H | 7.61 | 7.66 | 7.81 | 7.91 | 7.92 | 7.88 |
| <u>Day 3</u> | | | | | | |
| 0 H | 7.53 | 7.44 | 7.48 | 7.43 | 7.51 | 7.56 |
| <u>Day 4</u> | | | | | | |
| 24 H | 7.57 | 7.37 | 7.46 | 7.52 | 7.44 | 7.59 |

TABLE A41-2. FETAX BEACH POINT BUFFERED GROUNDWATER TOXICITY
TEST DATA (TEST NO. 2) - PERCENT EMBRYO SURVIVAL
AND MALFORMATIONS AFTER 96 HOURS OF EXPOSURE

| Conc (% by Vol) | Rep | Number Embryos Alive | Percent Survival | Number Embryos Malformed | Percent Malformed |
|-----------------------|-----|----------------------------|---------------------|--------------------------------|----------------------|
| UMD/WREC Control | 1 | 24 | 96 | 2 | 8.3 |
| | 2 | 23 | 92 | 1 | 4.3 |
| | 3 | 24 | 96 | 2 | 4.3 |
| | 4 | 24 | 96 | 2 | 8.3 |
| 10 | 1 | 24 | 96 | 3 | 12.5 |
| | 2 | 23 | 92 | 1 | 4.3 |
| 18 | 1 | 23 | 92 | 2 | 8.7 |
| | 2 | 24 | 96 | 2 | 8.3 |
| 32 | 1 | 22 | 88 | 4 | 18.1 |
| | 2 | 23 | 92 | 6 | 26.1 |
| 56 | 1 | 21 | 84 | 6 | 28.6 |
| | 2 | 22 | 88 | 3 | 13.6 |
| 100 | 1 | 20 | 80 | 4 | 20.0 |
| | 2 | 21 | 84 | 7 | 33.3 |

TABLE A41-3. FETAX BEACH POINT BUFFERED GROUNDWATER TOXICITY
TEST STATISTICAL ANALYSIS (TEST NO. 21) - PERCENT
EMBRYO SURVIVAL AFTER 96 HOURS OF EXPOSURE

Data Transformation:

Arc sine square root

Chi-Square Test for Normality:

| | |
|----------------------------|---|
| Calculated test statistic: | 5.69 |
| Alpha value: | 0.01 |
| Critical value: | 13.28 |
| Conclusion: | Fail to reject the null hypothesis that the data are normally distributed |

Bartlett's Test for Homogeneity of Variances:

| | |
|----------------------------|--|
| Calculated test statistic: | 0.45 |
| Alpha value: | 0.01 |
| Critical value: | 13.28 |
| Conclusion: | Fail to reject the null hypothesis that the variances are homogenous |

ANOVA:

| | |
|----------------------------|--|
| Calculated test statistic: | 9.00 |
| Alpha value: | 0.05 |
| Critical value: | 3.69 |
| Conclusion: | Reject the null hypothesis that all groups are equal |

Bonferroni's T-test:

| | |
|----------------------------|--|
| Calculated test statistic: | See Table A41-4 |
| Alpha value: | 0.05 |
| Critical value: | 2.90 |
| Conclusion: | Reject the null hypothesis that all groups are equal |

TABLE A41-4. FETAX BEACH POINT BUFFERED GROUNDWATER TOXICITY
STATISTICAL ANALYSIS (TEST NO. 2) - RESULTS OF
BONFERRONI'S T-TEST ON EMBRYO MORTALITY AFTER 96
HOURS OF EXPOSURE

| Conc (% by Vol) | No. of Reps | Mean Embryo Survival (%) _a | T Statistic | Significance |
|-----------------------|-------------------|--|-------------|--------------|
| UMD/WREC Control | 4 | 95.0 | | |
| 10 | 2 | 94.0 | 0.55 | |
| 18 | 2 | 94.0 | 0.55 | |
| 32 | 2 | 90.0 | 2.52 | |
| 56 | 2 | 86.0 | 4.13 | * |
| 100 | 2 | 82.0 | 5.55 | * |

_a Values given are actual percent mean surviving embryos rather than arc sine square root transformed means which were used in the statistical analysis.

* Significantly different at Alpha = 0.05 (Bonferroni's critical value = 2.90)

TABLE A41-5. FETAX BEACH POINT BUFFERED GROUNDWATER TOXICITY
TEST STATISTICAL ANALYSIS (TEST NO. 2) - PERCENT
EMBRYO MALFORMATIONS AFTER 96 HOURS OF EXPOSURE^a

Data Transformation:

Arc sine square root

Chi-Square Test for Normality:

| | |
|----------------------------|---|
| Calculated test statistic: | 10.66 |
| Alpha value: | 0.01 |
| Critical Value: | 13.28 |
| Conclusion: | Fail to reject the null hypothesis that the data are normally distributed |

Bartlett's Test for Homogeneity of Variances:

| | |
|----------------------------|--|
| Calculated test statistic: | 5.85 |
| Alpha value: | 0.01 |
| Critical value: | 11.34 |
| Conclusion: | Fail to reject the null hypothesis that the variances are homogenous |

ANOVA:

| | |
|----------------------------|--|
| Calculated test statistic: | 6.50 |
| Alpha value: | 0.05 |
| Critical value: | 4.76 |
| Conclusion: | Reject the null hypothesis that all groups are equal |

Bonferroni's T-Test:

| | |
|----------------------------|--|
| Calculated test statistic: | See Table A41-6 |
| Alpha value: | 0.05 |
| Critical value: | 2.75 |
| Conclusion: | Reject the null hypothesis that all groups are equal |

^a The 56% and 100% buffered Beach Point groundwater treatment was not included in the statistical analyses because all concentrations above the NOEC for survival are excluded from the hypothesis test for malformation effects.

TABLE A41-6. FETAX BEACH POINT BUFFERED GROUNDWATER TOXICITY
STATISTICAL ANALYSIS (TEST NO. 2) - RESULTS OF
BONFERRONI'S T-TEST ON EMBRYO MALFORMATIONS AFTER
96 HOURS OF EXPOSURE

| Conc (% by Vol) | No. of Reps | Mean Normal Embryos (%) ^a | T Statistic | Significance |
|-----------------------|-------------------|---|-------------|--------------|
| UMD/WREC Control | 4 | 93.0 | | |
| 10 | 2 | 91.5 | 0.63 | |
| 18 | 2 | 91.5 | 0.83 | |
| 32 | 2 | 78.0 | 4.33 | * |

^a Values given are actual percent mean normal embryos rather than arc sine square root transformed means which were used in the statistical analysis.

* Significantly different at alpha = 0.05 (Bonferroni's critical value = 2.75).

TABLE A41-7. FETAX BEACH POINT BUFFERED GROUNDWATER TOXICITY
TEST DATA (TEST NO. 2) - TYPE AND NUMBER OF
MALFORMED EMBRYOS AFTER 96 HOURS OF EXPOSURE

| Malformation | Test Concentrations (% Groundwater by Volume) | | | | | | | | | | | | | |
|--------------|---|---|---|---|------------|---|------------|---|------------|---|------------|---|------------|---|
| | 0 | | | | 10 | | 18 | | 32 | | 56 | | 100 | |
| | <u>Rep</u> | | | | <u>Rep</u> | | <u>Rep</u> | | <u>Rep</u> | | <u>Rep</u> | | <u>Rep</u> | |
| | 1 | 2 | 3 | 4 | 1 | 2 | 1 | 2 | 1 | 2 | 1 | 2 | 1 | 2 |
| Severe | | | | | 1 | | 1 | | | | | | | |
| Gut, coiling | | | | | | | | | 3 | 3 | 3 | 2 | 3 | 3 |
| Edema: | | | | | | | | | | | | | | |
| Multiple | 2 | | 1 | 1 | 1 | | 1 | 1 | 1 | 3 | 3 | 1 | 1 | 1 |
| Cardiac | | | | | | | | | | | | | | |
| Abdominal | | | | | | 1 | | 1 | | | | | | 1 |
| Facial | | | | | | | | | | | | | | |
| Cephalic | | | | | | | | | | | | | | |
| Blisters | | | | | | | | | | | | | | |
| Tail | | | | | | | | | | | | | | |
| Notochord | | 1 | 1 | 1 | | 1 | | | | | | | | |
| Fin | | | | | | | | | | | | | | |
| Face | | | | | | | | | | | | | | 2 |
| Eye | | | | | | | | | | | | | | |
| Brain | | | | | | | | | | | | | | |
| Hemorrhage | | | | | | | | | | | | | | |
| Cardiac | | | | | | | | | | | | | | |
| Other | | | | | | | | | | | | | | |

APPENDIX 42

FROG EMBRYO TERATOGENESIS ASSAY - Xenopus (FETAX)
CONDUCTED ON BEACH POINT RAW (pH \approx 4)
GROUNDWATER (WELL 33B)
(TEST NO. 3)

| | |
|---------------------------|---|
| Test Method: | ASTM Designation E 1439-91 ASTM (1991) |
| Type of Test: | Static renewal (every 24 h) |
| Date: | August 26-30, 1993 |
| Investigator: | S. D. Turley |
| Laboratory: | UMD/WREC |
| Groundwater: | |
| Source: | APG-EA Beach Point Well 33B |
| Chemical Characteristics: | See Appendix 48 |
| Test Medium: | |
| Source: | FETAX solution |
| pH characteristics: | See Table A42-1 |
| Test Organism: | |
| Scientific Name: | <u>Xenopus laevis</u> |
| Age at Start of Test: | Stage 8 blastula to stage 11 gastrulae |
| Source: | UMD/WREC culture |
| Experimental Chambers: | |
| Material: | Glass petri dishes |
| Test Solution Volume: | 10 mL |
| No. Organisms/Replicate: | 25 |
| No. Organisms/Treatment: | Control: 100 Groundwater: 50 |
| Loading: | n/a |
| Lighting: | Fluorescent; 60-85 foot candles |
| Aeration: | Prior to renewals |

Endpoints:

Mortality; malformation

Test Temperature:

24 ± 0.2 °C

Results:

Mortality:

The raw groundwater did not affect embryo survival. The data are summarized in Tables A42-2 and A42-3.

Malformations:

Significant ($\alpha = 0.05$) embryo malformations occurred at all concentrations down to 18% raw groundwater by volume (see Tables A42-2, A42-4, and A42-5). The NOEC and LOEC for the embryos, based on increased numbers of malformations, are as follows:

NOEC = 10% raw groundwater by volume.

LOEC = 18% raw groundwater by volume.

The types of malformed embryos are given in Table A42-6.

TABLE A42-1. SUMMARY OF THE BEACH POINT RAW GROUNDWATER
BIOASSAY pH (STANDARD UNITS) DATA FOR FETAX (TEST
NO. 3)

| | <u>Test Concentrations (Percent Groundwater by Volume)</u> | | | | | |
|--------------|--|------|------|------|------|------|
| | 0 | 10 | 18 | 32 | 56 | 100 |
| <u>Day 0</u> | | | | | | |
| 0 H | 7.70 | 7.51 | 7.21 | 6.93 | 6.32 | 3.97 |
| <u>Day 1</u> | | | | | | |
| 0 H | 7.62 | 7.38 | 7.09 | 6.77 | 6.28 | 4.07 |
| <u>Day 2</u> | | | | | | |
| 0 H | 7.59 | 7.26 | 7.17 | 7.00 | 6.33 | 4.02 |
| <u>Day 3</u> | | | | | | |
| 0 H | 7.65 | 7.32 | 7.11 | 6.69 | 6.07 | 3.95 |
| <u>Day 4</u> | | | | | | |
| 24 H | 7.51 | 7.20 | 7.01 | 6.58 | 6.01 | 4.01 |

TABLE A42-2. FETAX BEACH POINT RAW GROUNDWATER TOXICITY TEST
DATA (TEST NO. 3) - PERCENT EMBRYO SURVIVAL AND
MALFORMATIONS AFTER 96 HOURS OF EXPOSURE

| Conc (% by Vol) | Rep | Number Embryos Alive | Percent Survival | Number Embryos Malformed | Percent Malformed |
|-----------------------|-----|----------------------------|---------------------|--------------------------------|----------------------|
| UMD/WREC Control | 1 | 25 | 100 | 1 | 4.0 |
| | 2 | 24 | 96 | 2 | 8.3 |
| | 3 | 24 | 96 | 0 | 0.0 |
| | 4 | 24 | 96 | 1 | 4.2 |
| APG-EA Control | 1 | 24 | 96 | 2 | 8.3 |
| | 2 | 25 | 100 | 1 | 4.0 |
| 10 | 1 | 24 | 96 | 2 | 8.3 |
| | 2 | 23 | 92 | 2 | 8.7 |
| 18 | 1 | 23 | 92 | 3 | 13.0 |
| | 2 | 25 | 100 | 3 | 12.0 |
| 32 | 1 | 23 | 92 | 3 | 13.0 |
| | 2 | 24 | 96 | 4 | 16.7 |
| 56 | 1 | 25 | 100 | 4 | 16.0 |
| | 2 | 22 | 88 | 4 | 18.2 |
| 100 | 1 | 24 | 96 | 9 | 37.5 |
| | 2 | 22 | 88 | 10 | 45.5 |

TABLE A42-3. FETAX BEACH POINT RAW GROUNDWATER TOXICITY TEST
STATISTICAL ANALYSIS (TEST NO. 3) - PERCENT EMBRYO
SURVIVAL AFTER 96 HOURS OF EXPOSURE

Data Transformation:

Arc sine square root

Chi-Square Test for Normality:

| | |
|----------------------------|---|
| Calculated test statistic: | 7.43 |
| Alpha value: | 0.01 |
| Critical value: | 13.28 |
| Conclusion: | Fail to reject the null hypothesis that the data are normally distributed |

Bartlett's Test for Homogeneity of Variances:

| | |
|----------------------------|--|
| Calculated test statistic: | 2.74 |
| Alpha value: | 0.01 |
| Critical value: | 16.81 |
| Conclusion: | Fail to reject the null hypothesis that the variances are homogenous |

ANOVA:

| | |
|----------------------------|--|
| Calculated test statistic: | 0.44 |
| Alpha value: | 0.05 |
| Critical value: | 3.37 |
| Conclusion: | Fail to reject the null hypothesis that all groups are equal |

TABLE A42-4. FETAX BEACH POINT RAW GROUNDWATER TOXICITY TEST
STATISTICAL ANALYSIS (TEST NO. 3) - PERCENT EMBRYO
MALFORMATIONS AFTER 96 HOURS OF EXPOSURE

Data Transformation:

Arc sine square root

Chi-Square Test for Normality:

| | |
|----------------------------|---|
| Calculated test statistic: | 9.96 |
| Alpha value: | 0.01 |
| Critical Value: | 13.28 |
| Conclusion: | Fail to reject the null hypothesis that the data are normally distributed |

Bartlett's Test for Homogeneity of Variances:

| | |
|----------------------------|--|
| Calculated test statistic: | 7.46 |
| Alpha value: | 0.01 |
| Critical value: | 16.81 |
| Conclusion: | Fail to reject the null hypothesis that the variances are homogenous |

ANOVA:

| | |
|----------------------------|--|
| Calculated test statistic: | 21.00 |
| Alpha value: | 0.05 |
| Critical value: | 3.37 |
| Conclusion: | Reject the null hypothesis that all groups are equal |

Bonferroni's T-test:

| | |
|----------------------------|--|
| Calculated test statistic: | See Table A42-5 |
| Alpha value: | 0.05 |
| Critical value: | 2.94 |
| Conclusion: | Reject the null hypothesis that all groups are equal |

TABLE A42-5. FETAX BEACH POINT RAW GROUNDWATER TOXICITY TEST
STATISTICAL ANALYSIS (TEST NO. 3) - RESULTS OF
BONFERRONI'S T-TEST ON EMBRYO MALFORMATIONS AFTER
96 HOURS OF EXPOSURE

| Conc (% by Vol) | No. of Reps | Mean Normal Embryos (%) ^a | T Statistic | Significance |
|-----------------------|-------------------|---|-------------|--------------|
| UMD/WREC Control | 4 | 95.9 | | |
| APG-EA | 2 | 93.9 | 0.99 | |
| 10 | 2 | 91.5 | 2.02 | |
| 18 | 2 | 87.5 | 3.40 | * |
| 32 | 2 | 85.1 | 4.11 | * |
| 56 | 2 | 83.0 | 4.77 | * |
| 100 | 2 | 58.7 | 10.53 | * |

^a Values given are actual percent mean normal embryos rather than arc sine square root transformed means which were used in the statistical analysis.

* Significantly different at alpha = 0.05 (Bonferroni's critical value = 2.94).

TABLE A42-6. FETAX BEACH POINT RAW GROUNDWATER TOXICITY TEST DATA (TEST NO. 3) - TYPE AND NUMBER OF MALFORMED EMBRYOS AFTER 96 HOURS OF EXPOSURE.

| Malformation | Test Concentrations (% Groundwater by Volume) | | | | | | | | | | | |
|--------------|---|---|---|---|------------------|---|------------|------------|------------|------------|------------|---|
| | UMD ^a | | | | APG ^b | | 10 | 18 | 32 | 56 | 100 | |
| | <u>Rep</u> | | | | <u>Rep</u> | | <u>Rep</u> | <u>Rep</u> | <u>Rep</u> | <u>Rep</u> | <u>Rep</u> | |
| | 1 | 2 | 3 | 4 | 1 | 2 | 1 | 2 | 1 | 2 | 1 | 2 |
| <hr/> | | | | | | | | | | | | |
| Severe | | | | | | | | | | | | |
| Gut, coiling | 1 | | | 1 | 2 | | 1 | 2 | 1 | 2 | 1 | 3 |
| Edema: | | | | | | | | | | | | |
| Multiple | | 2 | | | 1 | 1 | 2 | 1 | 1 | 2 | 2 | 7 |
| Cardiac | | | | | | | | | | | | |
| Abdominal | | | | | | | | | | 1 | | 1 |
| Facial | | | | | | | | | | | | |
| Cephalic | | | | | | | | | | | | |
| Blisters | | | | | | | | | | | | |
| Tail | | | | | | | | | | | | |
| Notochord | | | | | | | | | | | | |
| Fin | | | | | | | | | | | | |
| Face | | | | | | | | 1 | | | 1 | 1 |
| Eye | | | | | | | | | | | | |
| Brain | | | | | | | | | | | | |
| Hemorrhage | | | | | | | | | | | | |
| Cardiac | | | | | | | | | | | | |
| Other | | | | | | | | | | | | |

^a UMD/WREC control.

^b APG-EA control.

APPENDIX 43

FROG EMBRYO TERATOGENESIS ASSAY - Xenopus (FETAX) CONDUCTED ON BEACH POINT BUFFERED (pH \approx 7) GROUNDWATER (WELL 33B) (TEST NO. 3)

| | |
|---------------------------|---|
| Test Method: | ASTM Designation E 1439-91 ASTM (1991) |
| Type of Test: | Static renewal (every 24 h) |
| Date: | August 26-30, 1993 |
| Investigator: | S. D. Turley |
| Laboratory: | UMD/WREC |
| Groundwater: | |
| Source: | APG-EA Beach Point Well 33B |
| Chemical Characteristics: | See Appendix 48 |
| Test Medium: | |
| Source: | FETAX solution |
| pH characteristics: | See Table A43-1 |
| Test Organism: | |
| Scientific Name: | <u>Xenopus laevis</u> |
| Age at Start of Test: | Stage 8 blastula to stage 11 gastrulae |
| Source: | UMD/WREC culture |
| Experimental Chambers: | |
| Material: | Glass petri dishes |
| Test Solution Volume: | 10 mL |
| No. Organisms/Replicate: | 25 |
| No. Organisms/Treatment: | Groundwater: 50 Control: 100 |
| Loading: | n/a |
| Lighting: | Fluorescent; 60-85 foot candles |
| Aeration: | Prior to renewals |

| | |
|-------------------|--------------------------|
| pH Buffer: | 10 N NaOH |
| Endpoints: | Mortality; malformations |
| Test Temperature: | 24 ± 0.2 °C |

Results:

Mortality:

The buffered groundwater did not affect embryo survival.
The data are summarized in Tables A43-2 and A43-3.

Malformations:

Significant ($\alpha = 0.05$) embryo malformations occurred at all concentrations down to 32% buffered groundwater by volume (see Tables A43-2, A43-4, and A43-5). The NOEC and LOEC for the embryos, based on increased numbers of malformations, are as follows:

NOEC = 18% buffered groundwater by volume.

LOEC = 32% buffered groundwater by volume.

The types of malformed embryos are given in Table A43-6.

TABLE A43-1. SUMMARY OF THE BEACH POINT BUFFERED GROUNDWATER
BIOASSAY pH (STANDARD UNITS) DATA FOR FETAX (TEST
NO. 3)

| | <u>Test Concentrations (Percent Groundwater by Volume)</u> | | | | | |
|--------------|--|------|------|------|------|------|
| | 0 | 10 | 18 | 32 | 56 | 100 |
| <u>Day 0</u> | | | | | | |
| 0 H | 7.70 | 7.48 | 7.41 | 7.38 | 7.30 | 7.25 |
| <u>Day 1</u> | | | | | | |
| 0 H | 7.62 | 7.58 | 7.51 | 7.37 | 7.33 | 7.30 |
| <u>Day 2</u> | | | | | | |
| 0 H | 7.59 | 7.63 | 7.55 | 7.43 | 7.37 | 7.57 |
| <u>Day 3</u> | | | | | | |
| 0 H | 7.65 | 7.50 | 7.44 | 7.41 | 7.33 | 7.21 |
| <u>Day 4</u> | | | | | | |
| 24 H | 7.50 | 7.36 | 7.30 | 7.23 | 7.17 | 7.09 |

TABLE A43-2. FETAX BEACH POINT BUFFERED GROUNDWATER TOXICITY
TEST DATA (TEST NO. 3) - PERCENT EMBRYO SURVIVAL
AND MALFORMATIONS AFTER 96 HOURS OF EXPOSURE

| Conc (% by Vol) | Rep | Number Embryos Alive | Percent Survival | Number Embryos Malformed | Percent Malformed |
|-----------------------|-----|----------------------------|---------------------|--------------------------------|----------------------|
| UMD/WREC Control | 1 | 25 | 100 | 1 | 4.0 |
| | 2 | 24 | 96 | 2 | 8.3 |
| | 3 | 24 | 96 | 0 | 0.0 |
| | 4 | 24 | 96 | 1 | 4.2 |
| 10 | 1 | 24 | 96 | 1 | 4.2 |
| | 2 | 25 | 100 | 2 | 8.0 |
| 18 | 1 | 24 | 96 | 2 | 8.3 |
| | 2 | 23 | 92 | 3 | 13.0 |
| 32 | 1 | 24 | 96 | 3 | 12.5 |
| | 2 | 23 | 92 | 3 | 13.0 |
| 56 | 1 | 24 | 96 | 4 | 16.7 |
| | 2 | 25 | 100 | 4 | 16.0 |
| 100 | 1 | 24 | 96 | 5 | 20.8 |
| | 2 | 23 | 92 | 7 | 30.4 |

TABLE A43-3. FETAX BEACH POINT BUFFERED GROUNDWATER TOXICITY
TEST STATISTICAL ANALYSIS (TEST NO. 3) - PERCENT
EMBRYO SURVIVAL AFTER 96 HOURS OF EXPOSURE

Data Transformation:

Arc sine square root

Chi-Square Test for Normality:

| | |
|----------------------------|---|
| Calculated test statistic: | 5.69 |
| Alpha value: | 0.01 |
| Critical value: | 13.28 |
| Conclusion: | Fail to reject the null hypothesis that the data are normally distributed |

Bartlett's Test for Homogeneity of Variance:

| | |
|----------------------------|--|
| Calculated test statistic: | 0.12 |
| Alpha value: | 0.01 |
| Critical value: | 15.09 |
| Conclusion: | Fail to reject the null hypothesis that all groups are equal |

ANOVA:

| | |
|----------------------------|--|
| Calculated test statistic: | 1.33 |
| Alpha value: | 0.05 |
| Critical value: | 3.69 |
| Conclusion: | Fail to reject the null hypothesis that all groups are equal |

TABLE A43-4. FETAX BEACH POINT BUFFERED GROUNDWATER TOXICITY
TEST STATISTICAL ANALYSIS (TEST NO. 3) - PERCENT
EMBRYO MALFORMATIONS AFTER 96 HOURS OF EXPOSURE

Data Transformation:

Arc sine square root

Chi-Square Test for Normality:

| | |
|----------------------------|---|
| Calculated test statistic: | 8.00 |
| Alpha value: | 0.01 |
| Critical Value: | 13.28 |
| Conclusion: | Fail to reject the null hypothesis that the data are normally distributed |

Bartlett's Test for Homogeneity of Variances:

| | |
|----------------------------|--|
| Calculated test statistic: | 7.80 |
| Alpha value: | 0.01 |
| Critical value: | 15.09 |
| Conclusion: | Fail to reject the null hypothesis that the variances are homogenous |

ANOVA:

| | |
|----------------------------|--|
| Calculated test statistic: | 9.00 |
| Alpha value: | 0.05 |
| Critical value: | 3.69 |
| Conclusion: | Reject the null hypothesis that all groups are equal |

Bonferroni's T-Test:

| | |
|----------------------------|--|
| Calculated test statistic: | See Table A43-5 |
| Alpha value: | 0.05 |
| Critical value: | 2.90 |
| Conclusion: | Reject the null hypothesis that all groups are equal |

TABLE A43-5. FETAX BEACH POINT BUFFERED GROUNDWATER TOXICITY
TEST STATISTICAL ANALYSIS (TEST NO. 3) - RESULTS OF
BONFERRONI'S T-TEST ON EMBRYO MALFORMATIONS AFTER
96 HOURS OF EXPOSURE

| Conc (% by Vol) | No. of Reps | Mean Normal Embryos (%) ^a | T Statistic | Significance |
|-----------------------|-------------------|---|-------------|--------------|
| UMD/WREC Control | 4 | 95.9 | | |
| 10 | 2 | 93.9 | 0.85 | |
| 18 | 2 | 89.4 | 2.38 | |
| 32 | 2 | 87.2 | 3.01 | * |
| 56 | 2 | 83.7 | 3.95 | * |
| 100 | 2 | 74.5 | 6.00 | * |

^a Values given are actual percent mean normal embryos rather than arc sine square root transformed means which were used in the statistical analysis.

* Significantly different at alpha = 0.05 (Bonferroni's critical value = 2.90).

TABLE A43-6. FETAX BEACH POINT BUFFERED GROUNDWATER TOXICITY
TEST DATA (TEST NO. 3) - TYPE AND NUMBER OF
MALFORMED EMBRYOS AFTER 96 HOURS OF EXPOSURE

| Malformation | <u>Test Concentrations (% Groundwater by Volume)</u> | | | | | | | | | |
|--------------|--|---|---|---|------------|---|------------|---|------------|---|
| | 0 | | | | 10 | | 18 | | 32 | |
| | <u>Rep</u> | | | | <u>Rep</u> | | <u>Rep</u> | | <u>Rep</u> | |
| | 1 | 2 | 3 | 4 | 1 | 2 | 1 | 2 | 1 | 2 |
| Severe | | | | | | | | | | |
| Gut, coiling | 1 | | 1 | | | | 1 | 1 | 2 | 1 |
| Edema: | | | | | | | | | | |
| Multiple | | 2 | | | | | 1 | 1 | 2 | |
| Cardiac | | | | | | | | | | |
| Abdominal | | | | | 1 | | 1 | | 3 | 1 |
| Facial | | | | | | | | | | |
| Cephalic | | | | | | | | | | |
| Blisters | | | | | | | | | | |
| Tail | | | | | | | | | | |
| Notochord | | | | | 1 | | | | | |
| Fin | | | | | | | | | | |
| Face | | | | | 1 | | | | | 1 |
| Eye | | | | | | | | | | |
| Brain | | | | | | | | | | |
| Hemorrhage | | | | | | | | | | |
| Cardiac | | | | | | | | | | |
| Other | | | | | | | | | | |

APPENDIX 44

FROG EMBRYO TERATOGENESIS ASSAY - Xenopus (FETAX)
CONDUCTED ON BEACH POINT RAW (pH \approx 4)
GROUNDWATER (WELL 33B)
(TEST NO. 4)

| | |
|---------------------------|--|
| Test Method: | ASTM Designation E 1439-91 (ASTM, 1991) |
| Type of Test: | Static renewal (every 24 h) |
| Date: | December 3 - 7, 1993 |
| Investigator: | S. D. Turley |
| Laboratory: | UMD/WREC |
| Groundwater: | |
| Source: | APG-EA Beach Point Well 33B |
| Chemical Characteristics: | See Appendix 48 |
| Test Medium: | |
| Source: | FETAX solution |
| pH characteristics: | See Table A44-1 |
| Test Organism: | |
| Scientific Name: | <u>Xenopus laevis</u> |
| Age at Start of Test: | Stage 8 blastula to stage 11 gastrulae |
| Source: | UMD/WREC culture |
| Experimental Chambers: | |
| Material: | Glass petri dishes |
| Test Solution Volume: | 10 mL |
| No. Organisms/Replicate: | 25 |
| No. Organisms/Treatment: | Control: 100 Groundwater: 50 |
| Loading: | n/a |
| Lighting: | Fluorescent; 60-85 foot candles |
| Aeration: | Prior to renewals |

Endpoints:

Mortality; malformation

Test Temperature:

24 ± 0.2 °C

Results:

Mortality:

Significant ($\alpha = 0.05$) mortality occurred in frog embryos exposed for 96 h to 100% Beach Point raw groundwater. 100% Beach Point raw groundwater killed 26% of the frog embryos; thus, a 96-h LC50 could not be obtained. The data are summarized in Tables A44-2, A44-3, and A44-4.

Malformations:

Significant ($\alpha = 0.05$) embryo malformations occurred at a concentration of 56% raw groundwater by volume (see Tables A44-2, A44-4, and A44-5). The NOEC and LOEC for the embryos, based on increased numbers of malformations, are as follows:

NOEC = 32% raw groundwater by volume.

LOEC = 56% raw groundwater by volume.

The types of malformed embryos are given in Table A44-6.

TABLE A44-1. SUMMARY OF THE BEACH POINT RAW GROUNDWATER
BIOASSAY pH (STANDARD UNITS) DATA FOR FETAX (TEST
NO. 4)

| | <u>Test Concentrations (Percent Groundwater by Volume)</u> | | | | | |
|--------------|--|------|------|------|------|------|
| | 0 | 10 | 18 | 32 | 56 | 100 |
| <u>Day 0</u> | | | | | | |
| 0 H | 7.51 | 7.33 | 7.18 | 6.99 | 6.54 | 3.91 |
| <u>Day 1</u> | | | | | | |
| 0 H | 7.50 | 7.40 | 7.23 | 7.06 | 6.61 | 3.99 |
| <u>Day 2</u> | | | | | | |
| 0 H | 7.44 | 7.19 | 7.11 | 6.91 | 6.44 | 3.87 |
| <u>Day 3</u> | | | | | | |
| 0 H | 7.49 | 7.27 | 7.14 | 7.00 | 6.52 | 3.88 |

TABLE A44-2. FETAX BEACH POINT RAW GROUNDWATER TOXICITY TEST
DATA (TEST NO. 4) - PERCENT EMBRYO SURVIVAL AND
MALFORMATIONS AFTER 96 HOURS OF EXPOSURE

| Conc (% by Vol) | Rep | Number Embryos Alive | Percent Survival | Number Embryos Malformed | Percent Malformed |
|----------------------------|-----|----------------------------|---------------------|--------------------------------|----------------------|
| UMD/WREC Control | 1 | 25 | 100 | 2 | 8.0 |
| | 2 | 24 | 96 | 2 | 8.3 |
| | 3 | 25 | 100 | 1 | 4.0 |
| | 4 | 24 | 96 | 1 | 4.2 |
| APG-EA Diluent Water | 1 | 23 | 92 | 2 | 8.7 |
| | 2 | 25 | 100 | 2 | 8.0 |
| 10 | 1 | 25 | 100 | 1 | 4.0 |
| | 2 | 24 | 96 | 2 | 8.3 |
| 18 | 1 | 24 | 96 | 1 | 4.2 |
| | 2 | 25 | 100 | 2 | 8.0 |
| 32 | 1 | 23 | 92 | 3 | 13.0 |
| | 2 | 25 | 100 | 2 | 8.0 |
| 56 | 1 | 22 | 88 | 4 | 18.2 |
| | 2 | 23 | 92 | 4 | 17.4 |
| 100 | 1 | 18 | 72 | 9 | 50.0 |
| | 2 | 19 | 76 | 10 | 52.6 |

TABLE A44-3. FETAX BEACH POINT RAW GROUNDWATER TOXICITY TEST
STATISTICAL ANALYSIS (TEST NO. 4) - PERCENT EMBRYO
SURVIVAL AFTER 96 HOURS OF EXPOSURE

Data Transformation:

Arc sine square root

Shapiro-Wilk's Test for Normality:

| | |
|----------------------------|---|
| Calculated test statistic: | 0.91 |
| Alpha value: | 0.01 |
| Critical value: | 0.84 |
| Conclusion: | Fail to reject the null hypothesis that the data are normally distributed |

Bartlett's Test for Homogeneity of Variances:

| | |
|----------------------------|--|
| Calculated test statistic: | 2.92 |
| Alpha value: | 0.01 |
| Critical value: | 16.59 |
| Conclusion: | Fail to reject the null hypothesis that the variances are homogenous |

ANOVA:

| | |
|----------------------------|--|
| Calculated test statistic: | 31.14 |
| Alpha value: | 0.05 |
| Critical value: | 3.37 |
| Conclusion: | Reject the null hypothesis that all groups are equal |

Bonferroni T-Test:

| | |
|----------------------------|--|
| Calculated test statistic: | See Table A44-4 |
| Alpha value: | 0.05 |
| Critical value: | 2.94 |
| Conclusion: | Reject the null hypothesis that all groups are equal |

TABLE A44-4. FETAX BEACH POINT RAW GROUNDWATER TOXICITY TEST
STATISTICAL ANALYSIS (TEST NO. 4) - RESULT'S OF
BONFERRONI'S T-TEST ON EMBRYO MORTALITY AFTER 96
HOURS OF EXPOSURE

| Conc (% by Vol) | No. of Reps | Mean Embryo Mortality (%) ^a | T-Statistic | Significance |
|----------------------------|-------------------|---|-------------|--------------|
| UMD/WREC Control | 4 | 2.0 | | |
| APG-EA Diluent Water | 2 | 4.0 | 0.589 | |
| 10 | 2 | 2.0 | 0.000 | |
| 18 | 2 | 2.0 | 0.000 | |
| 32 | 2 | 4.0 | 0.589 | |
| 56 | 2 | 10.0 | 2.339 | |
| 100 | 2 | 26.0 | 12.218 | * |

^a Values given are actual percent mean mortality rather than arc
sine square root transformed means which were used in the
statistical analysis.

* Significantly different at $\alpha = 0.05$ (Bonferroni critical
value = 2.94).

TABLE A44-5. FETAX BEACH POINT RAW GROUNDWATER TOXICITY TEST
STATISTICAL ANALYSIS (TEST NO. 4) - PERCENT EMBRYO
MALFORMATIONS AFTER 96 HOURS OF EXPOSURE

Data Transformation:

Arc sine square root

Shapiro-Wilk's Test for Normality:

| | |
|----------------------------|---|
| Calculated test statistic: | 0.84 |
| Alpha value: | 0.01 |
| Critical Value: | 0.83 |
| Conclusion: | Fail to reject the null hypothesis that the data are normally distributed |

Bartlett's Test for Homogeneity of Variances:

| | |
|----------------------------|--|
| Calculated test statistic: | 5.69 |
| Alpha value: | 0.01 |
| Critical value: | 15.09 |
| Conclusion: | Fail to reject the null hypothesis that the variances are homogenous |

ANOVA:

| | |
|----------------------------|--|
| Calculated test statistic: | 6.00 |
| Alpha value: | 0.05 |
| Critical value: | 3.69 |
| Conclusion: | Reject the null hypothesis that all groups are equal |

Bonferroni's T-test:

| | |
|----------------------------|--|
| Calculated test statistic: | See Table A44-5 |
| Alpha value: | 0.05 |
| Critical value: | 2.90 |
| Conclusion: | Reject the null hypothesis that all groups are equal |

TABLE A44-6. FETAX BEACH POINT RAW GROUNDWATER TOXICITY TEST DATA (TEST NO. 4) - TYPE AND NUMBER OF MALFORMED EMBRYOS AFTER 96 HOURS OF EXPOSURE.

| Malformation | Test Concentrations (% Groundwater by Volume) | | | | | | | | | | | | | | | |
|--------------|---|---|---|---|------------------|---|------------|---|------------|---|------------|---|------------|---|------------|---|
| | UMD ^a | | | | APG ^b | | 10 | | 18 | | 32 | | 56 | | 100 | |
| | <u>Rep</u> | | | | <u>Rep</u> | | <u>Rep</u> | | <u>Rep</u> | | <u>Rep</u> | | <u>Rep</u> | | <u>Rep</u> | |
| | 1 | 2 | 3 | 4 | 1 | 2 | 1 | 2 | 1 | 2 | 1 | 2 | 1 | 2 | 1 | 2 |
| Severe | | | | | | | | | | | | | 1 | | 1 | |
| Gut, coiling | 1 | 1 | | | 2 | 1 | 1 | 1 | | | | | 1 | | 2 | 2 |
| Edema: | | | | | | | | | | | | | | | | |
| Multiple | 1 | | 1 | 1 | 1 | | 1 | | 1 | 1 | 1 | | 2 | 1 | 4 | 5 |
| Cardiac | | | | | | | | | | | | | | | | 1 |
| Abdominal | | | | | | | | | | | 1 | 2 | 1 | 2 | | 1 |
| Facial | | | | | | | | | | | | | | | | |
| Cephalic | | | | | | | | | | | | | | | | |
| Blisters | | | | | | | | | | | | | | | | |
| Tail | | | | | | | | | | | | | | | | |
| Notochord | | 1 | | | | | | | 1 | | 1 | | | | 1 | 1 |
| Fin | | | | | | | | | | | | | | | | |
| Face | | | | | | | | | | | | | | | | 1 |
| Eye | | | | | | | | | | | | | | | | |
| Brain | | | | | | | | | | | | | | | | |
| Hemorrhage | | | | | | | | | | | | | | | | |
| Cardiac | | | | | | | | | | | | | | | | |
| Other | | | | | | | | | | | | | | | | |

^a UMD/WREC control.

^b APG-EA diluent water.

APPENDIX 45

FROG EMBRYO TERATOGENESIS ASSAY - Xenopus (FETAX) CONDUCTED ON BEACH POINT BUFFERED (pH \approx 7) GROUNDWATER (WELL 33B) (TEST NO. 4)

| | |
|---------------------------|--|
| Test Method: | ASTM Designation E 1439-91 (ASTM, 1991) |
| Type of Test: | Static renewal (every 24 h) |
| Date: | December 3 - 7, 1993 |
| Investigator: | S. D. Turley |
| Laboratory: | UMD/WREC |
| Groundwater: | |
| Source: | APG-EA Beach Point Well 33B |
| Chemical Characteristics: | See Appendix 48 |
| Test Medium: | |
| Source: | FETAX solution |
| pH characteristics: | See Table A45-1 |
| Test Organism: | |
| Scientific Name: | <u>Xenopus laevis</u> |
| Age at Start of Test: | Stage 8 blastula to stage 11 gastrulae |
| Source: | UMD/WREC culture |
| Experimental Chambers: | |
| Material: | Glass petri dishes |
| Test Solution Volume: | 10 mL |
| No. Organisms/Replicate: | 25 |
| No. Organisms/Treatment: | Groundwater: 50 Control: 100 |
| Loading: | n/a |
| Lighting: | Fluorescent; 60-85 foot candles |
| Aeration: | Prior to renewals |

| | |
|-------------------|--------------------------|
| pH Buffer: | 10 N NaOH |
| Endpoints: | Mortality; malformations |
| Test Temperature: | 24 ± 0.2 °C |

Results:

Mortality:

The 100% Beach Point buffered groundwater caused a significant ($\alpha = 0.05$) reduction in embryo survival. 100% Beach Point buffered groundwater only killed 13% of the frog embryos; thus, a 96-h LC50 could not be obtained. The data are summarized in Tables A45-2, A45-3, and A45-4.

Malformations:

Significant ($\alpha = 0.05$) embryo malformations occurred at all treatments down to 32% buffered groundwater by volume (see Tables A45-2, A45-5, and A45-6). The NOEC and LOEC for the embryos, based on increased numbers of malformations, are as follows:

NOEC = 18% buffered groundwater by volume.

LOEC = 32% buffered groundwater by volume.

The types of malformed embryos are given in Table A45-7.

TABLE A45-1. SUMMARY OF THE BEACH POINT BUFFERED GROUNDWATER
BIOASSAY pH (STANDARD UNITS) DATA FOR FETAX (TEST
NO. 4)

| | <u>Test Concentrations (Percent Groundwater by Volume)</u> | | | | | |
|--------------|--|------|------|------|------|------|
| | 0 | 10 | 18 | 32 | 56 | 100 |
| <u>Day 0</u> | | | | | | |
| 0 H | 7.51 | 7.40 | 7.28 | 7.32 | 7.32 | 7.40 |
| <u>Day 1</u> | | | | | | |
| 0 H | 7.50 | 7.44 | 7.33 | 7.28 | 7.19 | 7.11 |
| <u>Day 2</u> | | | | | | |
| 0 H | 7.44 | 7.40 | 7.30 | 7.26 | 7.17 | 7.10 |
| <u>Day 3</u> | | | | | | |
| 0 H | 7.49 | 7.51 | 7.50 | 7.40 | 7.43 | 7.39 |
| <u>Day 4</u> | | | | | | |
| 24 H | 7.58 | 7.47 | 7.40 | 7.35 | 7.30 | 7.27 |

TABLE A45-2. FETAX BEACH POINT BUFFERED GROUNDWATER TOXICITY
TEST DATA (TEST NO. 4) - PERCENT EMBRYO SURVIVAL
AND MALFORMATIONS AFTER 96 HOURS OF EXPOSURE

| Conc (% by Vol) | Rep | Number Embryos Alive | Percent Survival | Number Embryos Malformed | Percent Malformed |
|-----------------------|-----|----------------------------|---------------------|--------------------------------|----------------------|
| UMD/WREC Control | 1 | 25 | 100 | 2 | 8.0 |
| | 2 | 24 | 96 | 2 | 8.3 |
| | 3 | 25 | 100 | 1 | 4.0 |
| | 4 | 24 | 96 | 1 | 4.2 |
| 10 | 1 | 24 | 96 | 2 | 8.3 |
| | 2 | 25 | 100 | 1 | 4.0 |
| 18 | 1 | 23 | 92 | 2 | 8.7 |
| | 2 | 25 | 100 | 2 | 8.0 |
| 32 | 1 | 24 | 96 | 3 | 12.5 |
| | 2 | 23 | 92 | 4 | 17.4 |
| 56 | 1 | 24 | 96 | 5 | 20.8 |
| | 2 | 22 | 88 | 4 | 18.1 |
| 100 | 1 | 20 | 80 | 7 | 20.8 |
| | 2 | 23 | 92 | 7 | 30.4 |

TABLE A45-3. FETAX BEACH POINT BUFFERED GROUNDWATER TOXICITY
TEST STATISTICAL ANALYSIS (TEST NO. 4) - PERCENT
EMBRYO SURVIVAL AFTER 96 HOURS OF EXPOSURE

Data Transformation:

Arc sine square root

Shapiro-Wilk's Test for Normality:

| | |
|----------------------------|---|
| Calculated test statistic: | 0.86 |
| Alpha value: | 0.01 |
| Critical value: | 0.83 |
| Conclusion: | Fail to reject the null hypothesis that the data are normally distributed |

Bartlett's Test for Homogeneity of Variance:

| | |
|----------------------------|--|
| Calculated test statistic: | 1.32 |
| Alpha value: | 0.01 |
| Critical value: | 15.09 |
| Conclusion: | Fail to reject the null hypothesis that all groups are equal |

ANOVA:

| | |
|----------------------------|--|
| Calculated test statistic: | 2.13 |
| Alpha value: | 0.05 |
| Critical value: | 3.69 |
| Conclusion: | Reject the null hypothesis that all groups are equal |

Bonferroni's T-Test

| | |
|----------------------------|--|
| Calculated test statistic: | See Table A45-4 |
| Alpha value: | 0.01 |
| Critical value: | 2.90 |
| Conclusion: | Reject the null hypothesis that all groups are equal |

TABLE A45-4. FETAX BEACH POINT BUFFERED GROUNDWATER TOXICITY
TEST STATISTICAL ANALYSIS (TEST NO. 4) - RESULT'S
OF BONFERRONI'S T-TEST ON EMBRYO MORTALITY AFTER 96
HOURS OF EXPOSURE

| Conc (% by Vol) | No. of Reps | Mean Embryo Mortality (%) ^a | T-Statistic | Significance |
|-----------------------|-------------------|---|-------------|--------------|
| UMD/WREC Control | 4 | 2.0 | | |
| 10 | 2 | 2.0 | 0.000 | |
| 18 | 2 | 4.0 | 0.551 | |
| 32 | 2 | 6.0 | 1.204 | |
| 56 | 2 | 8.0 | 1.637 | |
| 100 | 2 | 14.0 | 2.898 | * |

^a Values given are actual percent mean mortality rather than arc sine square root transformed means which were used in the statistical analysis.

* Significantly different at alpha = 0.05 (Bonferroni's critical value = 2.90).

TABLE A45-5. FETAX BEACH POINT BUFFERED GROUNDWATER TOXICITY
TEST STATISTICAL ANALYSIS (TEST NO. 4) - PERCENT
EMBRYO MALFORMATIONS AFTER 96 HOURS OF EXPOSURE

Data Transformation:

Arc sine square root

Chi-Square Test for Normality:

| | |
|----------------------------|---|
| Calculated test statistic: | 6.09 |
| Alpha value: | 0.01 |
| Critical Value: | 13.28 |
| Conclusion: | Fail to reject the null hypothesis that the data are normally distributed |

Bartlett's Test for Homogeneity of Variances:

| | |
|----------------------------|--|
| Calculated test statistic: | 3.39 |
| Alpha value: | 0.01 |
| Critical value: | 13.28 |
| Conclusion: | Fail to reject the null hypothesis that the variances are homogenous |

ANOVA:

| | |
|----------------------------|--|
| Calculated test statistic: | 9.00 |
| Alpha value: | 0.05 |
| Critical value: | 4.12 |
| Conclusion: | Reject the null hypothesis that all groups are equal |

Bonferroni's T-Test:

| | |
|----------------------------|--|
| Calculated test statistic: | See Table A45-5 |
| Alpha value: | 0.05 |
| Critical value: | 2.84 |
| Conclusion: | Reject the null hypothesis that all groups are equal |

TABLE A45-6. FETAX BEACH POINT BUFFERED GROUNDWATER TOXICITY TEST STATISTICAL ANALYSIS (TEST NO. 4) - RESULTS OF BONFERRONI'S T-TEST ON EMBRYO MALFORMATIONS AFTER 96 HOURS OF EXPOSURE^a

| Conc (% by Vol) | No. of Reps | Mean Normal Embryos (%) ^b | T Statistic | Significance |
|-----------------------|-------------------|---|-------------|--------------|
| UMD/WREC Control | 4 | 92.9 | | |
| 10 | 2 | 93.9 | -0.516 | |
| 18 | 2 | 91.7 | 0.679 | |
| 32 | 2 | 85.1 | 3.332 | * |
| 56 | 2 | 80.6 | 4.899 | * |

^a The 100% Beach Point raw groundwater treatment was not included in the statistical analyses because all concentrations above the NOEC for survival are excluded from the hypothesis test for malformations.

^b Values given are actual percent mean normal embryos rather than arc sine square root transformed means which were used in the statistical analysis.

* Significantly different at alpha = 0.05 (Bonferroni's critical value = 2.84).

TABLE A45-7. FETAX BEACH POINT BUFFERED GROUNDWATER TOXICITY
TEST DATA (TEST NO. 4) - TYPE AND NUMBER OF
MALFORMED EMBRYOS AFTER 96 HOURS OF EXPOSURE

| Malformation | Test Concentrations (% Groundwater by Volume) | | | | | | | | | |
|--------------|---|---|---|---|------------|---|------------|---|------------|---|
| | 0 | | | | 10 | | 18 | | 32 | |
| | <u>Rep</u> | | | | <u>Rep</u> | | <u>Rep</u> | | <u>Rep</u> | |
| | 1 | 2 | 3 | 4 | 1 | 2 | 1 | 2 | 1 | 2 |
| Severe | | | | | | | | | 1 | |
| Gut, coiling | 1 | 1 | | | 2 | | 1 | | | |
| Edema: | | | | | | | | | | |
| Multiple | 1 | | 1 | 1 | 1 | | 2 | 1 | 2 | 1 |
| Cardiac | | | | | | | | | | |
| Abdominal | | | | | | | | | 1 | 2 |
| Facial | | | | | | | | | | |
| Cephalic | | | | | | | | | | |
| Blisters | | | | | | | | | | |
| Tail | | | | | | | | | | |
| Notochord | | | | | | | | | | |
| Fin | | | | | | | | | | |
| Face | | | | | | | | | | |
| Eye | | | | | | | | | | |
| Brain | | | | | | | | | | |
| Hemorrhage | | | | | | | | | | |
| Cardiac | | | | | | | | | | |
| Other | | | | | | | | | | |

APPENDIX 46

SIX-MONTH INTERIM AND NINE-MONTH FINAL JAPANESE MEDAKA
CHRONIC HISTOPATHOLOGY GROWTH DATA

TABLE A46-1. SIX-MONTH INTERIM GROWTH DATA OF JAPANESE MEDAKA DURING THE CHRONIC HISTOPATHOLOGY EXPOSURE STUDY (SACRIFICED ON AUGUST 25, 1993; TEST DAY 182) - DILUENT WATER

| Tank 1 | | | Tank 2 | | | Tank 3 | | | Tank 4 | | |
|----------|-----------------|----------------------|----------|-----------------|----------------------|----------|-----------------|----------------------|----------|-----------------|----------------------|
| Fish No. | Wet Weight (mg) | Standard Length (mm) | Fish No. | Wet Weight (mg) | Standard Length (mm) | Fish No. | Wet Weight (mg) | Standard Length (mm) | Fish No. | Wet Weight (mg) | Standard Length (mm) |
| 1 | 573 | 33 | 1 | 401 | 30 | 1 | 314 | 28 | 1 | 520 | 30 |
| 2 | 423 | 31 | 2 | 349 | 27 | 2 | 416 | 32 | 2 | 367 | 28 |
| 3 | 484 | 32 | 3 | 530 | 34 | 3 | 423 | 33 | 3 | 499 | 31 |
| 4 | 473 | 32 | 4 | 714 | 34 | 4 | 627 | 34 | 4 | 545 | 31 |
| 5 | 271 | 26 | 5 | 330 | 30 | 5 | 463 | 31 | 5 | 529 | 32 |
| 6 | 489 | 31 | 6 | 364 | 30 | 6 | 548 | 31 | 6 | 568 | 32 |
| 7 | 579 | 34 | 7 | 480 | 31 | 7 | 388 | 29 | 7 | 433 | 31 |
| 8 | 495 | 29 | 8 | 508 | 32 | 8 | 323 | 27 | 8 | 557 | 31 |
| 9 | 442 | 32 | 9 | 568 | 32 | 9 | 427 | 32 | 9 | 688 | 33 |
| 10 | 458 | 30 | 10 | 423 | 29 | 10 | 526 | 31 | 10 | 475 | 31 |
| 11 | 528 | 31 | 11 | 496 | 32 | 11 | 308 | 29 | 11 | 542 | 31 |
| 12 | 593 | 32 | 12 | 371 | 28 | 12 | 513 | 30 | 12 | 447 | 29 |
| 13 | 482 | 31 | 13 | 372 | 30 | 13 | 514 | 31 | 13 | 519 | 30 |
| 14 | 612 | 32 | 14 | 495 | 31 | 14 | 517 | 30 | 14 | 504 | 31 |
| 15 | 446 | 32 | 15 | 401 | 29 | 15 | 362 | 29 | 15 | 470 | 30 |
| 16 | 516 | 32 | 16 | 459 | 30 | 16 | 512 | 31 | 16 | 406 | 29 |
| 17 | 425 | 32 | 17 | 413 | 31 | 17 | 617 | 32 | 17 | 384 | 27 |
| 18 | 620 | 34 | 18 | 339 | 27 | 18 | 545 | 31 | 18 | 423 | 32 |
| 19 | 395 | 28 | 19 | 722 | 34 | 19 | 511 | 30 | 19 | 323 | 27 |
| 20 | 354 | 29 | 20 | 614 | 33 | 20 | 525 | 32 | 20 | 589 | 32 |
| | | | | | | | | | | | |
| Mean | 483 | 31 | | 467 | 31 | | 469 | 31 | | 489 | 30 |
| Min | 271 | 26 | | 330 | 27 | | 308 | 27 | | 323 | 27 |
| Max | 620 | 34 | | 722 | 34 | | 627 | 34 | | 688 | 33 |
| S.D. | 86.1 | 1.9 | | 113.3 | 2.1 | | 92.5 | 1.7 | | 83.8 | 1.6 |
| N | 20 | 20 | | 20 | 20 | | 20 | 20 | | 20 | 20 |

TABLE A46-1. (CONTINUED) - 1% GROUNDWATER

| Tank 5 | | | Tank 6 | | | Tank 7 | | | Tank 8 | | |
|----------|-----------------|----------------------|----------|-----------------|----------------------|----------|-----------------|----------------------|----------|-----------------|----------------------|
| Fish No. | Wet Weight (mg) | Standard Length (mm) | Fish No. | Wet Weight (mg) | Standard Length (mm) | Fish No. | Wet Weight (mg) | Standard Length (mm) | Fish No. | Wet Weight (mg) | Standard Length (mm) |
| 1 | 412 | 27 | 1 | 512 | 32 | 1 | 425 | 34 | 1 | 396 | 28 |
| 2 | 283 | 27 | 2 | 677 | 32 | 2 | 433 | 30 | 2 | 457 | 30 |
| 3 | 772 | 32 | 3 | 484 | 29 | 3 | 503 | 30 | 3 | 596 | 21 |
| 4 | 389 | 29 | 4 | 808 | 35 | 4 | 348 | 30 | 4 | 435 | 30 |
| 5 | 349 | 29 | 5 | 472 | 31 | 5 | 523 | 32 | 5 | 573 | 32 |
| 6 | 499 | 31 | 6 | 584 | 32 | 6 | 422 | 28 | 6 | 549 | 31 |
| 7 | 632 | 31 | 7 | 345 | 29 | 7 | 531 | 31 | 7 | 422 | 29 |
| 8 | 474 | 30 | 8 | 263 | 26 | 8 | 377 | 28 | 8 | 434 | 32 |
| 9 | 429 | 28 | 9 | 403 | 28 | 9 | 417 | 30 | 9 | 604 | 32 |
| 10 | 423 | 32 | 10 | 657 | 35 | 10 | 479 | 31 | 10 | 363 | 27 |
| 11 | 640 | 32 | 11 | 478 | 30 | 11 | 388 | 30 | 11 | 669 | 32 |
| 12 | 457 | 30 | 12 | 376 | 29 | 12 | 405 | 30 | 12 | 507 | 32 |
| 13 | 591 | 32 | 13 | 403 | 31 | 13 | 539 | 30 | 13 | 497 | 32 |
| 14 | 508 | 30 | 14 | 605 | 34 | 14 | 602 | 32 | 14 | 563 | 32 |
| 15 | 524 | 30 | 15 | 462 | 32 | 15 | 416 | 31 | 15 | 642 | 32 |
| 16 | 639 | 31 | 16 | 344 | 28 | 16 | 481 | 30 | 16 | 456 | 30 |
| 17 | 473 | 29 | 17 | 383 | 30 | 17 | 601 | 32 | 17 | 506 | 29 |
| 18 | 651 | 31 | 18 | 492 | 30 | 18 | 403 | 29 | 18 | 520 | 32 |
| 19 | 348 | 28 | 19 | 489 | 30 | 19 | 377 | 29 | 19 | 457 | 31 |
| 20 | 506 | 30 | 20 | 430 | 30 | 20 | 293 | 28 | 20 | 415 | 30 |
| 21 | 507 | 31 | | | | | | | | | |
| | | | | | | | | | | | |
| Mean | 500 | 30 | | 483 | 31 | | 448 | 30 | | 503 | 30 |
| Min | 283 | 27 | | 263 | 26 | | 293 | 28 | | 363 | 21 |
| Max | 772 | 32 | | 808 | 35 | | 602 | 34 | | 669 | 32 |
| S.D. | 117.4 | 1.5 | | 127.3 | 2.3 | | 80.0 | 1.5 | | 83.0 | 2.6 |
| N | 21 | 21 | | 20 | 20 | | 20 | 20 | | 20 | 20 |

TABLE A46-1. (CONTINUED) - 10% GROUNDWATER

| Tank 9 | | | Tank 10 | | | Tank 11 | | | Tank 12 | | |
|----------|-----------------|----------------------|----------|-----------------|----------------------|----------|-----------------|----------------------|----------|-----------------|----------------------|
| Fish No. | Wet Weight (mg) | Standard Length (mm) | Fish No. | Wet Weight (mg) | Standard Length (mm) | Fish No. | Wet Weight (mg) | Standard Length (mm) | Fish No. | Wet Weight (mg) | Standard Length (mm) |
| 1 | 467 | 30 | 1 | 607 | 31 | 1 | 364 | 31 | 1 | 565 | 32 |
| 2 | 610 | 32 | 2 | 478 | 30 | 2 | 418 | 31 | 2 | 586 | 30 |
| 3 | 545 | 32 | 3 | 446 | 30 | 3 | 478 | 32 | 3 | 609 | 32 |
| 4 | 392 | 31 | 4 | 493 | 31 | 4 | 560 | 32 | 4 | 427 | 31 |
| 5 | 591 | 33 | 5 | 397 | 30 | 5 | 423 | 30 | 5 | 510 | 31 |
| 6 | 450 | 31 | 6 | 565 | 32 | 6 | 718 | 34 | 6 | 430 | 32 |
| 7 | 490 | 31 | 7 | 564 | 32 | 7 | 294 | 27 | 7 | 462 | 31 |
| 8 | 583 | 31 | 8 | 532 | 31 | 8 | 291 | 29 | 8 | 461 | 30 |
| 9 | 599 | 32 | 9 | 437 | 30 | 9 | 416 | 28 | 9 | 573 | 31 |
| 10 | 555 | 31 | 10 | 510 | 31 | 10 | 555 | 31 | 10 | 466 | 31 |
| 11 | 411 | 29 | 11 | 418 | 30 | 11 | 548 | 33 | 11 | 325 | 27 |
| 12 | 586 | 31 | 12 | 419 | 30 | 12 | 451 | 28 | 12 | 469 | 32 |
| 13 | 587 | 31 | 13 | 593 | 33 | 13 | 680 | 33 | 13 | 387 | 31 |
| 14 | 303 | 27 | 14 | 493 | 31 | 14 | 576 | 30 | 14 | 712 | 34 |
| 15 | 476 | 29 | 15 | 480 | 31 | 15 | 536 | 33 | 15 | 444 | 30 |
| 16 | 541 | 31 | 16 | 338 | 28 | 16 | 578 | 32 | 16 | 380 | 29 |
| 17 | 496 | 31 | 17 | 640 | 33 | 17 | 296 | 28 | 17 | 427 | 31 |
| 18 | 400 | 31 | 18 | 471 | 32 | 18 | 505 | 31 | 18 | 392 | 31 |
| 19 | 573 | 32 | 19 | 548 | 33 | 19 | 444 | 32 | 19 | 437 | 29 |
| 20 | 659 | 33 | 20 | 420 | 29 | 20 | 320 | 29 | 20 | 302 | 29 |
| Mean | 516 | 31 | | 492 | 31 | | 473 | 31 | | 468 | 31 |
| Min | 303 | 27 | | 338 | 28 | | 291 | 27 | | 302 | 27 |
| Max | 659 | 33 | | 640 | 33 | | 718 | 34 | | 712 | 34 |
| S.D. | 88.6 | 1.4 | | 76.0 | 1.3 | | 120.7 | 2.0 | | 97.5 | 1.5 |
| N | 20 | 20 | | 20 | 20 | | 20 | 20 | | 20 | 20 |

TABLE A46-2. JAPANESE MEDAKA BEACH POINT GROUNDWATER CHRONIC
HISTOPATHOLOGY ASSAY STATISTICAL ANALYSIS - WET
WEIGHT AFTER SIX MONTHS OF EXPOSURE

Data Transformation:

None

Kolmogorov-Smirnov Test for Normality:

Data were normally distributed ($P = 0.127$)

Levene Median Test for Homogeneity of Variances:

Variances were homogenous ($P = 0.682$)

General Linear Model Type III Test- Replicates within Treatment
as the Error Term:

| Source | DF | Sum of Squares | Mean Square | F Value | P > F |
|-----------|----|----------------|-------------|---------|-------|
| Treatment | 2 | 4162.5043 | 2081.2521 | 0.25 | 0.781 |

Results:

No difference in wet weight occurred between the Japanese medaka exposed for six months to 10% groundwater by volume, 1% groundwater by volume, and diluent water.

TABLE A46-2. (CONTINUED) - STANDARD LENGTH AFTER SIX MONTHS OF EXPOSURE

Data Transformation:

None

Kolmogorov-Smirnov Test for Normality:

Data were not normally distributed ($P < 0.001$)

Kruskal Wallis One Way ANOVA on Ranks:

| Group | N | Median | H | P |
|---------|----|--------|-------|-------|
| Control | 80 | 31.0 | | |
| 10% | 80 | 30.0 | | |
| 1% | 81 | 30.0 | 4.042 | 0.133 |

Results:

No difference in standard length occurred between the Japanese medaka exposed for six months to 10% groundwater by volume, 1% groundwater by volume, and diluent water.

TABLE A46-3. NINE-MONTH FINAL GROWTH DATA OF JAPANESE MEDAKA DURING THE CHRONIC HISTOPATHOLGY EXPOSURE STUDY (SACRIFICED ON NOVEMBER 29, 1993; TEST DAY 278) - DILUENT WATER

| Tank 1 | | | Tank 2 | | | Tank 3 | | | Tank 4 | | |
|----------|-----------------|----------------------|----------|-----------------|----------------------|----------|-----------------|----------------------|----------|-----------------|----------------------|
| Fish No. | Wet Weight (mg) | Standard Length (mm) | Fish No. | Wet Weight (mg) | Standard Length (mm) | Fish No. | Wet Weight (mg) | Standard Length (mm) | Fish No. | Wet Weight (mg) | Standard Length (mm) |
| 1 | 321 | 30 | 1 | 346 | 28 | 1 | 362 | 27 | 1 | 459 | 29 |
| 2 | 505 | 31 | 2 | 377 | 27 | 2 | 482 | 32 | 2 | 773 | 34 |
| 3 | 524 | 30 | 3 | 661 | 33 | 3 | 521 | 30 | 3 | 521 | 30 |
| 4 | 580 | 31 | 4 | 576 | 32 | 4 | 723 | 35 | 4 | 424 | 29 |
| 5 | 574 | 32 | 5 | 725 | 34 | 5 | 453 | 32 | 5 | 503 | 33 |
| 6 | 782 | 34 | 6 | 686 | 32 | 6 | 569 | 34 | 6 | 702 | 35 |
| 7 | 628 | 33 | 7 | 413 | 31 | 7 | 709 | 36 | 7 | 599 | 30 |
| 8 | 447 | 32 | 8 | 522 | 31 | 8 | 641 | 31 | 8 | 452 | 31 |
| 9 | 525 | 32 | 9 | 610 | 34 | 9 | 424 | 31 | 9 | 777 | 32 |
| 10 | 689 | 34 | 10 | 450 | 28 | 10 | 664 | 32 | 10 | 752 | 35 |
| 11 | 715 | 34 | 11 | 600 | 33 | 11 | 474 | 34 | 11 | 569 | 32 |
| 12 | 680 | 34 | 12 | 303 | 26 | 12 | 464 | 32 | 12 | 798 | 36 |
| 13 | 1044 | 37 | 13 | 592 | 33 | 13 | 365 | 29 | 13 | 393 | 29 |
| 14 | 610 | 32 | 14 | 366 | 27 | 14 | 372 | 30 | 14 | 692 | 36 |
| 15 | 736 | 32 | 15 | 429 | 31 | 15 | 647 | 32 | 15 | 309 | 25 |
| 16 | 456 | 32 | 16 | 668 | 34 | 16 | 536 | 34 | 16 | 459 | 30 |
| 17 | 586 | 32 | 17 | 699 | 32 | 17 | 547 | 32 | 17 | 476 | 31 |
| 18 | 507 | 31 | 18 | 470 | 31 | 18 | 587 | 33 | 18 | 473 | 31 |
| 19 | 656 | 32 | 19 | 797 | 34 | 19 | 626 | 32 | 19 | 477 | 31 |
| 20 | 628 | 32 | 20 | 724 | 35 | 20 | 902 | 36 | 20 | 679 | 34 |
| 21 | 385 | 30 | 21 | 741 | 34 | | | | 21 | 646 | 32 |
| 22 | 481 | 32 | 22 | 806 | 36 | | | | 22 | 496 | 30 |
| 23 | 593 | 32 | 23 | 459 | 31 | | | | 23 | 553 | 32 |
| 24 | 796 | 33 | 24 | 693 | 34 | | | | 24 | 804 | 34 |
| | | | 25 | 778 | 34 | | | | 25 | 736 | 34 |

TABLE A46-3. (CONTINUED) - DILUENT WATER CONT

| Tank 1 | | | Tank 2 | | | Tank 3 | | | Tank 4 | | |
|----------|-----------------|----------------------|----------|-----------------|----------------------|----------|-----------------|----------------------|----------|-----------------|----------------------|
| Fish No. | Wet Weight (mg) | Standard Length (mm) | Fish No. | Wet Weight (mg) | Standard Length (mm) | Fish No. | Wet Weight (mg) | Standard Length (mm) | Fish No. | Wet Weight (mg) | Standard Length (mm) |
| | | | 29 | 502 | 32 | | | | 29 | 480 | 29 |
| | | | 30 | 315 | 26 | | | | 30 | 439 | 32 |
| | | | 31 | 552 | 32 | | | | 31 | 658 | 34 |
| | | | | | | | | | 32 | 480 | 30 |
| | | | | | | | | | 33 | 557 | 32 |
| | | | | | | | | | 34 | 465 | 32 |
| | | | | | | | | | 35 | 695 | 31 |
| <hr/> | | | | | | | | | | | |
| Mean | 602 | 32 | | 573 | 32 | | 553 | 32 | | 565 | 32 |
| Min | 321 | 30 | | 303 | 26 | | 362 | 27 | | 309 | 25 |
| Max | 1044 | 37 | | 806 | 36 | | 902 | 36 | | 804 | 36 |
| S.D. | 148.9 | 1.5 | | 149.4 | 2.6 | | 134.6 | 2.2 | | 129.4 | 2.3 |
| N | 24 | 24 | | 31 | 31 | | 20 | 20 | | 35 | 35 |

TABLE A46-3. (CONTINUED) - 1% GROUNDWATER

| Tank 5 | | | Tank 6 | | | Tank 7 | | | Tank 8 | | |
|----------|-----------------|----------------------|----------|-----------------|----------------------|----------|-----------------|----------------------|----------|-----------------|----------------------|
| Fish No. | Wet Weight (mg) | Standard Length (mm) | Fish No. | Wet Weight (mg) | Standard Length (mm) | Fish No. | Wet Weight (mg) | Standard Length (mm) | Fish No. | Wet Weight (mg) | Standard Length (mm) |
| 1 | 546 | 31 | 1 | 549 | 32 | 1 | 695 | 33 | 1 | 384 | 29 |
| 2 | 350 | 36 | 2 | 572 | 31 | 2 | 578 | 34 | 2 | 346 | 31 |
| 3 | 658 | 31 | 3 | 641 | 34 | 3 | 438 | 32 | 3 | 510 | 32 |
| 4 | 640 | 32 | 4 | 555 | 31 | 4 | 522 | 30 | 4 | 474 | 32 |
| 5 | 683 | 32 | 5 | 828 | 34 | 5 | 563 | 33 | 5 | 575 | 32 |
| 6 | 562 | 31 | 6 | 484 | 32 | 6 | 662 | 32 | 6 | 518 | 30 |
| 7 | 607 | 32 | 7 | 297 | 28 | 7 | 622 | 33 | 7 | 884 | 34 |
| 8 | 706 | 32 | 8 | 548 | 30 | 8 | 587 | 32 | 8 | 626 | 33 |
| 9 | 612 | 31 | 9 | 491 | 30 | 9 | 349 | 30 | 9 | 618 | 32 |
| 10 | 390 | 28 | 10 | 740 | 30 | 10 | 668 | 32 | 10 | 609 | 34 |
| 11 | 630 | 32 | 11 | 830 | 32 | 11 | 553 | 33 | 11 | 625 | 32 |
| 12 | 923 | 35 | 12 | 642 | 33 | 12 | 708 | 35 | 12 | 615 | 33 |
| 13 | 519 | 30 | 13 | 548 | 30 | 13 | 549 | 31 | 13 | 440 | 32 |
| 14 | 468 | 29 | 14 | 487 | 30 | 14 | 784 | 35 | 14 | 473 | 32 |
| 15 | 488 | 30 | 15 | 589 | 34 | 15 | 517 | 31 | 15 | 658 | 33 |
| 16 | 502 | 30 | 16 | 629 | 32 | 16 | 568 | 31 | 16 | 325 | 27 |
| 17 | 464 | 31 | 17 | 654 | 34 | 17 | 393 | 30 | 17 | 678 | 32 |
| 18 | 806 | 34 | 18 | 371 | 30 | 18 | 373 | 29 | 18 | 551 | 30 |
| 19 | 535 | 30 | 19 | 600 | 35 | 19 | 603 | 31 | 19 | 525 | 32 |
| 20 | 773 | 35 | 20 | 416 | 30 | 20 | 806 | 34 | 20 | 527 | 30 |
| 21 | 544 | 32 | 21 | 422 | 28 | 21 | 456 | 31 | 21 | 501 | 32 |
| 22 | 641 | 34 | 22 | 533 | 32 | 22 | 456 | 31 | 22 | 470 | 30 |
| 23 | 704 | 32 | 23 | 604 | 33 | 23 | 588 | 32 | 23 | 405 | 30 |
| 24 | 590 | 32 | 24 | 468 | 30 | 24 | 616 | 32 | 24 | 539 | 31 |
| 25 | 550 | 33 | 25 | 331 | 34 | 25 | 566 | 32 | 25 | 428 | 30 |

TABLE A46-3. (CONTINUED) - 1% GROUNDWATER CONT

| Tank 5 | | | Tank 6 | | | Tank 7 | | | Tank 8 | | |
|----------|-----------------|----------------------|----------|-----------------|----------------------|----------|-----------------|----------------------|----------|-----------------|----------------------|
| Fish No. | Wet Weight (mg) | Standard Length (mm) | Fish No. | Wet Weight (mg) | Standard Length (mm) | Fish No. | Wet Weight (mg) | Standard Length (mm) | Fish No. | Wet Weight (mg) | Standard Length (mm) |
| 26 | 713 | 34 | 26 | 556 | 31 | 26 | 672 | 34 | 26 | 751 | 36 |
| 27 | 700 | 32 | 27 | 514 | 32 | 27 | 715 | 35 | 27 | 549 | 32 |
| 28 | 508 | 31 | 28 | 621 | 34 | 28 | 542 | 31 | 28 | 879 | 36 |
| 29 | 361 | 30 | 29 | 751 | 34 | | | | 29 | 467 | 31 |
| 30 | 520 | 31 | 30 | 431 | 31 | | | | 30 | 484 | 31 |
| 31 | 467 | 29 | 31 | 498 | 30 | | | | 31 | 603 | 30 |
| 32 | 503 | 30 | 32 | 432 | 30 | | | | 32 | 493 | 30 |
| | | | 33 | 749 | 34 | | | | 33 | 458 | 31 |
| | | | 34 | 467 | 30 | | | | 34 | 328 | 30 |
| | | | | | | | | | 35 | 595 | 35 |
| | | | | | | | | | 36 | 290 | 28 |
| | | | | | | | | | 37 | 531 | 30 |
| <hr/> | | | | | | | | | | | |
| Mean | 583 | 32 | | 554 | 32 | | 577 | 32 | | 533 | 31 |
| Min | 350 | 28 | | 297 | 28 | | 349 | 29 | | 290 | 27 |
| Max | 923 | 36 | | 830 | 35 | | 806 | 35 | | 884 | 36 |
| S.D. | 126.5 | 1.8 | | 127.6 | 1.9 | | 113.4 | 1.6 | | 131.5 | 1.9 |
| N | 32 | 32 | | 34 | 34 | | 28 | 28 | | 37 | 37 |

TABLE A46-3. (CONTINUED) - 10% GROUNDWATER

| Tank 9 | | | Tank 10 | | | Tank 11 | | | Tank 12 | | |
|----------|-----------------|----------------------|----------|-----------------|----------------------|----------|-----------------|----------------------|----------|-----------------|----------------------|
| Fish No. | Wet Weight (mg) | Standard Length (mm) | Fish No. | Wet Weight (mg) | Standard Length (mm) | Fish No. | Wet Weight (mg) | Standard Length (mm) | Fish No. | Wet Weight (mg) | Standard Length (mm) |
| 1 | 580 | 32 | 1 | 789 | 36 | 1 | 413 | 31 | 1 | 318 | 29 |
| 2 | 762 | 34 | 2 | 378 | 29 | 2 | 543 | 32 | 2 | 379 | 30 |
| 3 | 736 | 34 | 3 | 390 | 30 | 3 | 508 | 31 | 3 | 608 | 32 |
| 4 | 483 | 31 | 4 | 760 | 36 | 4 | 687 | 35 | 4 | 616 | 32 |
| 5 | 662 | 34 | 5 | 542 | 30 | 5 | 678 | 33 | 5 | 544 | 30 |
| 6 | 782 | 34 | 6 | 451 | 31 | 6 | 757 | 33 | 6 | 489 | 32 |
| 7 | 697 | 34 | 7 | 496 | 35 | 7 | 441 | 31 | 7 | 459 | 34 |
| 8 | 692 | 35 | 8 | 687 | 36 | 8 | 547 | 31 | 8 | 450 | 30 |
| 9 | 691 | 34 | 9 | 552 | 32 | 9 | 691 | 34 | 9 | 659 | 33 |
| 10 | 648 | 30 | 10 | 728 | 36 | 10 | 478 | 31 | 10 | 442 | 30 |
| 11 | 496 | 32 | 11 | 836 | 35 | 11 | 532 | 31 | 11 | 604 | 32 |
| 12 | 655 | 34 | 12 | 458 | 31 | 12 | 393 | 31 | 12 | 582 | 35 |
| 13 | 534 | 30 | 13 | 604 | 33 | 13 | 722 | 34 | 13 | 592 | 35 |
| 14 | 606 | 34 | 14 | 502 | 30 | 14 | 551 | 33 | 14 | 731 | 35 |
| 15 | 582 | 35 | 15 | 872 | 36 | 15 | 845 | 34 | 15 | 453 | 32 |
| 16 | 626 | 34 | 16 | 625 | 32 | 16 | 461 | 31 | 16 | 665 | 36 |
| 17 | 731 | 35 | 17 | 361 | 30 | 17 | 412 | 30 | 17 | 497 | 32 |
| 18 | 691 | 36 | 18 | 682 | 34 | 18 | 535 | 30 | 18 | 347 | 27 |
| 19 | 681 | 32 | 19 | 835 | 35 | 19 | 631 | 32 | 19 | 403 | 30 |
| 20 | 526 | 32 | 20 | 366 | 30 | 20 | 604 | 32 | 20 | 594 | 33 |
| 21 | 550 | 34 | 21 | 522 | 34 | 21 | 472 | 31 | 21 | 825 | 36 |
| 22 | 558 | 32 | 22 | 606 | 34 | 22 | 619 | 34 | 22 | 615 | 33 |
| 23 | 448 | 32 | 23 | 553 | 33 | 23 | 363 | 30 | 23 | 741 | 34 |
| 24 | 777 | 37 | 24 | 432 | 30 | 24 | 462 | 30 | 24 | 468 | 28 |
| 25 | 562 | 31 | 25 | 873 | 36 | 25 | 496 | 31 | 25 | 616 | 33 |

TABLE A46-3. (CONTINUED) - 10% GROUNDWATER CONT

| Tank 9 | | | Tank 10 | | | Tank 11 | | | Tank 12 | | |
|----------|-----------------|----------------------|----------|-----------------|----------------------|----------|-----------------|----------------------|----------|-----------------|----------------------|
| Fish No. | Wet Weight (mg) | Standard Length (mm) | Fish No. | Wet Weight (mg) | Standard Length (mm) | Fish No. | Wet Weight (mg) | Standard Length (mm) | Fish No. | Wet Weight (mg) | Standard Length (mm) |
| 26 | 690 | 34 | 26 | 775 | 34 | 26 | 690 | 35 | 26 | 560 | 31 |
| 27 | 828 | 36 | 27 | 651 | 35 | 27 | 403 | 30 | 27 | 557 | 31 |
| 28 | 469 | 33 | 28 | 743 | 35 | 28 | 368 | 26 | 28 | 531 | 32 |
| 29 | 356 | 30 | 29 | 498 | 33 | 29 | 522 | 32 | 29 | 706 | 26 |
| 30 | 610 | 32 | 30 | 853 | 34 | 30 | 438 | 30 | 30 | 463 | 31 |
| 31 | 661 | 34 | 31 | 693 | 33 | 31 | 642 | 33 | 31 | 689 | 34 |
| 32 | 685 | 35 | 32 | 428 | 30 | 32 | 608 | 34 | 32 | 388 | 31 |
| 33 | 406 | 34 | 33 | 560 | 30 | 33 | 437 | 30 | 33 | 704 | 35 |
| 34 | 471 | 29 | | | | 34 | 585 | 32 | 34 | 508 | 34 |
| 35 | 671 | 34 | | | | 35 | 766 | 34 | 35 | 474 | 30 |
| | | | | | | 36 | 672 | 36 | 36 | 629 | 42 |
| Mean | 617 | 33 | | 609 | 33 | | 555 | 32 | | 553 | 32 |
| Min | 356 | 29 | | 361 | 29 | | 363 | 26 | | 318 | 26 |
| Max | 828 | 37 | | 873 | 36 | | 845 | 36 | | 825 | 42 |
| S.D. | 111.8 | 1.9 | | 158.4 | 2.3 | | 123.0 | 1.9 | | 118.7 | 2.9 |
| N | 35 | 35 | | 33 | 33 | | 36 | 36 | | 36 | 36 |

TABLE A46-4. JAPANESE MEDAKA BEACH POINT GROUNDWATER CHRONIC HISTOPATHOLOGY ASSAY STATISTICAL ANALYSIS - WET WEIGHT AFTER NINE MONTHS OF EXPOSURE

Data Transformation:

None

Kolmogorov-Smirnov Test for Normality:

Data were normally distributed ($P = 0.119$)

Levene Median Test for Homogeneity of Variances:

Variances were homogenous ($P = 0.216$)

General Linear Model Type III Test- Replicates within Treatment as the Error Term:

| Source | DF | Sum of Squares | Mean Square | F Value | P > F |
|-----------|----|----------------|-------------|---------|-------|
| Treatment | 2 | 31435.8821 | 15717.9411 | 0.68 | 0.530 |

Results:

No difference in wet weight occurred between the Japanese medaka exposed for nine months to 10% groundwater by volume, 1% groundwater by volume, and diluent water.

TABLE A46-4. (CONTINUED) - STANDARD LENGTH AFTER NINE MONTHS OF EXPOSURE

Data Transformation:

None

Kolmogorov-Smirnov Test for Normality:

Data were not normally distributed ($P < 0.001$)

Kruskal Wallis One Way ANOVA on Ranks:

| Group | N | Median | H | P |
|---------|-----|--------|--------|-------|
| Control | 110 | 32.0 | | |
| 10% | 140 | 32.5 | | |
| 1% | 131 | 32.0 | 11.389 | 0.003 |

Dunn's Multiple Comparison Test:

| Comparison | Difference of Rank | Q | P |
|-----------------|--------------------|-------|------------|
| 10% vs. control | 28.482 | 2.033 | <0.05 |
| 1% vs. control | 15.990 | | Not Tested |

Results:

The Kruskal Wallis test indicates that a significant difference ($\alpha = 0.003$) in standard length occurred between the Japanese medaka exposed for nine months to 10% groundwater by volume, 1% groundwater by volume, and diluent water. However, when Dunn's multiple comparison test is used to enumerate the difference between the control and each experimental treatment, no statistically significant difference ($\alpha = 0.05$) occurs.

APPENDIX 47

SUMMARY OF THE SIX-MONTH INTERIM AND NINE-MONTH
FINAL JAPANESE MEDAKA CHRONIC
HISTOPATHOLOGY RESULTS

EPL[®]

EXPERIMENTAL PATHOLOGY LABORATORIES, INC.

U.S. ARMY BIOMEDICAL RESEARCH
AND DEVELOPMENT LABORATORY
TEST 401-001R
EPL PROJECT NUMBER 406-027

BEACH POINT GROUNDWATER STUDY WITH MEDAKA

AMENDED PATHOLOGY REPORT

VOLUME 1

Submitted by:

Experimental Pathology Laboratories, Inc.
P.O. Box 474
Herndon, VA 22070
(703) 471-7060

Submitted to:

U.S. Army Biomedical Research
and Development Laboratory
Fort Detrick, MD 21702-5010

August 26, 1994

U.S. ARMY BIOMEDICAL RESEARCH AND DEVELOPMENT LABORATORY
TEST 401-001R
EPL PROJECT NUMBER 406-027

BEACH POINT GROUNDWATER STUDY WITH MEDAKA
AMENDED PATHOLOGY SUMMARY

A histopathological evaluation was performed on tissues from fish of the species *Oryzias latipes* (Medaka) exposed to groundwater at Beach Point, Aberdeen Proving Ground, to indicate the toxic or carcinogenic potential of the groundwater and to determine the need for remediation. Medaka that were 14 days old were either exposed to 10 mg/l diethylnitrosamine (DEN) in water for 48 hours or were held under identical conditions with no DEN present. At 17 days of age the medaka were exposed to 0, 1 or 10% groundwater in the biomonitoring trailer at Beach Point. Laboratory controls were maintained at Ft. Detrick. After six months of exposure approximately one-third of the original 60 fish in each exposure group were euthanized and processed for histopathologic evaluation. The remainder of the fish in each exposure group were euthanized and evaluated after nine months of exposure. The study design is as follows:

| Group No. | Treatment mg/L DEN; % Groundwater | No. of Fish for Evaluation 6 Mo. Sacrifice | No. of Fish for Evaluation 9 Mo. Sacrifice |
|-----------|--------------------------------------|--|--|
| 1 | 0;0 | 20 | 27 |
| 2 | 0;0 | 20 | 34 |
| 3 | 10;0 | 20 | 25 |
| 4 | 10;0 | 20 | 36 |
| 5 | 0;1 | 21 | 36 |
| 6 | 0;1 | 20 | 37 |
| 7 | 10;1 | 20 | 33 |
| 8 | 10;1 | 20 | 38 |
| 9 | 0;10 | 20 | 37 |
| 10 | 0;10 | 20 | 34 |
| 11 | 10;10 | 20 | 33 |
| 12 | 10;10 | 20 | 40 |
| 17 | 0;Lab control | 20 | 38 |
| 18 | 0;Lab control | 20 | 38 |
| 19 | 10;Lab control | 20 | 35 |
| 20 | 10;Lab control | 20 | 34 |

Hematoxylin and eosin stained slides of each fish were prepared by Experimental Pathology Laboratories, Inc. Most fish were sampled by cutting five step sections through the fish in a longitudinal plane. In other fish, transverse sections were cut in certain areas (e.g. the head) in order to capture gross lesions that were described. In certain fish the lesions were sectioned separately. The following tissues were evaluated: bone (vertebra), brain, chromaffin tissue, corpuscle of Stannius, esophagus, eye, gallbladder, gill, heart,

hematopoietic tissue, interrenal tissue, intestine, kidney, liver, nares, ovary, pancreas, peripheral nerve, pineal organ, pituitary gland, pseudobranch, skeletal muscle, skin, spinal cord, spleen, stato-acoustic organ, swim bladder, testis, thymus, thyroid tissue, urinary bladder and gross lesions.

Tissues that were not present for examination in one or more medaka included bone (vertebra), chromaffin tissue, corpuscle of Stannius, eye, gallbladder, heart, interrenal tissue, nares, pineal organ, pituitary, spinal cord, spleen, thymus and urinary bladder. Occasional absence of these tissues is a condition inherent in the sectioning method.

Microscopic findings for each tissue examined from each medaka are listed in the Histopathology Incidence Tables by group and by sex. Inflammatory, degenerative and hyperplastic changes were graded from 1 to 5 depending upon severity. Neoplasms, cysts, developmental anomalies and other nongradable changes were designated as present (P). Tissues that were autolyzed are indicated with an "A," with or without additional diagnosis depending on the severity of the autolysis. Tissues of insufficient quantity for evaluation are indicated with an "I." All lesions are summarized by treatment group in the Incidence/Examined Summary by sex and with sexes combined with incidence expressed as numerator/denominator and percentage. Summary Incidence Tables by group and by disposition and sex, and by disposition with sexes combined, separate medaka which are part of a scheduled sacrifice from

those which died early. All neoplasms are also listed in the Neoplasm Summary Incidence Tables by group and by sex. A correlation of gross observations with the corresponding microscopic findings is presented in the Correlation of Gross and Microscopic Findings tables.

Medaka identified as Nos. 93-327-4-31 and 93-327-4-32 were not found among the fish submitted for histopathologic evaluation and are considered missing. Medaka from replicate Groups 1 and 2 probably intermingled and from replicate Groups 3 and 4 probably intermingled during a water backup in the biomonitoring trailer. Groups 1 and 2 did not intermingle with Groups 3 and 4.

RESULTS

6-Month Interim Sacrifice

At six months, liver neoplasms (hepatocellular adenoma and carcinoma) occurred only among medaka in the biomonitoring trailer that were pre-exposed to DEN. Most neoplasms occurred in medaka that were also exposed to 10% groundwater. Hepatocellular carcinomas occurred only in medaka exposed to DEN and 10% groundwater. No liver neoplasia was present among the laboratory controls. Foci of cellular alteration occurred in groups exposed to DEN and in groups not exposed to DEN, but the incidence was higher in groups exposed to DEN. See Table I for a tabulation of liver neoplasia and foci of cellular alteration in medaka after six months of exposure to groundwater. The replicate groups are combined in this synopsis.

U.S. Army Biomedical Research and Development Laboratory Test 401-001R

TABLE I

INCIDENCE OF LIVER NEOPLASIA AND FOCI OF CELLULAR ALTERATION
IN MEDAKA AFTER SIX MONTHS OF EXPOSURE TO GROUNDWATER

| Exposure: DEN | No DEN | DEN | No DEN | DEN | No DEN | DEN | No DEN | DEN |
|---------------------------|--------|-------|--------|-------|--------|---------|----------|----------|
| Groundwater | 0% | 0% | 1% | 1% | 10% | 10% | None | None |
| Group Number | 1 & 2 | 3 & 4 | 5 & 6 | 7 & 8 | 9 & 10 | 11 & 12 | 17 & 18* | 19 & 20* |
| LIVER (Total No. Fish) | (40) | (40) | (41) | (40) | (40) | (40) | (40) | (40) |
| Hepatocellular Adenoma | - | 2 | - | 1 | - | 4 | - | - |
| Hepatocellular Carcinoma | - | - | - | - | - | 3 | - | - |
| Basophilic Focus (Foci) | - | 1 | 1 | 2 | - | 1 | - | - |
| Eosinophilic Focus (Foci) | - | 2 | - | 5 | - | 3 | 1 | 3 |
| Eosinophilic Area | 1 | - | 1 | - | - | - | - | - |

*Groups 17, 18, 19 and 20 are laboratory controls.

Neoplasms unrelated to treatment that occurred in individual medaka included a lymphosarcoma in the thymus of a Group 3 medaka, a seminoma in the testis of a Group 1 medaka and a seminoma in the ovary of a Group 10 medaka.

Granulomatous inflammation of the subcutaneous tissue of the lower jaw and/or upper jaw which was unrelated to exposure occurred only in male medaka. One to three medaka out of 20 or 21 total fish were affected in each of Groups 1, 2, 3, 4, 5, 6, 9 and 11, but not in any of the laboratory control groups.

Granulomas and/or granulomatous inflammation occurred sporadically among males and females in a variety of other tissues including adipose tissue of the head and cranial cavity, choroid rete of the eye, heart, hematopoietic tissue, liver, mesentery, oral cavity

submucosa, ovary, pericardial cavity, pseudobranch, skeletal muscle and urinary bladder. These lesions were not related to exposure.

Increased basophilia of thyroid tissue was diagnosed in medaka in all groups, but the incidence was greater among males than among females. The change was characterized by basophilic cuboidal follicular cells that form numerous small follicles with a small amount of colloid in contrast to follicles that are variable in size, distended with colloid and lined by flattened follicular cells. The reason for this basophilic appearance of the thyroid tissue, especially among male medaka, is not known.

Other changes occurred in a variety of tissues in medaka of both control and exposed groups, but these lesions either were present in small numbers of medaka or were distributed among the control and exposed groups such that there was no apparent relationship to exposure.

9-Month Final Sacrifice

At nine months, liver neoplasms [hepatocellular adenoma and carcinoma; carcinoma, not otherwise specified (NOS); and cholangioma] occurred in medaka pre-exposed to DEN that were subsequently exposed to 0, 1 and 10% groundwater and in two Group 9 medaka exposed to 10% groundwater and no DEN. The occurrence of hepatocellular adenomas and carcinomas is neither dose related nor exposure related among the medaka pre-exposed to DEN and housed in the biomonitoring trailer. Among the control medaka held at the Fort Detrick laboratory, 3 of 69 fish pre-exposed to DEN had hepatocellular adenomas. Foci of cellular alteration

occurred in groups exposed to DEN and in groups not exposed to DEN, but the incidence was higher in groups exposed to DEN. See Table II for a tabulation of liver neoplasia and foci of cellular alteration in medaka after nine months of exposure to groundwater. The replicate groups are combined in this synopsis.

TABLE II

INCIDENCE OF LIVER NEOPLASIA AND FOCI OF CELLULAR ALTERATION
IN MEDAKA AFTER NINE MONTHS OF EXPOSURE TO GROUNDWATER

| Exposure: DEN | No DEN | DEN | No DEN | DEN | No DEN | DEN | No DEN | DEN |
|---------------------------------------|--------|-------|--------|-------|--------|---------|----------|----------|
| Groundwater | 0% | 0% | 1% | 1% | 10% | 10% | None | None |
| Group Number | 1 & 2 | 3 & 4 | 5 & 6 | 7 & 8 | 9 & 10 | 11 & 12 | 17 & 18* | 19 & 20* |
| LIVER (Total No.)** | (61) | (61) | (73) | (71) | (71) | (73) | (76) | (69) |
| Carcinoma, NOS | - | - | - | - | - | 1 | - | - |
| Hepatocellular Adenoma | - | 4 | - | 4 | 1 | 5 | - | 3 |
| Hepatocellular Adenoma, Multiple | - | 1 | - | 1 | - | 1 | - | - |
| Hepatocellular Carcinoma | - | 2 | - | - | 1 | 2 | - | - |
| Hepatocellular Carcinoma, Multiple | - | 1 | - | - | - | - | - | - |
| Cholangioma | - | - | - | - | - | 1 | - | - |
| Basophilic Area | - | - | - | 1 | - | 1 | - | - |
| Basophilic Focus (Foci) | - | 3 | 1 | 3 | 1 | 5 | - | 1 |
| Clear Cell Focus | - | - | - | - | 1 | - | - | - |
| Eosinophilic Focus (Foci) | - | 4 | - | 4 | 1 | 4 | 1 | 5 |

*Groups 17, 18, 19 and 20 are laboratory controls.

**Total number of tissues evaluated includes those from medaka that died early or were sacrificed at nine months.

U.S. Army Biomedical Research and Development Laboratory Test 401-001R

Neoplasms unrelated to treatment that occurred in individual medaka included lymphosarcoma in a single medaka of Groups 1, 3, 11 and 18 and in two medaka from Group 7; a thyroid follicular cell adenoma in a Group 5 medaka; a thyroid follicular cell carcinoma in a Group 9 medaka; a testicular seminoma in a Group 12 medaka; a hemangioma in the gills of a Group 17 medaka; a gallbladder papilloma in a Group 17 medaka and an ovarian seminoma in a Group 20 medaka.

Granulomatous inflammation of the lower and/or upper jaw occurred among medaka in all exposure groups, and was more commonly seen in male medaka than female medaka. The lesion was often described grossly as an enlarged or thickened lower or upper jaw or lip. Associated with the inflammation in some cases was an increase in the amount of connective tissue in the area in the form of cartilage, bone or fibrous connective tissue. A number of medaka, but not all, with granulomatous inflammation of the jaw also had granulomas or granulomatous inflammation in one or more other tissues such as the choroid rete of the eye, heart, hematopoietic tissue, liver, mesentery, oral cavity, pseudobranch and spleen. Other medaka had granulomatous lesions of one or more tissues, but no jaw lesions.

Tissues from two Group 2 male medaka (2-25 and 2-34), one Group 3 female medaka (3-50), one Group 18 male medaka (18-34) and one Group 20 female medaka (20-52) were stained with an acid-fast stain. In the Group 2 medaka, acid-fast positive organisms were present in the

U.S. Army Biomedical Research and Development Laboratory Test 401-001R

granulomatous lesions of the jaws as well as in other tissues with granulomatous lesions. In the Group 18 medaka, acid-fast organisms were present in the granulomas that were described only in the hematopoietic tissue. In the Group 3 and Group 20 medaka, acid-fast organisms were found in several tissues including gills, hematopoietic tissue, liver and mesentery.

Granulomas in the heart, hematopoietic tissue, liver and spleen occurred in medaka in almost all groups housed in the biomonitoring trailer. Among the laboratory controls, granulomas in one or more of these tissues were found only in Groups 18, 19 and 20 and among these three groups, the highest incidence was in Group 20 medaka, e.g., 15 of 34 had granulomas in the hematopoietic tissue and/or liver). With a few exceptions in granulomas of the heart, liver and spleen, and without exception in the hematopoietic tissue, the incidence of these granulomas was higher among male medaka than among female medaka. Among the groups housed in the trailer, the highest incidence of granulomas in these four tissues occurred among the control groups exposed to 0% groundwater, and the lowest incidence occurred among the groups exposed to 10% groundwater. See Table III for a tabulation of the incidence of these granulomas in the medaka housed in the biomonitoring trailer and the laboratory with the incidence from the replicate groups combined and males and females combined.

U.S. Army Biomedical Research and Development Laboratory Test 401-001R

TABLE III

INCIDENCE OF GRANULOMAS IN SELECTED TISSUES FROM MEDAKA
AFTER NINE MONTHS OF EXPOSURE TO GROUNDWATER

| Exposure: DEN | No DEN | DEN | No DEN | DEN | No DEN | DEN | No DEN | DEN |
|---------------------------------------|-----------|-------|-----------|-------|-----------|---------|-----------|----------|
| Groundwater | 0% | 0% | 1% | 1% | 10% | 10% | None | None |
| Group Number | 1 & 2 | 3 & 4 | 5 & 6 | 7 & 8 | 9 & 10 | 11 & 12 | 17 & 18* | 19 & 20* |
| HEART (Total No.)** | (61) | (61) | (73) | (69) | (71) | (73) | (76) | (69) |
| Granulomas | 12 | 22 | 9 | 9 | 5 | 4 | 0 | 6 |
| HEMATOPOIETIC TISSUE (Total No.)** | (61) | (61) | (73) | (71) | (71) | (73) | (76) | (69) |
| Granulomas | 29 | 28 | 17 | 14 | 7 | 7 | 1 | 16 |
| LIVER (Total No.)** | (61) | (61) | (73) | (71) | (71) | (73) | (76) | (69) |
| Granulomas | 37 | 31 | 24 | 22 | 10 | 13 | 0 | 16 |
| SPLEEN (Total No.)** | (60) | (56) | (64) | (67) | (63) | (62) | (65) | (61) |
| Granulomas | 12 | 12 | 11 | 5 | 5 | 2 | 0 | 8 |

*Groups 17, 18, 19 and 20 are laboratory controls.

**Total number of tissues evaluated includes those from medaka that died early or were sacrificed at nine months.

Of the 35 medaka that died early and were evaluated histologically, nine were females and 26 were males. None were from the laboratory control groups. They consisted of one or more fish from all exposure groups in the biomonitoring trailer except for Group 11. None of these 35 fish had liver neoplasms. All of the males that died early except one had granulomatous lesions of two or more tissues. In male medaka No. 1-34 the cause of death was not apparent. Male medaka No. 7-30 had lymphosarcoma and No. 9-23 had a thyroid follicular cell carcinoma in addition to granulomatous lesions. Of the nine females that died early, three had granulomatous lesions of numerous tissues. Medaka No. 2-27 had a granulomatous lesion of the caudal fin.

Lymphosarcoma occurred in medaka Nos. 1-27 and 7-26. Medaka Nos. 5-24 and 7-29 had egg impaction, and in medaka No. 8-23 moderately severe degenerative ova may have caused the bloated condition described before the fish was euthanized.

Cystic degeneration (spongiosis hepatis) in the liver occurred in all exposure groups and was more common among females than males. The incidence in the laboratory controls overall (29 of 145 total fish) was less than in the groups in the biomonitoring trailer exposed to 0% (54 of 122), 1% (58 of 144) or 10% (46 of 144) groundwater. The severity varied from minimal to moderately severe, but was minimal to mild in the majority of the medaka with the lesion.

Hepatic cysts occurred in the liver in all exposure groups, but the incidence in the laboratory controls overall was less than in the groups in the biomonitoring trailer.

Vacuolated hepatocyte foci in the liver occurred in all exposure groups, but the incidence was less in controls (18 of 145 laboratory controls, 12 of 122 trailer controls) than in medaka exposed to 1% (32 of 144) or 10% (26 of 144) groundwater.

Fusion of gill lamellae occurred in male and female medaka in all the exposure groups and was more common among females than males. There was a slight increase in incidence of the gill change overall in the medaka exposed to 10% groundwater (Groups 9, 10, 11, 12); however, the severity of the change, which was usually minimal to mild, was similar among all the exposure groups.

Increased basophilia of thyroid tissue, characterized as stated in the discussion of the medaka from the six-month interim sacrifice, was diagnosed in medaka from all exposure groups. The incidence was approximately three times greater among males than among females but in almost all cases the severity was minimal to mild.

Other changes occurred in a variety of tissues in medaka of both control and exposed groups, but these lesions either were present in small numbers of medaka or were distributed among the groups such that there was no apparent relationship to exposure.

CONCLUSIONS

After six months of exposure to 0, 1 or 10% groundwater, liver neoplasia (hepatocellular adenoma and carcinoma) occurred in medaka that had been previously initiated with DEN. Seven of 40 medaka were affected in the groups exposed to 10% groundwater as compared to two of 40 among the controls and one of 40 medaka exposed to 1% groundwater. There were no liver neoplasms among the laboratory controls.

After nine months, hepatocellular adenomas or carcinomas, single or multiple, occurred in DEN-initiated medaka that had been exposed to 0, 1 and 10% groundwaters such that there were as many neoplasms among the controls (8 of 122) as among the 10% exposure group (8 of 144). There were two liver neoplasms (one hepatocellular adenoma and one carcinoma) in medaka exposed to 10% groundwater, but no DEN. Two additional liver neoplasms not related to hepatocytes occurred in

the DEN-10% groundwater exposure Group 12. One was a cholangioma and the other was a carcinoma, cell of origin unknown. In the DEN-1% groundwater exposure groups there were five (5 of 144) hepatocellular neoplasms, all adenomas. At nine months there was no dose-related response for liver neoplasia in medaka exposed to groundwater. There were three hepatocellular adenomas diagnosed among the DEN-initiated laboratory controls.

Other neoplasms that occurred sporadically among the various exposure groups at six months and nine months included lymphosarcoma, seminoma, thyroid follicular cell adenoma, thyroid follicular cell carcinoma, hemangioma of the gills and a gallbladder papilloma.

Granulomatous inflammation of the subcutaneous tissue of the lower and/or upper jaw occurred in small numbers of male medaka in the biomonitoring trailer (one to three medaka in each group except Groups 7, 8, 10 and 12) at six months. The same lesion occurred among medaka of all exposure groups, including laboratory controls, at nine months. The incidence was substantially greater among males than among females; for example, 144 males overall versus 17 females were diagnosed with granulomatous inflammation of the lower jaw. Granulomatous inflammation or granulomas occurred in a variety of additional tissues among medaka of all exposure groups. Acid-fast staining of five representative medaka was positive for the presence of acid-fast organisms in granulomatous lesions. The reason for the higher incidence of these lesions in males versus females is not known. Trauma may be an

U.S. Army Biomedical Research and Development Laboratory Test 401-001R

initiating factor in the granulomatous lesions of the jaws, but the reason why the jaws of these medaka should be traumatized is not known.

Granulomas in the heart, hematopoietic tissue, liver and spleen occurred in the highest incidence in controls in the biomonitoring trailer and in the lowest incidence in medaka exposed to 10% groundwater. See Table III. The reasons for this lower incidence among medaka exposed to 10% groundwater as compared to controls is not known.

Marilyn J. Wolfe
MARILYN J. WOLFE, D.V.M., Ph.D.
Pathologist

August 26, 1994

MJW/adu

APPENDIX 48

COMPREHENSIVE CHEMICAL ANALYSES CONDUCTED ON RAW
(pH \approx 4) BEACH POINT GROUNDWATER (WELL CC-33B)
AND CHRONIC HISTOPATHOLOGY EXPOSURE TANKS

TABLE A48-1. RESULTS OF THE APG-EA BEACH POINT GROUNDWATER AND DILUENT WATER
COMPREHENSIVE CHEMICAL ANALYSES (TEST NO. 1) - GENERAL WATER QUALITY^a

| Parameter | Method | Detection Limits | 100% G.W. | 10% G.W. | 1% G.W. | Diluent H ₂ O |
|---------------------------------------|-----------|---------------------|--------------|-------------|------------|-----------------------------|
| Total Alkalinity (CaCO ₃) | EPA 310.1 | 5 | b | 21 | 28 | 26 |
| Ammonia (N) | EPA 350.1 | 0.02 | 0.029 | 0.025 | 0.029 | 0.027 |
| Chloride (Cl) | EPA 300.0 | 0.5 | 490 | 86.7 | 31.2 | 21.9 |
| Total Cyanide (CN) | EPA 335.2 | 10 | ND | ND | ND | ND |
| Fluoride (F) | EPA 300.0 | 0.2 | 1.2 | 0.66 | 0.55 | 0.84 |
| Hardness, (CaCO ₃) | SM2540B | | 170 | 73 | 62 | 59 |
| Nitrate (N) | EPA 300.0 | 0.1 | 0.14 | 1.5 | 1.7 | 4.3 |
| Nitrite (N) | b | b | b | b | b | b |
| Total Phosphate (P) | EPA 365.4 | 0.01 | 2.64 | 3.13 | 0.26 | 0.54 |
| Total Suspended Solids | EPA 160.2 | 5 | ND | ND | ND | ND |
| Specific Conductance | EPA 120.1 | | 1600 | 265 | 175 | 169 |
| Sulfate (SO ₄) | EPA 300.0 | 0.5 | 91.2 | 34.2 | 29.3 | 28.1 |
| Sulfide (S) | EPA 376.1 | 5 | 180 | 350 | 360 | 330 |
| TOC | EPA 415.2 | 1 | 4 | 2 | ND | ND |
| Bromide (Br) | EPA 300.0 | 0.1 | 1.5 | 0.16 | ND | ND |

^a All results expressed as mg/L except for specific conductance which is expressed as $\mu\text{mhos/cm}$.

^b Analysis not conducted by Vendor.

TABLE A48-1. (CONTINUED) - METALS^a

| Parameter | Method | Detection Limits | 100% G.W. | 10% G.W. | 1% G.W. | Diluent H ₂ O |
|-----------------|-----------|------------------|-----------|----------|---------|--------------------------|
| Aluminum (Al) | EPA 200.7 | 40 | 810 | 64 | ND | ND |
| Antimony (Sb) | EPA 200.7 | 20 | ND | ND | ND | ND |
| Arsenic (As) | EPA 206.2 | 5 | ND | ND | ND | ND |
| Beryllium (Be) | EPA 200.7 | 5 | ND | ND | ND | ND |
| Boron (B) | EPA 200.7 | 50 | 54 | 51 | ND | ND |
| Cadmium (Cd) | EPA 213.2 | 0.5 | 0.76 | ND | ND | ND |
| Calcium (Ca) | EPA 200.7 | 500 | 16500 | 17300 | 17200 | 16700 |
| Chromium (Cr) | EPA 200.7 | 5 | ND | ND | ND | ND |
| Cobalt (Co) | EPA 200.7 | 40 | 110 | ND | ND | ND |
| Copper (Cu) | EPA 200.7 | 5 | 5.4 | ND | ND | ND |
| Iron (Fe) | EPA 200.7 | 100 | 4600 | 340 | ND | ND |
| Lead (Pb) | EPA 239.2 | 5 | ND | ND | ND | ND |
| Magnesium (Mg) | EPA 200.7 | 500 | 31300 | 7290 | 4590 | 4230 |
| Manganese (Mn) | EPA 200.7 | 15 | 1860 | 200 | 22 | ND |
| Mercury (Hg) | EPA 245.1 | 0.5 | ND | ND | ND | ND |
| Molybdenum (Mo) | EPA 200.7 | 40 | ND | ND | ND | ND |
| Nickel (Ni) | EPA 200.7 | 8 | 120 | 13 | ND | ND |
| Potassium (K) | EPA 258.1 | 100 | 4130 | 2050 | 1760 | 1710 |
| Selenium (Se) | EPA 270.2 | 5 | ND | ND | ND | ND |
| Silver (Ag) | EPA 272.2 | 0.5 | ND | ND | ND | ND |
| Sodium (Na) | EPA 200.7 | 500 | 257000 | 39200 | 12400 | 9260 |
| Thallium (Tl) | EPA 279.2 | 5 | ND | ND | ND | ND |
| Tin (Sn) | EPA 200.7 | 100 | ND | ND | ND | ND |
| Zinc (Zn) | EPA 200.7 | 5 | 210 | 170 | 160 | 160 |

^a Results expressed as µg/L.

TABLE A48-1. (CONTINUED) - PURGEABLE ORGANIC COMPOUNDS^a

| Parameter | Method | Detection Limits | 100% G.W. | 10% G.W. | 1% G.W. | Diluent H ₂ O |
|-----------------------------|-----------|------------------|-----------|----------|---------|--------------------------|
| Benzene | EPA 524.2 | 1 | ND | ND | ND | ND |
| Bromobenzene | EPA 524.2 | 1 | ND | ND | ND | ND |
| Bromochloromethane | EPA 524.2 | 1 | ND | ND | ND | ND |
| Bromodichloromethane | EPA 524.2 | 1 | ND | ND | ND | ND |
| Bromoform | EPA 524.2 | 1 | ND | ND | ND | ND |
| Bromomethane | EPA 524.2 | 1 | ND | ND | ND | ND |
| Carbon tetrachloride | EPA 524.2 | 1 | ND | ND | ND | ND |
| Chlorobenzene | EPA 524.2 | 1 | 2 | ND | ND | ND |
| Chloroethane | EPA 524.2 | 1 | ND | ND | ND | ND |
| Chloroform | EPA 524.2 | 1 | 10 | 1 | ND | ND |
| Chloromethane | EPA 524.2 | 1 | ND | ND | ND | ND |
| 2-Chlorotoluene | EPA 524.2 | 1 | ND | ND | ND | ND |
| 4-Chlorotoluene | EPA 524.2 | 1 | ND | ND | ND | ND |
| Dibromochloromethane | EPA 524.2 | 1 | ND | ND | ND | ND |
| 1,2-Dibromo-3-chloropropane | EPA 524.2 | 1 | ND | ND | ND | ND |
| 1,2-Dibromoethane | EPA 524.2 | 1 | ND | ND | ND | ND |
| Dibromomethane | EPA 524.2 | 1 | ND | ND | ND | ND |
| 1,2-Dichlorobenzene | EPA 524.2 | 1 | ND | ND | ND | ND |
| 1,3-Dichlorobenzene | EPA 524.2 | 1 | ND | ND | ND | ND |
| 1,4-Dichlorobenzene | EPA 524.2 | 1 | ND | ND | ND | ND |
| Dichlorodifluoromethane | EPA 524.2 | 1 | ND | ND | ND | ND |
| 1,1-Dichloroethane | EPA 524.2 | 1 | ND | ND | ND | ND |
| 1,2-Dichloroethane | EPA 524.2 | 1 | ND | ND | ND | ND |
| 1,1-Dichloroethene | EPA 524.2 | 1 | 1 | ND | ND | ND |
| cis-1,2-Dichloroethene | EPA 524.2 | 1 | 430 | 27 | ND | ND |
| trans-1,2-Dichloroethene | EPA 524.2 | 1 | ND | ND | ND | ND |
| 1,2-Dichloropropane | EPA 524.2 | 1 | ND | ND | ND | ND |

^a Results expressed as µg/L.

TABLE A48-1. (CONTINUED) - PURGEABLE ORGANIC COMPOUNDS CON'T^a

| Parameter | Method | Detection Limits | 100% G.W. | 10% G.W. | 1% G.W. | Diluent H ₂ O |
|---------------------------|-----------|------------------|-----------|----------|---------|--------------------------|
| 1,3-Dichloropropane | EPA 524.2 | 1 | ND | ND | ND | ND |
| 2,2-Dichloropropane | EPA 524.2 | 1 | ND | ND | ND | ND |
| 1,1-Dichloropropene | EPA 524.2 | 1 | ND | ND | ND | ND |
| cis-1,3-Dichloropropene | EPA 524.2 | 1 | ND | ND | ND | ND |
| trans-1,3-Dichloropropene | EPA 524.2 | 1 | ND | ND | ND | ND |
| Ethylbenzene | EPA 524.2 | 1 | ND | ND | ND | ND |
| Methylene chloride | EPA 524.2 | 1 | ND | ND | ND | ND |
| Styrene | EPA 524.2 | 1 | ND | ND | ND | ND |
| 1,1,1,2-Tetrachloroethane | EPA 524.2 | 1 | 3 | ND | ND | ND |
| 1,1,2,2-Tetrachloroethane | EPA 524.2 | 1 | 10000 | 1400 | ND | 7 |
| Tetrachloroethene | EPA 524.2 | 1 | 90 | 6 | ND | ND |
| Toluene | EPA 524.2 | 1 | ND | ND | ND | ND |
| 1,1,1-Trichloroethane | EPA 524.2 | 1 | ND | ND | ND | ND |
| 1,1,2-Trichloroethane | EPA 524.2 | 1 | 90 | 8 | ND | 1 |
| Trichloroethene | EPA 524.2 | 1 | 1400 | 80 | ND | ND |
| Trichlorofluoromethane | EPA 524.2 | 1 | ND | ND | ND | ND |
| 1,2,3-Trichloropropane | EPA 524.2 | 1 | ND | ND | ND | ND |
| Vinyl chloride | EPA 524.2 | 1 | 2 | ND | ND | ND |
| o-Xylene | EPA 524.2 | 1 | ND | ND | ND | ND |
| m-Xylene | EPA 524.2 | 1 | ND | ND | ND | ND |
| p-Xylene | EPA 524.2 | 1 | ND | ND | ND | ND |

^a Results expressed as µg/L.

TABLE A48-1. (CONTINUED) - ORGANIC COMPOUNDS^a

| Parameter | Method | Detection Limits | 100% G.W. | 10% G.W. | 1% G.W. | Diluent H ₂ O |
|-------------------------------|---------|------------------|-----------|----------|---------|--------------------------|
| Acenaphthene | EPA 525 | 10 | ND | ND | ND | ND |
| Acenaphthylene | EPA 525 | 10 | ND | ND | ND | ND |
| Anthracene | EPA 525 | 10 | ND | ND | ND | ND |
| Benzidine | EPA 525 | 20 | ND | ND | ND | ND |
| Benzo[a]anthracene | EPA 525 | 10 | ND | ND | ND | ND |
| Benzo[b]fluoranthene | EPA 525 | 10 | ND | ND | ND | ND |
| Benzo[k]fluoranthene | EPA 525 | 10 | ND | ND | ND | ND |
| Benzo[a]pyrene | EPA 525 | 10 | ND | ND | ND | ND |
| Benzo[g,h,i]perylene | EPA 525 | 10 | ND | ND | ND | ND |
| Butyl Benzyl Phthalate | EPA 525 | 10 | ND | ND | ND | ND |
| Bis (2-chloroethyl) Ether | EPA 525 | 10 | ND | ND | ND | ND |
| Bis (2-chloroethoxy) Methane | EPA 525 | 10 | ND | ND | ND | ND |
| Bis (2-ethylhexyl) Phthalate | EPA 525 | 10 | ND | ND | ND | ND |
| Bis (2-chloroisopropyl) Ether | EPA 525 | 10 | ND | ND | ND | ND |
| 4-Bromophenyl Phenyl Ether | EPA 525 | 10 | ND | ND | ND | ND |
| 2-Chloronaphthalene | EPA 525 | 10 | ND | ND | ND | ND |
| 4-Chlorophenyl Phenyl Ether | EPA 525 | 10 | ND | ND | ND | ND |
| Chrysene | EPA 525 | 10 | ND | ND | ND | ND |
| Dibenzo (a,h) Anthracene | EPA 525 | 10 | ND | ND | ND | ND |
| Di-n-Butylphthalate | EPA 525 | 10 | ND | ND | ND | ND |
| 1,3-Dichlorobenzene | EPA 525 | 10 | ND | ND | ND | ND |
| 1,2-Dichlorobenzene | EPA 525 | 10 | ND | ND | ND | ND |
| 1,4-Dichlorobenzene | EPA 525 | 10 | ND | ND | ND | ND |
| 3,3'-Dichlorobenzidine | EPA 525 | 20 | ND | ND | ND | ND |
| 1,2-Diphenylhydrazine | EPA 525 | 20 | ND | ND | ND | ND |
| Diethyl Phthalate | EPA 525 | 10 | ND | ND | ND | ND |
| Dimethyl Phthalate | EPA 525 | 10 | ND | ND | ND | ND |
| 2,4-Dinitrotoluene | EPA 525 | 10 | ND | ND | ND | ND |
| 2,6-Dinitrotoluene | EPA 525 | 10 | ND | ND | ND | ND |

^a Results expressed as µg/L.

TABLE A48-1. (CONTINUED) - ORGANIC COMPOUNDS CON'T^a

| Parameter | Method | Detection Limits | 100% G.W. | 10% G.W. | 1% G.W. | Diluent H ₂ O |
|----------------------------|---------|------------------|-----------|----------|---------|--------------------------|
| Di-n-octylphthalate | EPA 525 | 10 | ND | ND | ND | ND |
| Fluoranthene | EPA 525 | 10 | ND | ND | ND | ND |
| Fluorene | EPA 525 | 10 | ND | ND | ND | ND |
| Hexachlorobenzene | EPA 525 | 10 | ND | ND | ND | ND |
| Hexachlorobutadiene | EPA 525 | 10 | ND | ND | ND | ND |
| Hexachloroethane | EPA 525 | 10 | ND | ND | ND | ND |
| Hexachlorocyclopentadiene | EPA 525 | 10 | ND | ND | ND | ND |
| Indeno[1,2,3,c,d]pyrene | EPA 525 | 10 | ND | ND | ND | ND |
| Isophorone | EPA 525 | 10 | ND | ND | ND | ND |
| Naphthalene | EPA 525 | 10 | ND | ND | ND | ND |
| Nitrobenzene | EPA 525 | 10 | ND | ND | ND | ND |
| N-Nitrosodi-n-propylamine | EPA 525 | 10 | ND | ND | ND | ND |
| N-Nitrosodimethylamine | EPA 525 | 10 | ND | ND | ND | ND |
| N-Nitrosodiphenylamine | EPA 525 | 10 | ND | ND | ND | ND |
| Phenanthrene | EPA 525 | 10 | ND | ND | ND | ND |
| Pyrene | EPA 525 | 10 | ND | ND | ND | ND |
| 1,2,4-Trichlorobenzene | EPA 525 | 10 | ND | ND | ND | ND |
| 4-Chloro-3-methylphenol | EPA 525 | 10 | ND | ND | ND | ND |
| 2-Chlorophenol | EPA 525 | 10 | ND | ND | ND | ND |
| 2,4-Dichlorophenol | EPA 525 | 10 | ND | ND | ND | ND |
| 2,4-Dimethylphenol | EPA 525 | 10 | ND | ND | ND | ND |
| 2,4-Dinitrophenol | EPA 525 | 50 | ND | ND | ND | ND |
| 2-Methyl-4,6-dinitrophenol | EPA 525 | 50 | ND | ND | ND | ND |
| 2-Nitrophenol | EPA 525 | 10 | ND | ND | ND | ND |
| 4-Nitrophenol | EPA 525 | 50 | ND | ND | ND | ND |
| Pentachlorophenol | EPA 525 | 50 | ND | ND | ND | ND |
| Phenol | EPA 525 | 10 | ND | ND | ND | ND |
| 2,4,6-Trichlorophenol | EPA 525 | 10 | ND | ND | ND | ND |

^a Results expressed as µg/L.

TABLE A48-1. (CONTINUED) - CHLORINATED PESTICIDES^a

| Parameter | Method | Detection Limits | 100% G.W. | 10% G.W. | 1% G.W. | Diluent H ₂ O |
|--------------------|---------|------------------|-----------|----------|---------|--------------------------|
| Aldrin | EPA 508 | 0.01 | ND | ND | ND | ND |
| Chlordane-alpha | EPA 508 | 0.01 | ND | ND | ND | ND |
| Chlordane-gamma | EPA 508 | 0.01 | ND | ND | ND | ND |
| Chloroneb | EPA 508 | 0.01 | ND | ND | ND | ND |
| Chlorobenzilate(a) | EPA 508 | 0.01 | ND | ND | ND | ND |
| Chlorothalonil | EPA 508 | 0.01 | ND | ND | ND | ND |
| DCPA | EPA 508 | 0.01 | ND | ND | ND | ND |
| 4,4'-DDD | EPA 508 | 0.01 | ND | ND | ND | ND |
| 4,4'-DDE | EPA 508 | 0.01 | ND | ND | ND | ND |
| 4,4'-DDT | EPA 508 | 0.01 | ND | ND | ND | ND |
| Dieldrin | EPA 508 | 0.01 | ND | ND | ND | ND |
| Endosulfan I | EPA 508 | 0.01 | ND | ND | ND | ND |
| Endosulfan II | EPA 508 | 0.01 | ND | ND | ND | ND |
| Endosulfan sulfate | EPA 508 | 0.01 | ND | ND | ND | ND |
| Endrin | EPA 508 | 0.01 | ND | ND | ND | ND |
| Endrin aldehyde | EPA 508 | 0.01 | ND | ND | ND | ND |
| Etridiazole | EPA 508 | 0.01 | ND | ND | ND | ND |
| HCH-alpha | EPA 508 | 0.01 | ND | ND | ND | ND |
| HCH-beta | EPA 508 | 0.01 | ND | ND | ND | ND |
| HCH-delta(a) | EPA 508 | 0.01 | ND | ND | ND | ND |
| HCH-gamma(Lindane) | EPA 508 | 0.01 | ND | ND | ND | ND |
| Heptachlor | EPA 508 | 0.01 | ND | ND | ND | ND |
| Heptachlor epoxide | EPA 508 | 0.01 | ND | ND | ND | ND |
| Hexachlorobenzene | EPA 508 | 0.01 | ND | ND | ND | ND |
| Methoxychlor | EPA 508 | 0.01 | ND | ND | ND | ND |
| cis-Permethrin | EPA 508 | 0.01 | ND | ND | ND | ND |
| trans-Permethrin | EPA 508 | 0.01 | ND | ND | ND | ND |
| Propachlor | EPA 508 | 0.01 | ND | ND | ND | ND |

^a Results expressed as µg/L.

TABLE A48-1. (CONTINUED) - CHLORINATED PESTICIDES CON'T^a

| Parameter | Method | Detection Limits | 100% G.W. | 10% G.W. | 1% G.W. | Diluent H ₂ O |
|--------------|---------|------------------|-----------|----------|---------|--------------------------|
| Trifluralin | EPA 508 | 0.01 | ND | ND | ND | ND |
| Toxaphene | EPA 508 | 0.5 | ND | ND | ND | ND |
| Chlordane | EPA 508 | 0.08 | ND | ND | ND | ND |
| Aroclor 1016 | EPA 508 | 0.1 | ND | ND | ND | ND |
| Aroclor 1221 | EPA 508 | 0.1 | ND | ND | ND | ND |
| Aroclor 1232 | EPA 508 | 0.1 | ND | ND | ND | ND |
| Aroclor 1242 | EPA 508 | 0.1 | ND | ND | ND | ND |
| Aroclor 1248 | EPA 508 | 0.1 | ND | ND | ND | ND |
| Aroclor 1254 | EPA 508 | 0.1 | ND | ND | ND | ND |
| Aroclor 1260 | EPA 508 | 0.1 | ND | ND | ND | ND |

^a Results expressed as µg/L.

TABLE A48-1. (CONTINUED) - ORGANOPHOSPHORUS PESTICIDES AND CHLORINATED ACIDS^a

| Parameter | Method | Detection Limits | 100% G.W. | 10% G.W. | 1% G.W. | Diluent H ₂ O |
|-------------------------------------|-----------|------------------|-----------|----------|---------|--------------------------|
| <u>Organophosphorus Pesticides:</u> | | | | | | |
| Malathion | EPA 614 | 0.2 | ND | ND | ND | ND |
| Parathion | EPA 614 | 0.2 | ND | ND | ND | ND |
| Dursban | EPA 614 | 0.2 | ND | ND | ND | ND |
| <u>Chlorinated Acids:</u> | | | | | | |
| Acifluorfen | EPA 515.1 | 0.05 | ND | ND | ND | ND |
| Bentazon | EPA 515.1 | 0.05 | ND | ND | ND | ND |
| Chloramben | EPA 515.1 | 0.05 | ND | ND | ND | ND |
| 2,4-D | EPA 515.1 | 0.05 | ND | ND | ND | ND |
| Dalapon | EPA 515.1 | 0.05 | ND | ND | ND | ND |
| 2,4-DB | EPA 515.1 | 0.05 | ND | ND | ND | ND |
| Dicamba | EPA 515.1 | 0.05 | ND | ND | ND | ND |
| 3,5-Dichlorobenzoic acid | EPA 515.1 | 0.05 | ND | ND | ND | ND |
| Dichlorprop | EPA 515.1 | 0.05 | ND | ND | ND | ND |
| Dinoseb | EPA 515.1 | 0.05 | ND | ND | ND | ND |
| 5-Hydroxydicamba | EPA 515.1 | 0.05 | ND | ND | ND | ND |
| 4-Nitrophenol | EPA 515.1 | 0.05 | ND | ND | ND | ND |
| Pentachlorophenol (PCP) | EPA 515.1 | 0.05 | ND | ND | ND | ND |
| Picloram | EPA 515.1 | 0.05 | ND | ND | ND | ND |
| 2,4,5-T | EPA 515.1 | 0.05 | ND | ND | ND | ND |
| Methyl 2,4,5-TP | EPA 515.1 | 0.05 | ND | ND | ND | ND |
| DCPA acid metabolites(a) | EPA 515.1 | 0.05 | ND | ND | ND | ND |

^a Results expressed as µg/L.

TABLE A48-1. (CONTINUED) - NITROGEN AND PHOSPHOROUS-CONTAINING PESTICIDES^a

| Parameter | Method | Detection Limits | 100% G.W. | 10% G.W. | 1% G.W. | Diluent H ₂ O |
|--------------------|---------|------------------|-----------|----------|---------|--------------------------|
| Alachlor | EPA 507 | 0.2 | ND | ND | ND | ND |
| Ametryn | EPA 507 | 0.25 | ND | ND | ND | ND |
| Atraton | EPA 507 | 0.2 | ND | ND | ND | ND |
| Atrazine | EPA 507 | 0.25 | ND | ND | ND | ND |
| Bromacil | EPA 507 | 0.2 | ND | ND | ND | ND |
| Butachlor | EPA 507 | 0.2 | ND | ND | ND | ND |
| Butylate | EPA 507 | 0.2 | ND | ND | ND | ND |
| Carboxin | EPA 507 | 0.2 | ND | ND | ND | ND |
| Chlorpropham | EPA 507 | 0.2 | ND | ND | ND | ND |
| Cycloate | EPA 507 | 0.25 | ND | ND | ND | ND |
| Diazinon(a) | EPA 507 | 0.2 | ND | ND | ND | ND |
| Dichlorvos | EPA 507 | 0.2 | ND | ND | ND | ND |
| Diphenamid | EPA 507 | 0.25 | ND | ND | ND | ND |
| Disulfoton | EPA 507 | 0.25 | ND | ND | ND | ND |
| Disulfoton sulfone | EPA 507 | 0.2 | ND | ND | ND | ND |
| EPTC | EPA 507 | 0.25 | ND | ND | ND | ND |
| Ethoprop | EPA 507 | 0.25 | ND | ND | ND | ND |
| Fenamiphos | EPA 507 | 0.25 | ND | ND | ND | ND |
| Fenarimol | EPA 507 | 0.2 | ND | ND | ND | ND |
| Fluridone | EPA 507 | 1.0 | ND | ND | ND | ND |
| Hexazinone | EPA 507 | 0.2 | ND | ND | ND | ND |
| Merphos | EPA 507 | 0.25 | ND | ND | ND | ND |
| Metolachlor | EPA 507 | 0.2 | ND | ND | ND | ND |
| Metribuzin | EPA 507 | 0.2 | ND | ND | ND | ND |
| Mevinphos | EPA 507 | 0.25 | ND | ND | ND | ND |
| MGK 264 | EPA 507 | 0.2 | ND | ND | ND | ND |
| Molinate | EPA 507 | 0.2 | ND | ND | ND | ND |

^a Results expressed as µg/L.

TABLE A48-1. (CONTINUED) - NITROGEN AND PHOSPHOROUS-CONTAINING PESTICIDES CON'T^a

| Parameter | Method | Detection Limits | 100% G.W. | 10% G.W. | 1% G.W. | Diluent H ₂ O |
|--------------|---------|------------------|-----------|----------|---------|--------------------------|
| Napropamide | EPA 507 | 0.2 | ND | ND | ND | ND |
| Norflurazon | EPA 507 | 0.2 | ND | ND | ND | ND |
| Pebulate | EPA 507 | 0.2 | ND | ND | ND | ND |
| Prometon | EPA 507 | 0.25 | ND | ND | ND | ND |
| Prometryn | EPA 507 | 0.25 | ND | ND | ND | ND |
| Pronamide(a) | EPA 507 | 0.2 | ND | ND | ND | ND |
| Propazine | EPA 507 | 0.25 | ND | ND | ND | ND |
| Simazine | EPA 507 | 0.25 | ND | ND | ND | ND |
| Simetryn | EPA 507 | 0.2 | ND | ND | ND | ND |
| Stirofos | EPA 507 | 0.2 | ND | ND | ND | ND |
| Tebuthiuron | EPA 507 | 0.2 | ND | ND | ND | ND |
| Terbacil | EPA 507 | 0.2 | ND | ND | ND | ND |
| Terbufos(a) | EPA 507 | 0.2 | ND | ND | ND | ND |
| Terbutryn | EPA 507 | 0.25 | ND | ND | ND | ND |
| Triademefon | EPA 507 | 0.25 | ND | ND | ND | ND |
| Tricyclazole | EPA 507 | 1.0 | ND | ND | ND | ND |
| Vernolate | EPA 507 | 0.2 | ND | ND | ND | ND |

^a Results expressed as µg/L.

TABLE A48-2. RESULTS OF THE APG-EA BEACH POINT GROUNDWATER AND DILUENT WATER

COMPREHENSIVE CHEMICAL ANALYSES (TEST NO. 2) - GENERAL WATER QUALITY^a

| Parameter | Method | Detection Limits | 100% G.W. | 10% G.W. | 1% G.W. | Diluent H ₂ O |
|---------------------------------------|---------------------------|------------------|-----------|----------|---------|--------------------------|
| Total Alkalinity (CaCO ₃) | EPA 310.1 | 1 | ND | 25 | 30 | 31 |
| Ammonia (N) | SM 4500-NH ₃ E | 0.1 | ND | ND | ND | ND |
| Chloride (Cl) | EPA 325.3 | 1 | 580 | 68 | 26 | 21 |
| Total Cyanide (CN) | EPA 335.3 | 0.01 | 0.06 | ND | ND | ND |
| Fluoride (F) | EPA 340.2 | 0.05 | 0.36 | 0.88 | 0.94 | 0.94 |
| Hardness, (CaCO ₃) | EPA 130.2 | 1 | 220 | 80 | 70 | 74 |
| Nitrate (N) | EPA 353.2 | 0.1 | 2.5 | 3.2 | 2.3 | 2.5 |
| Nitrite (N) | EPA 354.1 | 0.01 | ND | ND | ND | ND |
| Total Phosphorus (P) | EPA 365.2 | 0.01 | 0.12 | 0.52 | 0.52 | 0.50 |
| Total Suspended Solids | EPA 160.2 | 5 | ND | ND | ND | ND |
| Specific Conductance | EPA 120.1 | N/A | 1760 | 356 | 213 | 200 |
| Sulfate (SO ₄) | EPA 375.4 | 1 | 62 | 17 | 12 | 12 |
| Sulfide (S) | EPA 376.1 | 1 | ND | ND | ND | ND |
| TOC | EPA 415.1 | 1 | 4 | ND | ND | ND |
| Bromide (Br) | SM 4500-Br | b | 1.5 | 0.44 | 0.44 | 0.37 |

^a All results expressed as mg/L except for specific conductance which is expressed as $\mu\text{mohs/cm}$.

^b The detection limit was 0.5 mg/L for the 100% groundwater sample and 0.1 mg/L for the 10% and 1% groundwater and dilution water samples.

TABLE A48-2. (CONTINUED) - METALS^a

| Parameter | Method | Detection Limits | 100% G.W. | 10% G.W. | 1% G.W. | Diluent H ₂ O |
|-----------------|-----------|------------------|-----------|----------|---------|--------------------------|
| Aluminum (Al) | EPA 200.7 | 0.1 | 1.1 | ND | ND | ND |
| Antimony (Sb) | EPA 204.2 | 0.005 | ND | ND | ND | ND |
| Arsenic (As) | EPA 206.2 | 0.005 | ND | ND | ND | ND |
| Beryllium (Be) | EPA 200.7 | 0.005 | ND | ND | ND | ND |
| Boron (B) | EPA 200.7 | 0.05 | ND | ND | ND | ND |
| Cadmium (Cd) | EPA 213.2 | 0.0005 | 0.0008 | 0.0006 | 0.0007 | 0.0008 |
| Calcium (Ca) | EPA 200.7 | 0.5 | 17 | 20 | 17 | 17 |
| Chromium (Cr) | EPA 218.2 | 0.002 | ND | ND | ND | ND |
| Cobalt (Co) | EPA 200.7 | 0.05 | 0.09 | ND | ND | ND |
| Copper (Cu) | EPA 220.2 | 0.005 | 0.008 | 0.006 | ND | ND |
| Iron (Fe) | EPA 200.7 | 0.1 | 5.4 | 0.5 | 0.1 | ND |
| Lead (Pb) | EPA 239.2 | 0.005 | ND | ND | ND | ND |
| Magnesium (Mg) | EPA 200.7 | 0.5 | 30 | 8.2 | 5.1 | 4.8 |
| Manganese (Mn) | EPA 200.7 | 0.01 | 1.7 | 0.20 | 0.02 | ND |
| Mercury (Hg) | EPA 245.1 | 0.0005 | ND | ND | ND | ND |
| Molybdenum (Mo) | EPA 200.7 | 0.01 | ND | ND | ND | ND |
| Nickel (Ni) | EPA 200.7 | 0.05 | 0.12 | ND | ND | ND |
| Potassium (K) | EPA 200.7 | 0.5 | 2.5 | 2.0 | 1.8 | 1.7 |
| Selenium (Se) | EPA 270.2 | 0.005 | ND | ND | ND | ND |
| Silver (Ag) | EPA 272.2 | 0.001 | ND | ND | ND | ND |
| Sodium (Na) | EPA 200.7 | 0.5 | 260 | 36 | 12 | 9.3 |
| Thallium (Tl) | EPA 279.2 | 0.005 | ND | ND | ND | ND |
| Tin (Sn) | EPA 282.2 | 0.005 | ND | ND | ND | ND |
| Zinc (Zn) | EPA 200.7 | 0.05 | 0.24 | 0.16 | 0.13 | 0.12 |

^a Results expressed as mg/L.

TABLE A48-2. (CONTINUED) - VOLATILES^a

| Parameter | Method | Detection Limits | 100% G.W. | 10% G.W. | 1% G.W. | Diluent H ₂ O |
|---------------------------|----------|------------------|--------------|----------|--------------|--------------------------|
| Chloromethane | EPA 8240 | 10 | ND | ND | ND | ND |
| Bromomethane | EPA 8240 | 10 | ND | ND | ND | ND |
| Vinyl chloride | EPA 8240 | 10 | ND | ND | ND | ND |
| Chloroethane | EPA 8240 | 10 | ND | ND | ND | ND |
| Methylene chloride | EPA 8240 | 5 | ND | ND | ND | ND |
| Acrolein | EPA 8240 | 100 | ND | ND | ND | ND |
| Acrylonitrile | EPA 8240 | 100 | ND | ND | ND | ND |
| Trichlorofluoromethane | EPA 8240 | 5 | ND | ND | ND | ND |
| 1,1-Dichloroethane | EPA 8240 | 5 | ND | ND | ND | ND |
| 1,2-Dichloroethene | EPA 8240 | 5 | 640 | 33 | ^b | ND |
| Chloroform | EPA 8240 | 5 | 11 | ND | ND | ND |
| 1,2-Dichloroethane | EPA 8240 | 5 | ND | ND | ND | ND |
| 1,1,1-Trichloroethane | EPA 8240 | 5 | ND | ND | ND | ND |
| Carbon tetrachloride | EPA 8240 | 5 | ND | ND | ND | ND |
| Bromodichloromethane | EPA 8240 | 5 | ND | ND | ND | ND |
| 1,2-Dichloropropane | EPA 8240 | 5 | ND | ND | ND | ND |
| 1,3-Dichloropropene | EPA 8240 | 5 | ND | ND | ND | ND |
| Dibromochloromethane | EPA 8240 | 5 | ND | ND | ND | ND |
| 1,1,2-Trichloroethane | EPA 8240 | 5 | 110 | 8 | ND | ND |
| 2-Chloroethylvinyl ether | EPA 8240 | 10 | ND | ND | ND | ND |
| Bromoform | EPA 8240 | 5 | ND | ND | ND | ND |
| Tetrachloroethene | EPA 8240 | 5 | 90 | 5 | ND | ND |
| 1,1,2,2-Tetrachloroethane | EPA 8240 | 5 | 17000 | 1300 | 160 | 9 |
| Ethylbenzene | EPA 8240 | 5 | ND | ND | ND | ND |
| 1,1-Dichloroethene | EPA 8240 | 5 | ND | ND | ND | ND |
| Trichloroethene | EPA 8240 | 5 | 1600 | 80 | 7 | ND |
| Benzene | EPA 8240 | 5 | ND | ND | ND | ND |
| Toluene | EPA 8240 | 5 | ND | ND | ND | ND |
| Chlorobenzene | EPA 8240 | 5 | ^b | ND | ND | ND |

^a Results expressed as µg/L.^b Detected below quantitation level.

TABLE A48-2. (CONTINUED) - BASE/NEUTRAL COMPOUNDS^a

| Parameter | Method | Detection Limits | 100% G.W. | 10% G.W. | 1% G.W. | Diluent H ₂ O |
|-----------------------------|----------|------------------|-----------|----------|---------|--------------------------|
| Acenaphthene | EPA 8270 | 10 | ND | ND | ND | ND |
| 2,4-Dinitrotoluene | EPA 8270 | 10 | ND | ND | ND | ND |
| 2,6-Dinitrotoluene | EPA 8270 | 10 | ND | ND | ND | ND |
| Diethyl phthalate | EPA 8270 | 10 | ND | ND | ND | ND |
| 4-Chlorophenyl phenyl ether | EPA 8270 | 10 | ND | ND | ND | ND |
| Fluorene | EPA 8270 | 10 | ND | ND | ND | ND |
| N-Nitrosodiphenylamine | EPA 8270 | 10 | ND | ND | ND | ND |
| 1,2-Diphenylhydrazine | EPA 8270 | 10 | ND | ND | ND | ND |
| 4-Bromophenyl phenyl ether | EPA 8270 | 10 | ND | ND | ND | ND |
| Hexachlorobenzene | EPA 8270 | 10 | ND | ND | ND | ND |
| Phenanthrene | EPA 8270 | 10 | ND | ND | ND | ND |
| Anthracene | EPA 8270 | 10 | ND | ND | ND | ND |
| Di-n-butyl phthalate | EPA 8270 | 10 | ND | ND | ND | ND |
| Fluoranthene | EPA 8270 | 10 | ND | ND | ND | ND |
| Pyrene | EPA 8270 | 10 | ND | ND | ND | ND |
| Benzidine | EPA 8270 | 50 | ND | ND | ND | ND |
| Butylbenzyl phthalate | EPA 8270 | 10 | ND | ND | ND | ND |
| 3,3'-Dichlorobenzidine | EPA 8270 | 20 | ND | ND | ND | ND |
| Benzo(a)anthracene | EPA 8270 | 10 | ND | ND | ND | ND |
| bis(2-Ethylhexyl) phthalate | EPA 8270 | 10 | ND | ND | ND | ND |
| Chrysene | EPA 8270 | 10 | ND | ND | ND | ND |
| Di-n-octyl phthalate | EPA 8270 | 10 | ND | ND | ND | ND |
| Benzo(b)fluoranthene | EPA 8270 | 10 | ND | ND | ND | ND |
| Benzo(k)fluoranthene | EPA 8270 | 10 | ND | ND | ND | ND |
| Benzo(a)pyrene | EPA 8270 | 10 | ND | ND | ND | ND |
| Indeno(1,2,3-cd)pyrene | EPA 8270 | 10 | ND | ND | ND | ND |
| Dibenz(a,h)anthracene | EPA 8270 | 10 | ND | ND | ND | ND |
| Benzo(g,h,i)perylene | EPA 8270 | 10 | ND | ND | ND | ND |

^a Results expressed as µg/L.

TABLE A48-2. (CONTINUED) - BASE/NEUTRAL COMPOUNDS CON'T^a

| Parameter | Method | Detection Limits | 100% G.W. | 10% G.W. | 1% G.W. | Diluent H ₂ O |
|------------------------------|----------|------------------|-----------|----------|---------|--------------------------|
| N-Nitrosodimethylamine | EPA 8270 | 10 | ND | ND | ND | ND |
| bis(2-Chloroethyl) ether | EPA 8270 | 10 | ND | ND | ND | ND |
| 1,3-Dichlorobenzene | EPA 8270 | 10 | ND | ND | ND | ND |
| 1,4-Dichlorobenzene | EPA 8270 | 10 | ND | ND | ND | ND |
| 1,2-Dichlorobenzene | EPA 8270 | 10 | ND | ND | ND | ND |
| bis(2-Chloroisopropyl) ether | EPA 8270 | 10 | ND | ND | ND | ND |
| N-Nitrosodi-n-propylamine | EPA 8270 | 10 | ND | ND | ND | ND |
| Hexachloroethane | EPA 8270 | 10 | ND | ND | ND | ND |
| Nitrobenzene | EPA 8270 | 10 | ND | ND | ND | ND |
| Isophorone | EPA 8270 | 10 | ND | ND | ND | ND |
| bis(2-Chloroethoxy)methane | EPA 8270 | 10 | ND | ND | ND | ND |
| 1,2,4-Trichlorobenzene | EPA 8270 | 10 | ND | ND | ND | ND |
| Naphthalene | EPA 8270 | 10 | ND | ND | ND | ND |
| Hexachlorobutadiene | EPA 8270 | 10 | ND | ND | ND | ND |
| Hexachlorocyclopentadiene | EPA 8270 | 10 | ND | ND | ND | ND |
| 2-Chloronaphthalene | EPA 8270 | 10 | ND | ND | ND | ND |
| Dimethyl phthalate | EPA 8270 | 10 | ND | ND | ND | ND |
| Acenaphthylene | EPA 8270 | 10 | ND | ND | ND | ND |

^a Results expressed as µg/L.

TABLE A48-2. (CONTINUED) - PESTICIDES^a

| Parameter | Method | Detection Limits | 100% G.W. | 10% G.W. | 1% G.W. | Diluent H ₂ O |
|--------------------|----------|------------------|-----------|----------|---------|--------------------------|
| a-BHC | EPA 8080 | 0.05 | ND | ND | ND | ND |
| b-BHC | EPA 8080 | 0.05 | ND | ND | ND | ND |
| g-BHC | EPA 8080 | 0.05 | ND | ND | ND | ND |
| d-BHC | EPA 8080 | 0.05 | ND | ND | ND | ND |
| Heptachlor | EPA 8080 | 0.05 | ND | ND | ND | ND |
| Aldrin | EPA 8080 | 0.05 | ND | ND | ND | ND |
| Heptachlor epoxide | EPA 8080 | 0.05 | ND | ND | ND | ND |
| a-Endosulfan | EPA 8080 | 0.05 | ND | ND | ND | ND |
| 4,4'-DDE | EPA 8080 | 0.1 | ND | ND | ND | ND |
| Dieldrin | EPA 8080 | 0.1 | ND | ND | ND | ND |
| Endrin | EPA 8080 | 0.1 | ND | ND | ND | ND |
| b-Endosulfan | EPA 8080 | 0.3 | ND | ND | ND | ND |
| Endrin aldehyde | EPA 8080 | 0.3 | ND | ND | ND | ND |
| 4,4'-DDD | EPA 8080 | 0.3 | ND | ND | ND | ND |
| Endosulfan sulfate | EPA 8080 | 0.3 | ND | ND | ND | ND |
| 4,4'-DDT | EPA 8080 | 0.3 | ND | ND | ND | ND |
| PCB-1242 | EPA 8080 | 1 | ND | ND | ND | ND |
| PCB-1254 | EPA 8080 | 1 | ND | ND | ND | ND |
| PCB-1221 | EPA 8080 | 1 | ND | ND | ND | ND |
| PCB-1232 | EPA 8080 | 1 | ND | ND | ND | ND |
| PCB-1248 | EPA 8080 | 1 | ND | ND | ND | ND |
| PCB-1260 | EPA 8080 | 1 | ND | ND | ND | ND |
| PCB-1016 | EPA 8080 | 1 | ND | ND | ND | ND |
| Chlordane | EPA 8080 | 1 | ND | ND | ND | ND |
| Toxaphene | EPA 8080 | 3 | ND | ND | ND | ND |

^a Results expressed as µg/L.

TABLE A48-2. (CONTINUED) - HERBICIDES AND ORGANOPHOSPHORUS PESTICIDES^a

| Parameter | Method | Detection Limits | 100% G.W. | 10% G.W. | 1% G.W. | Diluent |
|-------------------------------------|----------|------------------|-----------|----------|---------|---------|
| <u>Herbicides:</u> | | | | | | |
| 2,4-D | EPA 8150 | 0.5 | ND | ND | ND | ND |
| 2,4,5-TP (Silvex) | EPA 8150 | 0.2 | ND | ND | ND | ND |
| <u>Organophosphorus Pesticides:</u> | | | | | | |
| Malathion | EPA 8080 | 0.5 | ND | ND | ND | ND |
| Methyl azinphos | EPA 8080 | 3 | ND | ND | ND | ND |
| Coumaphos | EPA 8080 | 2 | ND | ND | ND | ND |
| Chlorpyrifos | EPA 8080 | 0.1 | ND | ND | ND | ND |
| Diazinon | EPA 8080 | 0.5 | ND | ND | ND | ND |
| Fenthion | EPA 8080 | 50 | ND | ND | ND | ND |
| Naled | EPA 8080 | 5 | ND | ND | ND | ND |
| Methyl parathion | EPA 8080 | 0.5 | ND | ND | ND | ND |
| Parathion | EPA 8080 | 0.5 | ND | ND | ND | ND |

^a Results expressed as µg/L.

TABLE A48-2. (CONTINUED) - ACID COMPOUNDS^a

| Parameter | Method | Detection Limits | 100% G.W. | 10% G.W. | 1% G.W. | Diluent H ₂ O |
|----------------------------|----------|---------------------|--------------|-------------|------------|-----------------------------|
| Phenol | EPA 8270 | 10 | ND | ND | ND | ND |
| 2-Chlorophenol | EPA 8270 | 10 | ND | ND | ND | ND |
| 2-Nitrophenol | EPA 8270 | 10 | ND | ND | ND | ND |
| 2,4-Dimethylphenol | EPA 8270 | 10 | ND | ND | ND | ND |
| 2,4-Dichlorophenol | EPA 8270 | 10 | ND | ND | ND | ND |
| 4-Chloro-3-methylphenol | EPA 8270 | 20 | ND | ND | ND | ND |
| 2,4,6-Trichlorophenol | EPA 8270 | 10 | ND | ND | ND | ND |
| 2,4-Dinitrophenol | EPA 8270 | 50 | ND | ND | ND | ND |
| 4-Nitrophenol | EPA 8270 | 50 | ND | ND | ND | ND |
| 4,6-Dinitro-2-methylphenol | EPA 8270 | 50 | ND | ND | ND | ND |
| Pentachlorophenol | EPA 8270 | 50 | ND | ND | ND | ND |

^a Results expressed as µg/L.

TABLE A48-3. RESULTS OF THE APG-EA BEACH POINT GROUNDWATER AND DILUENT WATER
COMPREHENSIVE CHEMICAL ANALYSES (TEST NO. 3) - GENERAL WATER QUALITY^a

| Parameter | Method | Detection Limits | 100% G.W. | 10% G.W. | 1% G.W. | Diluent H ₂ O |
|---------------------------------------|---------------------------|---------------------|--------------|-------------|------------|-----------------------------|
| Total Alkalinity (CaCO ₃) | EPA 310.1 | 1 | ND | 20 | 22 | 25 |
| Ammonia (N) | SM 4500-NH ₃ E | 0.1 | ND | ND | ND | ND |
| Chloride (Cl) | EPA 325.3 | b | 580 | 78 | 27 | 21 |
| Total Cyanide (CN) | EPA 335.3 | 0.01 | ND | ND | ND | ND |
| Fluoride (F) | EPA 340.2 | 0.05 | 0.24 | 0.88 | 0.98 | 0.94 |
| Hardness, (CaCO ₃) | EPA 130.2 | c | 300 | 92 | 76 | 81 |
| Nitrate (N) | EPA 353.2 | 0.1 | ND | 1.7 | 1.8 | 1.8 |
| Nitrite (N) | EPA 354.1 | 0.01 | ND | ND | ND | ND |
| Total Phosphorus (P) | EPA 365.2 | 0.01 | 0.01 | 0.39 | 0.44 | 0.43 |
| Total Suspended Solids | EPA 160.2 | 5 | ND | ND | ND | ND |
| Specific Conductance | EPA 120.1 | N/A | 1680 | 376 | 212 | 196 |
| Sulfate (SO ₄) | EPA 375.4 | 1 | 75 | 21 | 16 | 17 |
| Sulfide (S) | EPA 376.1 | 1 | ND | ND | ND | ND |
| TOC | EPA 415.1 | 1 | 2 | ND | ND | ND |
| Bromide (Br) | SM 4500-Br | 0.1 | 0.83 | 0.26 | ND | ND |

^a All results expressed as mg/L except for specific conductance which is expressed as $\mu\text{mhos/cm}$.

^b The detection limit was 10 mg/L for the 100% groundwater sample and 1 mg/L for the 10% and 1% groundwater and dilution water samples.

^c The detection limit was 5 mg/L for the 100% groundwater sample and 1 mg/L for the 10% and 1% groundwater and dilution water samples.

TABLE A48-3. (CONTINUED) - METALS^a

| Parameter | Method | Detection Limits | 100% G.W. | 10% G.W. | 1% G.W. | Diluent H ₂ O |
|-----------------|-----------|------------------|-----------|----------|---------|--------------------------|
| Aluminum (Al) | EPA 200.7 | 0.1 | 1.3 | 0.1 | ND | ND |
| Antimony (Sb) | EPA 204.2 | 0.005 | ND | ND | ND | ND |
| Arsenic (As) | EPA 206.2 | 0.005 | ND | ND | ND | ND |
| Beryllium (Be) | EPA 200.7 | 0.005 | ND | ND | ND | ND |
| Boron (B) | EPA 200.7 | 0.05 | 0.05 | ND | ND | ND |
| Cadmium (Cd) | EPA 213.2 | 0.0005 | 0.0014 | ND | ND | ND |
| Calcium (Ca) | EPA 200.7 | 0.5 | 19 | 16 | 16 | 16 |
| Chromium (Cr) | EPA 218.2 | 0.002 | ND | ND | ND | ND |
| Cobalt (Co) | EPA 200.7 | 0.02 | 0.10 | ND | ND | ND |
| Copper (Cu) | EPA 220.2 | 0.005 | 0.008 | ND | ND | ND |
| Iron (Fe) | EPA 200.7 | 0.1 | 7.5 | 0.62 | 0.13 | ND |
| Lead (Pb) | EPA 239.2 | 0.005 | ND | ND | ND | ND |
| Magnesium (Mg) | EPA 200.7 | 0.5 | 37 | 7.7 | 4.1 | 4.2 |
| Manganese (Mn) | EPA 200.7 | 0.01 | 2.1 | 0.22 | 0.02 | ND |
| Mercury (Hg) | EPA 245.1 | 0.0005 | ND | ND | ND | ND |
| Molybdenum (Mo) | EPA 200.7 | 0.02 | ND | ND | ND | ND |
| Nickel (Ni) | EPA 200.7 | 0.02 | 0.16 | 0.02 | 0.06 | 0.03 |
| Potassium (K) | EPA 200.7 | 0.5 | 3.1 | 2.5 | 1.7 | 2.0 |
| Selenium (Se) | EPA 270.2 | 0.005 | ND | ND | ND | ND |
| Silver (Ag) | EPA 272.2 | 0.001 | ND | ND | ND | ND |
| Sodium (Na) | EPA 200.7 | 0.5 | 300 | 40 | 12 | 10 |
| Thallium (Tl) | EPA 279.2 | 0.005 | ND | ND | ND | ND |
| Tin (Sn) | EPA 282.2 | 0.005 | ND | ND | ND | ND |
| Zinc (Zn) | EPA 200.7 | 0.05 | 0.30 | 0.44 | 0.45 | 0.47 |

^a Results expressed as mg/L.

TABLE A48-3. (CONTINUED) - VOLATILES^a

| Parameter | Method | Detection Limits | 100% G.W. | 10% G.W. | 1% G.W. | Diluent H ₂ O |
|---------------------------|----------|------------------|-----------|----------|---------|--------------------------|
| Chloromethane | EPA 8021 | 1 | ND | ND | ND | ND |
| Bromomethane | EPA 8021 | 1 | ND | ND | ND | ND |
| Vinyl chloride | EPA 8021 | 1 | ND | ND | ND | ND |
| Chloroethane | EPA 8021 | 1 | ND | ND | ND | ND |
| Methylene chloride | EPA 8021 | 1 | ND | ND | ND | ND |
| Trichlorofluoromethane | EPA 8021 | 1 | ND | ND | ND | ND |
| 1,1-Dichloroethane | EPA 8021 | 1 | ND | ND | ND | ND |
| 1,2-Dichloroethene | EPA 8021 | 1 | 140 | 6 | b | ND |
| Chloroform | EPA 8021 | 1 | 12 | 5 | 5 | 7 |
| 1,2-Dichloroethane | EPA 8021 | 1 | ND | ND | ND | ND |
| 1,1,1-Trichloroethane | EPA 8021 | 1 | ND | ND | ND | ND |
| Carbon tetrachloride | EPA 8021 | 1 | ND | ND | ND | ND |
| Bromodichloromethane | EPA 8021 | 1 | ND | ND | ND | 1 |
| 1,2-Dichloropropane | EPA 8021 | 1 | ND | ND | ND | ND |
| 1,3-Dichloropropene | EPA 8021 | 1 | ND | ND | ND | ND |
| Dibromochloromethane | EPA 8021 | 1 | ND | ND | ND | ND |
| 1,1,2-Trichloroethane | EPA 8021 | 1 | 37 | 5 | ND | ND |
| 2-Chloroethylvinyl ether | EPA 8021 | 5 | ND | ND | ND | ND |
| Bromoform | EPA 8021 | 2 | ND | ND | ND | ND |
| Tetrachloroethene | EPA 8021 | 1 | 30 | 4 | ND | ND |
| 1,1,2,2-Tetrachloroethane | EPA 8021 | c | 10000 | 900 | 100 | 8 |
| Ethylbenzene | EPA 8021 | 1 | ND | ND | ND | ND |
| 1,1-Dichloroethene | EPA 8021 | 1 | ND | ND | ND | ND |
| Trichloroethene | EPA 8021 | d | 450 | 50 | 5 | ND |
| Benzene | EPA 8021 | 1 | ND | ND | ND | ND |
| Toluene | EPA 8021 | 1 | ND | ND | ND | ND |
| Chlorobenzene | EPA 8021 | 1 | 2 | ND | ND | ND |

^a Results expressed as µg/L.^b Detected below quantitation level.^c Detection limits were as follows: 1000 µg/L for the 100% groundwater sample, 100 µg/L for the 10% groundwater sample, 10 µg/L for the 1% groundwater sample, and 1 µg/L for the diluent water sample.^d Detection limits were as follows: 20 µg/L for the 100% groundwater sample, 10 µg/L for the 10% groundwater sample, and 1 µg/L for the 1% groundwater and diluent water samples.

TABLE A48-3. (CONTINUED) - BASE/NEUTRAL COMPOUNDS^a

| Parameter | Method | Detection Limits | 100% G.W. | 10% G.W. | 1% G.W. | Diluent H ₂ O |
|-----------------------------|----------|------------------|-----------|----------|---------|--------------------------|
| Acenaphthene | EPA 8270 | 10 | ND | ND | ND | ND |
| 2,4-Dinitrotoluene | EPA 8270 | 10 | ND | ND | ND | ND |
| 2,6-Dinitrotoluene | EPA 8270 | 10 | ND | ND | ND | ND |
| Diethyl phthalate | EPA 8270 | 10 | ND | ND | ND | ND |
| 4-Chlorophenyl phenyl ether | EPA 8270 | 10 | ND | ND | ND | ND |
| Fluorene | EPA 8270 | 10 | ND | ND | ND | ND |
| N-Nitrosodiphenylamine | EPA 8270 | 10 | ND | ND | ND | ND |
| 1,2-Diphenylhydrazine | EPA 8270 | 10 | ND | ND | ND | ND |
| 4-Bromophenyl phenyl ether | EPA 8270 | 10 | ND | ND | ND | ND |
| Hexachlorobenzene | EPA 8270 | 10 | ND | ND | ND | ND |
| Phenanthrene | EPA 8270 | 10 | ND | ND | ND | ND |
| Anthracene | EPA 8270 | 10 | ND | ND | ND | ND |
| Di-n-butyl phthalate | EPA 8270 | 10 | ND | ND | ND | ND |
| Fluoranthene | EPA 8270 | 10 | ND | ND | ND | ND |
| Pyrene | EPA 8270 | 10 | ND | ND | ND | ND |
| Benzidine | EPA 8270 | 50 | ND | ND | ND | ND |
| Butylbenzyl phthalate | EPA 8270 | 10 | ND | ND | ND | ND |
| 3,3'-Dichlorobenzidine | EPA 8270 | 20 | ND | ND | ND | ND |
| Benzo(a)anthracene | EPA 8270 | 10 | ND | ND | ND | ND |
| bis(2-Ethylhexyl) phthalate | EPA 8270 | 10 | ND | ND | ND | ND |
| Chrysene | EPA 8270 | 10 | ND | ND | ND | ND |
| Di-n-octyl phthalate | EPA 8270 | 10 | ND | ND | ND | ND |
| Benzo(b)fluoranthene | EPA 8270 | 10 | ND | ND | ND | ND |
| Benzo(k)fluoranthene | EPA 8270 | 10 | ND | ND | ND | ND |
| Benzo(a)pyrene | EPA 8270 | 10 | ND | ND | ND | ND |
| Indeno(1,2,3-cd)pyrene | EPA 8270 | 10 | ND | ND | ND | ND |
| Dibenz(a,h)anthracene | EPA 8270 | 10 | ND | ND | ND | ND |
| Benzo(g,h,i)perylene | EPA 8270 | 10 | ND | ND | ND | ND |

^a Results expressed as µg/L.

TABLE A48-3. (CONTINUED) - BASE/NEUTRAL COMPOUNDS CON'T^a

| Parameter | Method | Detection Limits | 100% G.W. | 10% G.W. | 1% G.W. | Diluent |
|------------------------------|----------|------------------|-----------|----------|---------|---------|
| N-Nitrosodimethylamine | EPA 8270 | 10 | ND | ND | ND | ND |
| bis(2-Chloroethyl) ether | EPA 8270 | 10 | ND | ND | ND | ND |
| 1,3-Dichlorobenzene | EPA 8270 | 10 | ND | ND | ND | ND |
| 1,4-Dichlorobenzene | EPA 8270 | 10 | ND | ND | ND | ND |
| 1,2-Dichlorobenzene | EPA 8270 | 10 | ND | ND | ND | ND |
| bis(2-Chloroisopropyl) ether | EPA 8270 | 10 | ND | ND | ND | ND |
| N-Nitrosodi-n-propylamine | EPA 8270 | 10 | ND | ND | ND | ND |
| Hexachloroethane | EPA 8270 | 10 | ND | ND | ND | ND |
| Nitrobenzene | EPA 8270 | 10 | ND | ND | ND | ND |
| Isophorone | EPA 8270 | 10 | ND | ND | ND | ND |
| bis(2-Chloroethoxy)methane | EPA 8270 | 10 | ND | ND | ND | ND |
| 1,2,4-Trichlorobenzene | EPA 8270 | 10 | ND | ND | ND | ND |
| Naphthalene | EPA 8270 | 10 | ND | ND | ND | ND |
| Hexachlorobutadiene | EPA 8270 | 10 | ND | ND | ND | ND |
| Hexachlorocyclopentadiene | EPA 8270 | 10 | ND | ND | ND | ND |
| 2-Chloronaphthalene | EPA 8270 | 10 | ND | ND | ND | ND |
| Dimethyl phthalate | EPA 8270 | 10 | ND | ND | ND | ND |
| Acenaphthylene | EPA 8270 | 10 | ND | ND | ND | ND |

^a Results expressed as µg/L.

TABLE A48-3. (CONTINUED) - PESTICIDES^a

| Parameter | Method | Detection Limits | 100% G.W. | 10% G.W. | 1% G.W. | Diluent H ₂ O |
|--------------------|----------|------------------|-----------|----------|---------|--------------------------|
| a-BHC | EPA 8080 | 0.05 | ND | ND | ND | ND |
| b-BHC | EPA 8080 | 0.05 | ND | ND | ND | ND |
| g-BHC | EPA 8080 | 0.05 | ND | ND | ND | ND |
| d-BHC | EPA 8080 | 0.05 | ND | ND | ND | ND |
| Heptachlor | EPA 8080 | 0.05 | ND | ND | ND | ND |
| Aldrin | EPA 8080 | 0.05 | ND | ND | ND | ND |
| Heptachlor epoxide | EPA 8080 | 0.05 | ND | ND | ND | ND |
| a-Endosulfan | EPA 8080 | 0.1 | ND | ND | ND | ND |
| 4,4'-DDE | EPA 8080 | 0.1 | ND | ND | ND | ND |
| Dieldrin | EPA 8080 | 0.1 | ND | ND | ND | ND |
| Endrin | EPA 8080 | 0.1 | ND | ND | ND | ND |
| b-Endosulfan | EPA 8080 | 0.3 | ND | ND | ND | ND |
| Endrin aldehyde | EPA 8080 | 0.3 | ND | ND | ND | ND |
| 4,4'-DDD | EPA 8080 | 0.3 | ND | ND | ND | ND |
| Endosulfan sulfate | EPA 8080 | 0.3 | ND | ND | ND | ND |
| 4,4'-DDT | EPA 8080 | 0.3 | ND | ND | ND | ND |
| PCB-1242 | EPA 8080 | 1 | ND | ND | ND | ND |
| PCB-1254 | EPA 8080 | 1 | ND | ND | ND | ND |
| PCB-1221 | EPA 8080 | 1 | ND | ND | ND | ND |
| PCB-1232 | EPA 8080 | 1 | ND | ND | ND | ND |
| PCB-1248 | EPA 8080 | 1 | ND | ND | ND | ND |
| PCB-1260 | EPA 8080 | 1 | ND | ND | ND | ND |
| PCB-1016 | EPA 8080 | 1 | ND | ND | ND | ND |
| Chlordane | EPA 8080 | 1 | ND | ND | ND | ND |
| Toxaphene | EPA 8080 | 3 | ND | ND | ND | ND |

^a Results expressed as µg/L.

TABLE A48-3. (CONTINUED) - HERBICIDES AND ORGANOPHOSPHORUS PESTICIDES^a

| Parameter | Method | Detection Limits | 100% G.W. | 10% G.W. | 1% G.W. | Diluent H ₂ O |
|-------------------------------------|----------|------------------|-----------|----------|---------|--------------------------|
| <u>Herbicides:</u> | | | | | | |
| 2,4-D | EPA 8150 | 0.5 | ND | ND | ND | ND |
| 2,4,5-TP (Silvex) | EPA 8150 | 0.2 | ND | ND | ND | ND |
| <u>Organophosphorus Pesticides:</u> | | | | | | |
| Malathion | EPA 8080 | 0.5 | ND | ND | ND | ND |
| Methyl azinphos | EPA 8080 | 3 | ND | ND | ND | ND |
| Coumaphos | EPA 8080 | 2 | ND | ND | ND | ND |
| Chlorpyrifos | EPA 8080 | 0.1 | ND | ND | ND | ND |
| Diazinon | EPA 8080 | 0.5 | ND | ND | ND | ND |
| Fenthion | EPA 8080 | 50 | ND | ND | ND | ND |
| Naled | EPA 8080 | 5 | ND | ND | ND | ND |
| Methyl parathion | EPA 8080 | 0.5 | ND | ND | ND | ND |
| Parathion | EPA 8080 | 0.5 | ND | ND | ND | ND |

^a Results expressed as µg/L.

TABLE A48-3. (CONTINUED) - ACID COMPOUNDS^a

| Parameter | Method | Detection Limits | 100% G.W. | 10% G.W. | 1% G.W. | Diluent H ₂ O |
|----------------------------|----------|------------------|-----------|----------|---------|--------------------------|
| Phenol | EPA 8270 | 10 | ND | ND | ND | ND |
| 2-Chlorophenol | EPA 8270 | 10 | ND | ND | ND | ND |
| 2-Nitrophenol | EPA 8270 | 10 | ND | ND | ND | ND |
| 2,4-Dimethylphenol | EPA 8270 | 10 | ND | ND | ND | ND |
| 2,4-Dichlorophenol | EPA 8270 | 10 | ND | ND | ND | ND |
| 4-Chloro-3-methylphenol | EPA 8270 | 20 | ND | ND | ND | ND |
| 2,4,6-Trichlorophenol | EPA 8270 | 10 | ND | ND | ND | ND |
| 2,4-Dinitrophenol | EPA 8270 | 50 | ND | ND | ND | ND |
| 4-Nitrophenol | EPA 8270 | 50 | ND | ND | ND | ND |
| 4,6-Dinitro-2-methylphenol | EPA 8270 | 50 | ND | ND | ND | ND |
| Pentachlorophenol | EPA 8270 | 50 | ND | ND | ND | ND |

^a Results expressed as µg/L.

TABLE A48-4. RESULTS OF THE APG-EA BEACH POINT GROUNDWATER AND DILUENT WATER COMPREHENSIVE CHEMICAL ANALYSES (TEST NO. 4) - GENERAL WATER QUALITY^a

| Parameter | Method | Detection Limits | 100% G.W. | 10% G.W. | 1% G.W. | Diluent H ₂ O |
|---------------------------------------|---------------------------|------------------|-----------|----------|---------|--------------------------|
| Total Alkalinity (CaCO ₃) | EPA 310.1 | 1 | ND | 18 | 23 | 21 |
| Ammonia (N) | SM 4500-NH ₃ E | 0.1 | ND | ND | ND | ND |
| Chloride (Cl) | EPA 325.3 | ^b | 570 | 72 | 25 | 19 |
| Total Cyanide (CN) | EPA 335.3 | 0.01 | ND | ND | ND | ND |
| Fluoride (F) | EPA 340.2 | 0.1 | 0.2 | 0.6 | 0.7 | 0.7 |
| Hardness, (CaCO ₃) | EPA 130.2 | ^c | 230 | 89 | 68 | 76 |
| Nitrate (N) | EPA 353.2 | 0.1 | ND | 1.9 | 2.1 | 2.2 |
| Nitrite (N) | EPA 354.1 | 0.01 | ND | ND | ND | ND |
| Total Phosphorus (P) | EPA 365.2 | 0.01 | ND | 0.32 | 0.36 | 0.37 |
| Total Suspended Solids | EPA 160.2 | 5 | ND | ND | ND | ND |
| Specific Conductance | EPA 120.1 | N/A | 2020 | 388 | 210 | 180 |
| Sulfate (SO ₄) | EPA 375.4 | ^c | 82 | 29 | 22 | 19 |
| Sulfide (S) | EPA 376.1 | 1 | ND | ND | ND | ND |
| TOC | EPA 415.1 | 1 | 3 | ND | ND | ND |
| Bromide (Br) | SM 4500-Br | 0.1 | 0.4 | 0.3 | 0.1 | ND |

^a All results expressed as mg/L except for specific conductance which is expressed as $\mu\text{mohs/cm}$.

^b The detection limit was 10 mg/L for the 100% groundwater sample and 1 mg/L for the 10% and 1% groundwater and dilution water samples.

^c The detection limit was 2 mg/L for the 100% groundwater sample and 1 mg/L for the 10% and 1% groundwater and dilution water samples.

TABLE 48-4. (CONTINUED) - METALS^a

| Parameter | Method | Detection Limits | 100% G.W. | 10% G.W. | 1% G.W. | Diluent H ₂ O |
|-----------------|-----------|------------------|-----------|----------|---------|--------------------------|
| Aluminum (Al) | EPA 200.7 | 0.1 | 1.5 | 0.3 | ND | ND |
| Antimony (Sb) | EPA 204.2 | 0.005 | ND | ND | ND | ND |
| Arsenic (As) | EPA 206.2 | 0.005 | 0.007 | ND | ND | ND |
| Beryllium (Be) | EPA 200.7 | 0.005 | ND | ND | ND | ND |
| Boron (B) | EPA 200.7 | 0.05 | 0.07 | ND | ND | ND |
| Cadmium (Cd) | EPA 213.2 | 0.0005 | 0.0009 | ND | ND | ND |
| Calcium (Ca) | EPA 200.7 | 0.5 | 21 | 19 | 17 | 17 |
| Chromium (Cr) | EPA 218.2 | 0.002 | ND | ND | ND | ND |
| Cobalt (Co) | EPA 200.7 | 0.05 | 0.08 | ND | ND | ND |
| Copper (Cu) | EPA 220.2 | 0.005 | 0.006 | ND | ND | ND |
| Iron (Fe) | EPA 200.7 | 0.1 | 7.7 | 0.8 | 0.3 | ND |
| Lead (Pb) | EPA 239.2 | 0.005 | ND | ND | ND | ND |
| Magnesium (Mg) | EPA 200.7 | 0.5 | 43 | 9.9 | 6.1 | 5.5 |
| Manganese (Mn) | EPA 200.7 | 0.01 | 2.1 | 0.22 | 0.03 | ND |
| Mercury (Hg) | EPA 245.1 | 0.0005 | ND | ND | ND | ND |
| Molybdenum (Mo) | EPA 200.7 | 0.05 | ND | ND | ND | ND |
| Nickel (Ni) | EPA 200.7 | 0.05 | 0.18 | ND | ND | ND |
| Potassium (K) | EPA 200.7 | 0.5 | 3.9 | 2.9 | 2.9 | 2.7 |
| Selenium (Se) | EPA 270.2 | 0.005 | ND | ND | ND | ND |
| Silver (Ag) | EPA 272.2 | 0.001 | ND | ND | ND | ND |
| Sodium (Na) | EPA 200.7 | 0.5 | 350 | 41 | 13 | 9.1 |
| Thallium (Tl) | EPA 279.2 | 0.005 | ND | ND | ND | ND |
| Tin (Sn) | EPA 282.2 | 0.005 | ND | ND | ND | ND |
| Zinc (Zn) | EPA 200.7 | 0.05 | 0.31 | 0.62 | 0.66 | 0.60 |

^a Results expressed as mg/L.

TABLE 48-4. (CONTINUED) - VOLATILES^a

| Parameter | Method | Detection Limits | 100% G.W. | 10% G.W. | 1% G.W. | Diluent H ₂ O |
|---------------------------|----------|------------------|-----------|----------|---------|--------------------------|
| Chloromethane | EPA 8021 | 2 | ND | ND | ND | ND |
| Bromomethane | EPA 8021 | 5 | ND | ND | ND | ND |
| Vinyl chloride | EPA 8021 | 2 | ND | ND | ND | ND |
| Chloroethane | EPA 8021 | 1 | ND | ND | ND | ND |
| Methylene chloride | EPA 8021 | 1 | ND | ND | ND | ND |
| Trichlorofluoromethane | EPA 8021 | 1 | ND | ND | ND | ND |
| 1,1-Dichloroethane | EPA 8021 | 1 | ND | ND | ND | ND |
| 1,2-Dichloroethene | EPA 8021 | 1 | 110 | 4 | ND | ND |
| Chloroform | EPA 8021 | 1 | 12 | 4 | 3 | 3 |
| 1,2-Dichloroethane | EPA 8021 | 1 | ND | ND | ND | ND |
| 1,1,1-Trichloroethane | EPA 8021 | 1 | ND | ND | ND | ND |
| Carbon tetrachloride | EPA 8021 | 1 | ND | ND | ND | ND |
| Bromodichloromethane | EPA 8021 | 1 | ND | ND | ND | ND |
| 1,2-Dichloropropane | EPA 8021 | 1 | ND | ND | ND | ND |
| 1,3-Dichloropropene | EPA 8021 | 1 | ND | ND | ND | ND |
| Dibromochloromethane | EPA 8021 | 1 | ND | ND | ND | ND |
| 1,1,2-Trichloroethane | EPA 8021 | 1 | 70 | 4 | ND | ND |
| 2-Chloroethylvinyl ether | EPA 8021 | 5 | ND | ND | ND | ND |
| Bromoform | EPA 8021 | 5 | ND | ND | ND | ND |
| Tetrachloroethene | EPA 8021 | 1 | 80 | 3 | ND | ND |
| 1,1,2,2-Tetrachloroethane | EPA 8021 | b | 9000 | 700 | 91 | 4 |
| Ethylbenzene | EPA 8021 | 1 | ND | ND | ND | ND |
| 1,1-Dichloroethene | EPA 8021 | 1 | ND | ND | ND | ND |
| Trichloroethene | EPA 8021 | 1 | 780 | 40 | 4 | ND |
| Benzene | EPA 8021 | 1 | ND | ND | ND | ND |
| Toluene | EPA 8021 | 1 | ND | ND | ND | ND |
| Chlorobenzene | EPA 8021 | 1 | ND | ND | ND | ND |

^a Results expressed as $\mu\text{g/L}$.^b The detection limit was 10 $\mu\text{g/L}$ for the 100% groundwater sample and 1 $\mu\text{g/L}$ for the 10% and 1% groundwater and diluent water samples.

TABLE A48-4. (CONTINUED) - BASE/NEUTRAL COMPOUNDS^a

| Parameter | Method | Detection Limits | 100% G.W. | 10% G.W. | 1% G.W. | Diluent H ₂ O |
|-----------------------------|-----------|------------------|-----------|----------|---------|--------------------------|
| Acenaphthene | EPA 8270A | 10 | ND | ND | ND | ND |
| 2,4-Dinitrotoluene | EPA 8270A | 10 | ND | ND | ND | ND |
| 2,6-Dinitrotoluene | EPA 8270A | 10 | ND | ND | ND | ND |
| Diethyl phthalate | EPA 8270A | 10 | ND | ND | ND | ND |
| 4-Chlorophenyl phenyl ether | EPA 8270A | 10 | ND | ND | ND | ND |
| Fluorene | EPA 8270A | 10 | ND | ND | ND | ND |
| N-Nitrosodiphenylamine | EPA 8270A | 10 | ND | ND | ND | ND |
| 1,2-Diphenylhydrazine | EPA 8270A | 10 | ND | ND | ND | ND |
| 4-Bromophenyl phenyl ether | EPA 8270A | 10 | ND | ND | ND | ND |
| Hexachlorobenzene | EPA 8270A | 10 | ND | ND | ND | ND |
| Phenanthrene | EPA 8270A | 10 | ND | ND | ND | ND |
| Anthracene | EPA 8270A | 10 | ND | ND | ND | ND |
| Di-n-butyl phthalate | EPA 8270A | 10 | ND | ND | ND | ND |
| Fluoranthene | EPA 8270A | 10 | ND | ND | ND | ND |
| Pyrene | EPA 8270A | 10 | ND | ND | ND | ND |
| Benzidine | EPA 8270A | 50 | ND | ND | ND | ND |
| Butylbenzyl phthalate | EPA 8270A | 10 | ND | ND | ND | ND |
| 3,3'-Dichlorobenzidine | EPA 8270A | 20 | ND | ND | ND | ND |
| Benzo(a)anthracene | EPA 8270A | 10 | ND | ND | ND | ND |
| bis(2-Ethylhexyl) phthalate | EPA 8270A | 10 | ND | ND | ND | ND |
| Chrysene | EPA 8270A | 10 | ND | ND | ND | ND |
| Di-n-octyl phthalate | EPA 8270A | 10 | ND | ND | ND | ND |
| Benzo(b)fluoranthene | EPA 8270A | 10 | ND | ND | ND | ND |
| Benzo(k)fluoranthene | EPA 8270A | 10 | ND | ND | ND | ND |
| Benzo(a)pyrene | EPA 8270A | 10 | ND | ND | ND | ND |
| Indeno(1,2,3-cd)pyrene | EPA 8270A | 10 | ND | ND | ND | ND |
| Dibenz(a,h)anthracene | EPA 8270A | 10 | ND | ND | ND | ND |
| Benzo(g,h,i)perylene | EPA 8270A | 10 | ND | ND | ND | ND |

^a Results expressed as µg/L.

TABLE A48-4. (CONTINUED) - BASE/NEUTRAL COMPOUNDS CON'T^a

| Parameter | Method | Detection Limits | 100% G.W. | 10% G.W. | 1% G.W. | Diluent H ₂ O |
|------------------------------|-----------|------------------|-----------|----------|---------|--------------------------|
| N-Nitrosodimethylamine | EPA 8270A | 10 | ND | ND | ND | ND |
| bis(2-Chloroethyl) ether | EPA 8270A | 10 | ND | ND | ND | ND |
| 1,3-Dichlorobenzene | EPA 8270A | 10 | ND | ND | ND | ND |
| 1,4-Dichlorobenzene | EPA 8270A | 10 | ND | ND | ND | ND |
| 1,2-Dichlorobenzene | EPA 8270A | 10 | ND | ND | ND | ND |
| bis(2-Chloroisopropyl) ether | EPA 8270A | 10 | ND | ND | ND | ND |
| N-Nitrosodi-n-propylamine | EPA 8270A | 10 | ND | ND | ND | ND |
| Hexachloroethane | EPA 8270A | 10 | ND | ND | ND | ND |
| Nitrobenzene | EPA 8270A | 10 | ND | ND | ND | ND |
| Isophorone | EPA 8270A | 10 | ND | ND | ND | ND |
| bis(2-Chloroethoxy)methane | EPA 8270A | 10 | ND | ND | ND | ND |
| 1,2,4-Trichlorobenzene | EPA 8270A | 10 | ND | ND | ND | ND |
| Naphthalene | EPA 8270A | 10 | ND | ND | ND | ND |
| Hexachlorobutadiene | EPA 8270A | 10 | ND | ND | ND | ND |
| Hexachlorocyclopentadiene | EPA 8270A | 10 | ND | ND | ND | ND |
| 2-Chloronaphthalene | EPA 8270A | 10 | ND | ND | ND | ND |
| Dimethyl phthalate | EPA 8270A | 10 | ND | ND | ND | ND |
| Acenaphthylene | EPA 8270A | 10 | ND | ND | ND | ND |

^a Results expressed as µg/L.

TABLE A48-4. (CONTINUED) - PESTICIDES^a

| Parameter | Method | Detection Limits | 100% G.W. | 10% G.W. | 1% G.W. | Diluent H ₂ O |
|--------------------|----------|------------------|-----------|----------|---------|--------------------------|
| a-BHC | EPA 8080 | 0.05 | ND | ND | ND | ND |
| b-BHC | EPA 8080 | 0.05 | ND | ND | ND | ND |
| g-BHC | EPA 8080 | 0.05 | ND | ND | ND | ND |
| d-BHC | EPA 8080 | 0.05 | ND | ND | ND | ND |
| Heptachlor | EPA 8080 | 0.05 | ND | ND | ND | ND |
| Aldrin | EPA 8080 | 0.05 | ND | ND | ND | ND |
| Heptachlor epoxide | EPA 8080 | 0.05 | ND | ND | ND | ND |
| a-Endosulfan | EPA 8080 | 0.1 | ND | ND | ND | ND |
| 4,4'-DDE | EPA 8080 | 0.1 | ND | ND | ND | ND |
| Dieldrin | EPA 8080 | 0.1 | ND | ND | ND | ND |
| Endrin | EPA 8080 | 0.1 | ND | ND | ND | ND |
| b-Endosulfan | EPA 8080 | 0.3 | ND | ND | ND | ND |
| Endrin aldehyde | EPA 8080 | 0.3 | ND | ND | ND | ND |
| 4,4'-DDD | EPA 8080 | 0.3 | ND | ND | ND | ND |
| Endosulfan sulfate | EPA 8080 | 0.3 | ND | ND | ND | ND |
| 4,4'-DDT | EPA 8080 | 0.3 | ND | ND | ND | ND |
| PCB-1242 | EPA 8080 | 1 | ND | ND | ND | ND |
| PCB-1254 | EPA 8080 | 1 | ND | ND | ND | ND |
| PCB-1221 | EPA 8080 | 1 | ND | ND | ND | ND |
| PCB-1232 | EPA 8080 | 1 | ND | ND | ND | ND |
| PCB-1248 | EPA 8080 | 1 | ND | ND | ND | ND |
| PCB-1260 | EPA 8080 | 1 | ND | ND | ND | ND |
| PCB-1016 | EPA 8080 | 1 | ND | ND | ND | ND |
| Chlordane | EPA 8080 | 1 | ND | ND | ND | ND |
| Toxaphene | EPA 8080 | 3 | ND | ND | ND | ND |

^a Results expressed as µg/L.

TABLE A48-4. (CONTINUED) - HERBICIDES AND ORGANOPHOSPHORUS PESTICIDES^a

| Parameter | Method | Detection Limits | 100% G.W. | 10% G.W. | 1% G.W. | Diluent H ₂ O |
|-------------------------------------|-----------|------------------|-----------|----------|---------|--------------------------|
| <u>Herbicides:</u> | | | | | | |
| 2,4-D | EPA 8150A | 0.5 | ND | ND | ND | ND |
| 2,4,5-TP (Silvex) | EPA 8150A | 0.2 | ND | ND | ND | ND |
| <u>Organophosphorus Pesticides:</u> | | | | | | |
| Malathion | EPA 8080 | 0.5 | ND | ND | ND | ND |
| Methyl azinphos | EPA 8080 | 3 | ND | ND | ND | ND |
| Coumaphos | EPA 8080 | 2 | ND | ND | ND | ND |
| Chlorpyrifos | EPA 8080 | 0.1 | ND | ND | ND | ND |
| Diazinon | EPA 8080 | 0.5 | ND | ND | ND | ND |
| Fenthion | EPA 8080 | 50 | ND | ND | ND | ND |
| Naled | EPA 8080 | 5 | ND | ND | ND | ND |
| Methyl parathion | EPA 8080 | 0.5 | ND | ND | ND | ND |
| Parathion | EPA 8080 | 0.5 | ND | ND | ND | ND |

^a Results expressed as µg/L.

TABLE A48-4. (CONTINUED) - ACID COMPOUNDS^a

| Parameter | Method | Detection Limits | 100% G.W. | 10% G.W. | 1% G.W. | Diluent H ₂ O |
|----------------------------|-----------|---------------------|--------------|-------------|------------|-----------------------------|
| Phenol | EPA 8270A | 10 | ND | ND | ND | ND |
| 2-Chlorophenol | EPA 8270A | 10 | ND | ND | ND | ND |
| 2-Nitrophenol | EPA 8270A | 10 | ND | ND | ND | ND |
| 2,4-Dimethylphenol | EPA 8270A | 10 | ND | ND | ND | ND |
| 2,4-Dichlorophenol | EPA 8270A | 10 | ND | ND | ND | ND |
| 4-Chloro-3-methylphenol | EPA 8270A | 20 | ND | ND | ND | ND |
| 2,4,6-Trichlorophenol | EPA 8270A | 10 | ND | ND | ND | ND |
| 2,4-Dinitrophenol | EPA 8270A | 50 | ND | ND | ND | ND |
| 4-Nitrophenol | EPA 8270A | 50 | ND | ND | ND | ND |
| 4,6-Dinitro-2-methylphenol | EPA 8270A | 50 | ND | ND | ND | ND |
| Pentachlorophenol | EPA 8270A | 50 | ND | ND | ND | ND |

^a Results expressed as µg/L.

APPENDIX 49

ROUTINE WATER QUALITY OF THE CHRONIC
HISTOPATHOLOGY EXPOSURE TANKS

TANK No. 1
TANK CONCENTRATION: 100% DILUENT WATER
NO DEN EXPOSURE

| Date | Test Day | Time (Military) | Temperature (Celsius) | pH | Dissolved Oxygen (mg/L) | Alkalinity (mg/L as CaCO3) | Hardness (mg/L as CaCO3) | Conductivity (umohs/cm) | Total Residual Chlorine (mg/L) | Free Available Chlorine (mg/L) | Total Ammonia-Nitrogen (mg/L) | Total Unionized Ammonia-Nitrogen (mg/L) |
|----------|----------|-----------------|-----------------------|------|-------------------------|----------------------------|--------------------------|-------------------------|--------------------------------|--------------------------------|-------------------------------|---|
| 02-25-93 | 1 | 1500 | 24.8 | 6.63 | 7.60 | | | | | | | |
| 02-26-93 | 2 | 1000 | 24.1 | 7.17 | 7.66 | | | | | | | |
| 02-27-93 | 3 | 1000 | 24.5 | 7.11 | 7.59 | | | | | | | |
| 02-28-93 | 4 | 1100 | 24.6 | 7.09 | 7.54 | 34.0 | 85.5 | | | | | |
| 03-01-93 | 5 | 930 | 24.9 | 7.09 | 7.32 | | | | | | | |
| 03-02-93 | 6 | 1000 | 25.5 | 7.05 | 7.13 | | | | | | | |
| 03-03-93 | 7 | 1000 | 26.0 | 7.32 | 7.66 | | | | | | | |
| 03-04-93 | 8 | 1000 | 25.2 | 7.45 | 7.74 | | | | | | | |
| 03-05-93 | 9 | 800 | 25.5 | 7.43 | 7.49 | | | | | | | |
| 03-06-93 | 10 | 1000 | 24.9 | 7.28 | 7.81 | 34.0 | 85.5 | 294 | | | | |
| 03-07-93 | 11 | 1100 | 25.0 | 7.07 | 7.80 | | | | | | | |
| 03-08-93 | 12 | 900 | 25.0 | 7.06 | 7.73 | | | | | | | |
| 03-09-93 | 13 | 930 | 25.3 | 7.19 | 7.58 | | | | | | | |
| 03-10-93 | 14 | 900 | 24.9 | 7.07 | 7.73 | | | | | | | |
| 03-11-93 | 15 | 900 | 25.2 | 7.46 | 7.62 | | | | | | | |
| 03-12-93 | 16 | 800 | 24.6 | 7.41 | 8.10 | 54.4 | 85.5 | 194 | 0.00 | 0.00 | | |
| 03-13-93 | 17 | 1600 | 24.8 | 7.18 | 7.86 | | | | | | | |
| 03-14-93 | 18 | 1500 | 24.9 | 7.18 | 7.82 | | | | | | | |
| 03-15-93 | 19 | 900 | 24.7 | 7.11 | 7.97 | | | | | | | |
| 03-16-93 | 20 | 900 | 25.3 | 7.21 | 7.80 | | | | | | | |
| 03-17-93 | 21 | 900 | 25.5 | 7.22 | 7.43 | 34.0 | 85.5 | 411 | 0.00 | 0.00 | 0.019 | 0.00018 |
| 03-18-93 | 22 | 1030 | 25.3 | 7.05 | 7.85 | | | | | | | |
| 03-19-93 | 23 | 1000 | 25.2 | 7.02 | 7.83 | | | | | | | |
| 03-20-93 | 24 | 1100 | 27.0 | 7.15 | 7.59 | | | | | | | |
| 03-21-93 | 25 | 1230 | 25.5 | 7.15 | 7.98 | | | | | | | |
| 03-22-93 | 26 | 900 | 25.0 | 7.22 | 8.00 | | | | | | | |
| 03-23-93 | 27 | 900 | 24.7 | 7.16 | 8.05 | | | | | | | |
| 03-24-93 | 28 | 900 | 25.3 | 7.05 | 7.86 | | | | | | | |
| 03-25-93 | 29 | 900 | 25.2 | 7.00 | 7.78 | 34.0 | 68.4 | 196 | 0.00 | 0.00 | 0.000 | 0.00000 |
| 03-26-93 | 30 | 800 | 24.9 | 7.21 | 8.09 | | | | | | | |
| 03-27-93 | 31 | 1000 | 25.4 | 7.24 | 7.76 | | | | | | | |
| 03-28-93 | 32 | 1000 | 25.7 | 7.36 | 7.64 | | | | | | | |
| 03-29-93 | 33 | 1100 | 25.9 | 7.37 | 7.73 | | | | | | | |
| 03-30-93 | 34 | 1100 | 25.6 | 7.24 | 7.68 | | | | | | | |
| 03-31-93 | 35 | 1000 | 25.6 | 7.36 | 6.30 | | | | | | | |
| 04-01-93 | 36 | 1000 | 24.9 | 7.45 | 8.01 | 40.8 | 68.4 | 189 | 0.00 | 0.00 | 0.000 | 0.00000 |
| 04-02-93 | 37 | 1000 | 24.9 | 7.40 | 8.08 | | | | | | | |
| 04-03-93 | 38 | 1100 | 24.9 | 7.42 | 8.23 | | | | | | | |
| 04-04-93 | 39 | 1300 | 24.7 | 7.40 | 8.39 | | | | | | | |
| 04-05-93 | 40 | 900 | 24.8 | 7.38 | 8.47 | | | | | | | |
| 04-06-93 | 41 | 1100 | 24.4 | 7.42 | 8.21 | | | | | | | |

| | | | | | | | | | | | |
|----------|----|------|------|------|------|------|-------|-----|------|------|---------|
| 04-07-93 | 42 | 900 | 25.0 | 7.49 | 8.34 | 40.8 | 85.5 | 189 | 0.00 | 0.00 | 0.00000 |
| 04-08-93 | 43 | 900 | 25.3 | 7.49 | 8.20 | | | | | | |
| 04-09-93 | 44 | 900 | 25.3 | 7.50 | 7.97 | | | | | | |
| 04-10-93 | 45 | 1000 | 25.7 | 7.46 | 7.82 | | | | | | |
| 04-11-93 | 46 | 800 | 24.8 | 7.48 | 8.27 | | | | | | |
| 04-12-93 | 47 | 1000 | 25.2 | 7.44 | 8.04 | | | | | | |
| 04-13-93 | 48 | 900 | 25.0 | 7.48 | 7.87 | | | | | | |
| 04-14-93 | 49 | 600 | 25.1 | 7.48 | 7.59 | | | | | | |
| 04-15-93 | 50 | 900 | 25.3 | 7.47 | 7.59 | 40.8 | 85.5 | 189 | 0.00 | 0.00 | 0.00051 |
| 04-16-93 | 51 | 1000 | 25.7 | 7.46 | 8.53 | | | | | | |
| 04-17-93 | 52 | 1000 | 25.3 | 7.47 | 7.67 | | | | | | |
| 04-18-93 | 53 | 1000 | 24.8 | 7.50 | 7.83 | | | | | | |
| 04-19-93 | 54 | 900 | 25.1 | 7.52 | 7.89 | | | | | | |
| 04-20-93 | 55 | 900 | 25.5 | 7.46 | 7.81 | | | | | | |
| 04-21-93 | 56 | 1000 | 25.7 | 7.48 | 7.61 | | | | | | |
| 04-22-93 | 57 | 900 | 25.0 | 7.53 | 7.61 | 40.8 | 85.5 | 189 | 0.00 | 0.00 | 0.00000 |
| 04-23-93 | 58 | 900 | 24.9 | 7.51 | 7.48 | | | | | | |
| 04-24-93 | 59 | 1000 | 25.1 | 7.48 | 7.96 | | | | | | |
| 04-25-93 | 60 | 800 | 25.4 | 7.48 | 8.15 | | | | | | |
| 04-26-93 | 61 | 900 | 25.4 | 7.43 | 7.42 | | | | | | |
| 04-27-93 | 62 | 1000 | 24.4 | 7.54 | 8.06 | | | | | | |
| 04-28-93 | 63 | 900 | 25.0 | 7.39 | 7.54 | | | | | | |
| 04-29-93 | 64 | 1000 | 25.0 | 7.49 | 7.75 | 40.8 | 85.5 | 192 | 0.00 | 0.00 | 0.00035 |
| 04-30-93 | 65 | 1000 | 25.3 | 7.48 | 8.09 | | | | | | |
| 05-01-93 | 66 | 1000 | 25.5 | 7.45 | 8.31 | | | | | | |
| 05-02-93 | 67 | 1000 | 25.6 | 7.44 | 7.97 | | | | | | |
| 05-03-93 | 68 | 1000 | 25.6 | 7.42 | 8.06 | | | | | | |
| 05-04-93 | 69 | 1000 | 25.8 | 7.45 | 7.51 | | | | | | |
| 05-05-93 | 70 | 900 | 25.7 | 7.39 | 7.49 | | | | | | |
| 05-06-93 | 71 | 1000 | 25.2 | 7.40 | 7.44 | 40.8 | 85.5 | 198 | 0.00 | 0.00 | 0.00030 |
| 05-07-93 | 72 | 900 | 24.9 | 7.42 | 7.59 | | | | | | |
| 05-08-93 | 73 | 1640 | 25.3 | 7.46 | 5.33 | | | | | | |
| 05-09-93 | 74 | 1440 | 25.2 | 7.45 | 5.56 | | | | | | |
| 05-10-93 | 75 | 1000 | 24.9 | 7.40 | 7.13 | | | | | | |
| 05-11-93 | 76 | 900 | 25.2 | 7.45 | 7.70 | | | | | | |
| 05-12-93 | 77 | 1000 | 25.3 | 7.40 | 7.11 | | | | | | |
| 05-13-93 | 78 | 1000 | 25.2 | 7.44 | 7.20 | 40.8 | 85.5 | 195 | 0.00 | 0.00 | 0.00000 |
| 05-14-93 | 79 | 800 | 24.9 | 7.51 | 7.19 | | | | | | |
| 05-15-93 | 80 | 800 | 25.0 | 7.49 | 7.56 | | | | | | |
| 05-16-93 | 81 | 1100 | 25.2 | 7.47 | 7.76 | | | | | | |
| 05-17-93 | 82 | 900 | 25.0 | 7.35 | 7.67 | | | | | | |
| 05-18-93 | 83 | 1000 | 25.0 | 7.37 | 7.75 | | | | | | |
| 05-19-93 | 84 | 1000 | 25.0 | 7.49 | 7.41 | 40.8 | 102.6 | 204 | 0.00 | 0.00 | 0.00086 |
| 05-20-93 | 85 | 900 | 25.2 | 7.45 | 7.23 | | | | | | |
| 05-21-93 | 86 | 900 | 24.9 | 7.41 | 7.28 | | | | | | |
| 05-22-93 | 87 | 800 | 24.7 | 7.56 | 7.52 | | | | | | |
| 05-23-93 | 88 | 1000 | 24.6 | 7.46 | 7.41 | | | | | | |
| 05-24-93 | 89 | 800 | 24.7 | 7.45 | 7.31 | | | | | | |
| 05-25-93 | 90 | 1000 | 24.9 | 7.51 | 7.03 | | | | | | |
| 05-26-93 | 91 | 900 | 24.9 | 7.54 | 7.41 | | | | | | |
| 05-27-93 | 92 | 900 | 24.8 | 7.44 | 7.24 | 40.8 | 85.5 | 190 | 0.00 | 0.00 | 0.00085 |
| 05-28-93 | 93 | 1000 | 24.9 | 7.48 | 7.38 | | | | | | |
| 05-29-93 | 94 | 1200 | 25.1 | 7.42 | 7.54 | | | | | | |
| 05-30-93 | 95 | 900 | 24.9 | 7.44 | 6.96 | | | | | | |

| | | | | | | | | | | | |
|----------|-----|------|------|------|------|------|------|-----|------|-------|---------|
| 09-16-93 | 204 | 930 | 25.2 | 7.40 | 6.74 | 40.8 | 85.5 | 203 | 0.00 | 0.000 | 0.00000 |
| 09-17-93 | 205 | 930 | 24.6 | 7.30 | 6.73 | | | | | | |
| 09-18-93 | 206 | 800 | 24.7 | 7.44 | 7.13 | | | | | | |
| 09-19-93 | 207 | 1300 | 24.5 | 7.39 | 7.36 | | | | | | |
| 09-20-93 | 208 | 930 | 24.4 | 7.36 | 7.33 | | | | | | |
| 09-21-93 | 209 | 930 | 26.0 | 7.45 | 6.52 | | | | | | |
| 09-22-93 | 210 | 900 | 25.3 | 7.27 | 6.79 | | | | | | |
| 09-23-93 | 211 | 930 | 25.1 | 7.35 | 7.10 | 40.8 | 85.5 | 209 | 0.00 | 0.000 | 0.00000 |
| 09-24-93 | 212 | 1000 | 25.3 | 7.39 | 6.96 | | | | | | |
| 09-25-93 | 213 | 900 | 25.1 | 7.39 | 7.25 | | | | | | |
| 09-26-93 | 214 | 2100 | 25.7 | 7.52 | 7.37 | | | | | | |
| 09-27-93 | 215 | 1000 | 25.5 | 7.44 | 7.01 | | | | | | |
| 09-28-93 | 216 | 900 | 24.8 | 7.40 | 7.28 | | | | | | |
| 09-29-93 | 217 | 930 | 24.7 | 7.41 | 7.28 | 40.8 | 85.5 | 209 | 0.00 | 0.008 | 0.00011 |
| 09-30-93 | 218 | 1300 | 24.6 | 7.23 | 7.24 | | | | | | |
| 10-01-93 | 219 | 1100 | 24.1 | 7.32 | 8.73 | | | | | | |
| 10-02-93 | 220 | 830 | 24.6 | 7.26 | 7.17 | | | | | | |
| 10-03-93 | 221 | 830 | 24.7 | 7.28 | 7.05 | | | | | | |
| 10-04-93 | 222 | 1000 | 24.4 | 7.35 | 7.37 | | | | | | |
| 10-05-93 | 223 | 1000 | 24.9 | 7.37 | 7.13 | | | | | | |
| 10-06-93 | 224 | 930 | 24.4 | 7.44 | 7.41 | | | | | | |
| 10-07-93 | 225 | 930 | 24.9 | 7.48 | 7.23 | 47.6 | 85.5 | 206 | 0.00 | 0.000 | 0.00000 |
| 10-08-93 | 226 | 1000 | 25.0 | 7.40 | 7.27 | | | | | | |
| 10-09-93 | 227 | 1130 | 25.0 | 7.46 | 7.31 | | | | | | |
| 10-10-93 | 228 | 1130 | 24.7 | 7.46 | 7.49 | | | | | | |
| 10-11-93 | 229 | 1600 | 24.6 | 7.47 | 7.48 | | | | | | |
| 10-12-93 | 230 | 1000 | 24.5 | 7.37 | 7.42 | | | | | | |
| 10-13-93 | 231 | 930 | 24.7 | 7.41 | 7.27 | | | | | | |
| 10-14-93 | 232 | 1000 | 24.7 | 7.38 | 7.24 | 47.6 | 85.5 | 205 | 0.00 | 0.000 | 0.00000 |
| 10-15-93 | 233 | 1000 | 24.8 | 7.28 | 7.26 | | | | | | |
| 10-16-93 | 234 | 1230 | 24.7 | 7.30 | 7.34 | | | | | | |
| 10-17-93 | 235 | 1030 | 24.8 | 7.40 | 7.21 | | | | | | |
| 10-18-93 | 236 | 1000 | 24.7 | 7.33 | 7.15 | | | | | | |
| 10-19-93 | 237 | 900 | 24.8 | 7.29 | 7.19 | | | | | | |
| 10-20-93 | 238 | 1000 | 24.8 | 7.38 | 7.41 | | | | | | |
| 10-21-93 | 239 | 1000 | 24.8 | 7.39 | 7.20 | 47.6 | 85.5 | 206 | 0.00 | 0.012 | 0.00016 |
| 10-22-93 | 240 | 800 | 24.6 | 7.29 | 7.11 | | | | | | |
| 10-23-93 | 241 | 700 | 24.3 | 7.23 | 7.57 | | | | | | |
| 10-24-93 | 242 | 800 | 24.4 | 7.19 | 7.28 | | | | | | |
| 10-25-93 | 243 | 1000 | 24.5 | 7.23 | 6.81 | | | | | | |
| 10-26-93 | 244 | 1000 | 24.6 | 7.29 | 7.30 | | | | | | |
| 10-27-93 | 245 | 1000 | 24.5 | 7.34 | 7.31 | | | | | | |
| 10-28-93 | 246 | 1000 | 24.9 | 7.43 | 7.11 | 47.6 | 85.5 | 199 | 0.00 | 0.015 | 0.00018 |
| 10-29-93 | 247 | 1230 | 24.7 | 7.34 | 7.20 | | | | | | |
| 10-30-93 | 248 | 700 | 25.1 | 7.41 | 7.18 | | | | | | |
| 10-31-93 | 249 | 800 | 24.8 | 7.49 | 7.08 | | | | | | |
| 11-01-93 | 250 | 930 | 24.5 | 7.31 | 7.08 | | | | | | |
| 11-02-93 | 251 | 930 | 24.8 | 7.37 | 7.10 | | | | | | |
| 11-03-93 | 252 | 1000 | 25.1 | 7.32 | 7.00 | | | | | | |
| 11-04-93 | 253 | 1000 | 25.0 | 7.26 | 7.00 | 47.6 | 85.5 | 206 | 0.00 | 0.000 | 0.00000 |
| 11-05-93 | 254 | 1000 | 25.2 | 7.34 | 6.87 | | | | | | |
| 11-06-93 | 255 | 600 | 25.3 | 7.22 | 6.84 | | | | | | |
| 11-07-93 | 256 | 1000 | 25.0 | 7.28 | 6.73 | | | | | | |
| 11-08-93 | 257 | 1000 | 24.8 | 7.28 | 6.94 | | | | | | |

TANK No. 2
TANK CONCENTRATION: 100% DILUENT WATER
NO DEN EXPOSURE

| Date | Test Day | Time (Military) | Temperature (Celsius) | pH | Dissolved Oxygen (mg/L) | Alkalinity (mg/L as CaCO3) | Hardness (mg/L as CaCO3) | Conductivity (umohs/cm) | Total Residual Chlorine (mg/L) | Free Available Chlorine (mg/L) | Total Ammonia- Nitrogen (mg/L) | Total Unionized Ammonia- Nitrogen (mg/L) |
|----------|----------|--------------------|--------------------------|------|-------------------------------|----------------------------------|--------------------------------|----------------------------|---|---|---|--|
| 02-25-93 | 1 | 1500 | 24.8 | 6.68 | 7.60 | | | | | | | |
| 02-26-93 | 2 | 1000 | 24.2 | 7.16 | 7.59 | | | | | | | |
| 02-27-93 | 3 | 1000 | 24.5 | 7.11 | 7.54 | | | | | | | |
| 02-28-93 | 4 | 1100 | 24.7 | 7.09 | 7.32 | 34.0 | 85.5 | | | | | |
| 03-01-93 | 5 | 930 | 24.9 | 7.09 | 7.26 | | | | | | | |
| 03-02-93 | 6 | 1000 | 25.5 | 7.06 | 7.09 | | | | | | | |
| 03-03-93 | 7 | 1000 | 26.0 | 7.32 | 7.64 | | | | | | | |
| 03-04-93 | 8 | 1000 | 25.3 | 7.45 | 7.75 | | | | | | | |
| 03-05-93 | 9 | 800 | 25.5 | 7.42 | 7.52 | | | | | | | |
| 03-06-93 | 10 | 1000 | 24.9 | 7.28 | 7.79 | 34.0 | 85.5 | 291 | | | | |
| 03-07-93 | 11 | 1100 | 25.0 | 7.09 | 7.75 | | | | | | | |
| 03-08-93 | 12 | 900 | 25.0 | 7.05 | 7.71 | | | | | | | |
| 03-09-93 | 13 | 930 | 25.3 | 7.19 | 7.61 | | | | | | | |
| 03-10-93 | 14 | 900 | 24.9 | 7.07 | 7.73 | | | | | | | |
| 03-11-93 | 15 | 900 | 25.2 | 7.46 | 7.64 | 54.4 | 85.5 | 196 | 0.00 | 0.00 | | |
| 03-12-93 | 16 | 800 | 24.6 | 7.42 | 8.12 | | | | | | | |
| 03-13-93 | 17 | 1600 | 24.8 | 7.17 | 7.83 | | | | | | | |
| 03-14-93 | 18 | 1500 | 24.9 | 7.18 | 7.82 | | | | | | | |
| 03-15-93 | 19 | 900 | 24.7 | 7.11 | 8.04 | | | | | | | |
| 03-16-93 | 20 | 900 | 25.3 | 7.21 | 7.85 | | | | | | | |
| 03-17-93 | 21 | 900 | 25.5 | 7.23 | 7.48 | 34.0 | 85.5 | 408 | 0.00 | 0.00 | 0.017 | 0.00017 |
| 03-18-93 | 22 | 1030 | 25.3 | 7.05 | 7.86 | | | | | | | |
| 03-19-93 | 23 | 1000 | 25.2 | 7.02 | 7.75 | | | | | | | |
| 03-20-93 | 24 | 1100 | 27.0 | 7.15 | 7.67 | | | | | | | |
| 03-21-93 | 25 | 1230 | 25.5 | 7.15 | 8.04 | | | | | | | |
| 03-22-93 | 26 | 900 | 25.0 | 7.22 | 8.04 | | | | | | | |
| 03-23-93 | 27 | 900 | 24.7 | 7.16 | 8.05 | | | | | | | |
| 03-24-93 | 28 | 900 | 25.3 | 7.05 | 7.82 | | | | | | | |
| 03-25-93 | 29 | 900 | 25.2 | 7.00 | 7.81 | 34.0 | 68.4 | 195 | 0.00 | 0.00 | 0.0000 | 0.00000 |
| 03-26-93 | 30 | 800 | 24.9 | 7.21 | 8.09 | | | | | | | |
| 03-27-93 | 31 | 1000 | 25.4 | 7.24 | 7.74 | | | | | | | |
| 03-28-93 | 32 | 1000 | 25.7 | 7.36 | 7.72 | | | | | | | |
| 03-29-93 | 33 | 1100 | 25.9 | 7.37 | 7.78 | | | | | | | |
| 03-30-93 | 34 | 1100 | 25.6 | 7.28 | 7.73 | | | | | | | |
| 03-31-93 | 35 | 1000 | 25.6 | 7.36 | 6.29 | | | | | | | |
| 04-01-93 | 36 | 1000 | 24.9 | 7.47 | 8.12 | 40.8 | 68.4 | 188 | 0.00 | 0.00 | 0.0000 | 0.00000 |
| 04-02-93 | 37 | 1000 | 24.9 | 7.42 | 8.11 | | | | | | | |
| 04-03-93 | 38 | 1100 | 24.9 | 7.46 | 8.23 | | | | | | | |
| 04-04-93 | 39 | 1300 | 24.7 | 7.43 | 8.36 | | | | | | | |
| 04-05-93 | 40 | 900 | 24.8 | 7.40 | 8.34 | | | | | | | |
| 04-06-93 | 41 | 1100 | 24.4 | 7.44 | 8.23 | | | | | | | |

| | | | | | | | | | | | |
|----------|----|------|------|------|------|------|-------|-----|------|-------|---------|
| 04-07-93 | 42 | 900 | 25.0 | 7.52 | 8.49 | 40.8 | 85.5 | 189 | 0.00 | 0.00 | 0.00000 |
| 04-08-93 | 43 | 900 | 25.3 | 7.51 | 8.25 | | | | | | |
| 04-09-93 | 44 | 900 | 25.3 | 7.52 | 8.07 | | | | | | |
| 04-10-93 | 45 | 1000 | 25.7 | 7.50 | 7.95 | | | | | | |
| 04-11-93 | 46 | 800 | 24.8 | 7.50 | 8.43 | | | | | | |
| 04-12-93 | 47 | 1000 | 25.2 | 7.47 | 8.16 | | | | | | |
| 04-13-93 | 48 | 900 | 25.0 | 7.50 | 7.96 | | | | | | |
| 04-14-93 | 49 | 600 | 25.1 | 7.50 | 7.65 | | | | | | |
| 04-15-93 | 50 | 900 | 25.4 | 7.50 | 7.62 | | | | | | |
| 04-16-93 | 51 | 1000 | 25.7 | 7.50 | 8.59 | | | | | | |
| 04-17-93 | 52 | 1000 | 25.3 | 7.50 | 7.68 | | | | | | |
| 04-18-93 | 53 | 1000 | 24.8 | 7.50 | 7.87 | | | | | | |
| 04-19-93 | 54 | 900 | 25.1 | 7.52 | 7.89 | | | | | | |
| 04-20-93 | 55 | 900 | 25.6 | 7.46 | 7.79 | | | | | | |
| 04-21-93 | 56 | 1000 | 25.7 | 7.50 | 7.63 | | | | | | |
| 04-22-93 | 57 | 900 | 25.0 | 7.53 | | 40.8 | 85.5 | 188 | 0.00 | 0.000 | 0.00000 |
| 04-23-93 | 58 | 900 | 24.9 | 7.51 | 7.40 | | | | | | |
| 04-24-93 | 59 | 1000 | 25.1 | 7.48 | 8.00 | | | | | | |
| 04-25-93 | 60 | 800 | 25.4 | 7.48 | 8.14 | | | | | | |
| 04-26-93 | 61 | 900 | 25.4 | 7.43 | 7.44 | | | | | | |
| 04-27-93 | 62 | 1000 | 24.4 | 7.54 | 8.06 | | | | | | |
| 04-28-93 | 63 | 900 | 25.0 | 7.40 | 7.58 | | | | | | |
| 04-29-93 | 64 | 1000 | 25.0 | 7.49 | 7.76 | 40.8 | 85.5 | 193 | 0.00 | 0.017 | 0.00029 |
| 04-30-93 | 65 | 1000 | 25.3 | 7.48 | 7.95 | | | | | | |
| 05-01-93 | 66 | 1000 | 25.5 | 7.45 | 8.18 | | | | | | |
| 05-02-93 | 67 | 1000 | 25.6 | 7.44 | 7.94 | | | | | | |
| 05-03-93 | 68 | 1000 | 25.6 | 7.42 | 7.94 | | | | | | |
| 05-04-93 | 69 | 1000 | 25.8 | 7.45 | 7.45 | | | | | | |
| 05-05-93 | 70 | 900 | 25.7 | 7.36 | 7.13 | | | | | | |
| 05-06-93 | 71 | 1000 | 25.2 | 7.38 | 7.19 | 40.8 | 85.5 | 197 | 0.00 | 0.027 | 0.00037 |
| 05-07-93 | 72 | 900 | 24.9 | 7.42 | 7.41 | | | | | | |
| 05-08-93 | 73 | 1640 | 25.3 | 7.39 | 5.40 | | | | | | |
| 05-09-93 | 74 | 1440 | 25.2 | 7.44 | 5.22 | | | | | | |
| 05-10-93 | 75 | 1000 | 24.9 | 7.40 | 7.04 | | | | | | |
| 05-11-93 | 76 | 900 | 25.2 | 7.45 | 7.58 | | | | | | |
| 05-12-93 | 77 | 1000 | 25.3 | 7.40 | 7.08 | 40.8 | 85.5 | 194 | 0.00 | 0.000 | 0.00000 |
| 05-13-93 | 78 | 1000 | 25.2 | 7.44 | 7.16 | | | | | | |
| 05-14-93 | 79 | 800 | 24.9 | 7.51 | 7.15 | | | | | | |
| 05-15-93 | 80 | 800 | 25.0 | 7.49 | 7.54 | | | | | | |
| 05-16-93 | 81 | 1100 | 25.2 | 7.47 | 7.75 | | | | | | |
| 05-17-93 | 82 | 900 | 25.0 | 7.36 | 7.64 | | | | | | |
| 05-18-93 | 83 | 1000 | 25.0 | 7.37 | 7.75 | | | | | | |
| 05-19-93 | 84 | 1000 | 25.0 | 7.49 | 7.40 | | | | | | |
| 05-20-93 | 85 | 900 | 25.2 | 7.45 | 7.21 | 40.8 | 102.6 | 204 | 0.00 | 0.048 | 0.00077 |
| 05-21-93 | 86 | 900 | 25.0 | 7.42 | 7.26 | | | | | | |
| 05-22-93 | 87 | 800 | 24.7 | 7.56 | 7.48 | | | | | | |
| 05-23-93 | 88 | 1000 | 24.6 | 7.46 | 7.38 | | | | | | |
| 05-24-93 | 89 | 800 | 24.8 | 7.45 | 7.27 | | | | | | |
| 05-25-93 | 90 | 1000 | 24.9 | 7.51 | 7.01 | | | | | | |
| 05-26-93 | 91 | 900 | 24.9 | 7.54 | 7.40 | | | | | | |
| 05-27-93 | 92 | 900 | 24.8 | 7.44 | 7.21 | 40.8 | 85.5 | 190 | 0.00 | 0.053 | 0.00081 |
| 05-28-93 | 93 | 1000 | 24.9 | 7.48 | 7.38 | | | | | | |
| 05-29-93 | 94 | 1200 | 25.1 | 7.42 | 7.44 | | | | | | |
| 05-30-93 | 95 | 900 | 24.9 | 7.44 | 6.81 | | | | | | |

| | | | | | | | | | | | |
|----------|-----|------|------|------|------|------|------|-----|------|------|---------|
| 09-16-93 | 204 | 930 | 25.2 | 7.40 | 6.71 | 40.8 | 85.5 | 203 | 0.00 | 0.00 | 0.00000 |
| 09-17-93 | 205 | 930 | 24.6 | 7.30 | 6.96 | | | | | | |
| 09-18-93 | 206 | 800 | 24.7 | 7.44 | 7.01 | | | | | | |
| 09-19-93 | 207 | 1300 | 24.6 | 7.39 | 7.30 | | | | | | |
| 09-20-93 | 208 | 930 | 24.5 | 7.36 | 7.29 | | | | | | |
| 09-21-93 | 209 | 930 | 26.0 | 7.45 | 6.52 | | | | | | |
| 09-22-93 | 210 | 900 | 25.4 | 7.27 | 6.75 | | | | | | |
| 09-23-93 | 211 | 930 | 25.1 | 7.35 | 6.91 | 40.8 | 85.5 | 209 | 0.00 | 0.00 | 0.00000 |
| 09-24-93 | 212 | 1000 | 25.3 | 7.39 | 6.74 | | | | | | |
| 09-25-93 | 213 | 900 | 25.1 | 7.39 | 7.14 | | | | | | |
| 09-26-93 | 214 | 2100 | 25.7 | 7.52 | 7.38 | | | | | | |
| 09-27-93 | 215 | 1000 | 25.4 | 7.44 | 6.92 | | | | | | |
| 09-28-93 | 216 | 900 | 24.8 | 7.40 | 7.19 | | | | | | |
| 09-29-93 | 217 | 930 | 24.8 | 7.41 | 7.02 | 40.8 | 85.5 | 209 | 0.00 | 0.00 | 0.00014 |
| 09-30-93 | 218 | 1300 | 24.6 | 7.27 | 7.11 | | | | | | |
| 10-01-93 | 219 | 1100 | 24.1 | 7.28 | 8.38 | | | | | | |
| 10-02-93 | 220 | 830 | 24.7 | 7.30 | 7.18 | | | | | | |
| 10-03-93 | 221 | 830 | 24.8 | 7.30 | 6.66 | | | | | | |
| 10-04-93 | 222 | 1000 | 24.4 | 7.31 | 7.09 | | | | | | |
| 10-05-93 | 223 | 1000 | 24.9 | 7.33 | 6.90 | | | | | | |
| 10-06-93 | 224 | 930 | 24.4 | 7.42 | 7.26 | | | | | | |
| 10-07-93 | 225 | 930 | 24.9 | 7.46 | 7.09 | 47.6 | 85.5 | 206 | 0.00 | 0.00 | 0.00000 |
| 10-08-93 | 226 | 1000 | 25.0 | 7.36 | 7.11 | | | | | | |
| 10-09-93 | 227 | 1130 | 25.0 | 7.42 | 7.22 | | | | | | |
| 10-10-93 | 228 | 1130 | 24.7 | 7.46 | 7.37 | | | | | | |
| 10-11-93 | 229 | 1600 | 24.6 | 7.47 | 7.50 | | | | | | |
| 10-12-93 | 230 | 1000 | 24.5 | 7.36 | 7.41 | | | | | | |
| 10-13-93 | 231 | 930 | 24.7 | 7.41 | 7.27 | | | | | | |
| 10-14-93 | 232 | 1000 | 24.7 | 7.36 | 7.16 | 47.6 | 85.5 | 205 | 0.00 | 0.00 | 0.00000 |
| 10-15-93 | 233 | 1000 | 24.8 | 7.27 | 7.23 | | | | | | |
| 10-16-93 | 234 | 1230 | 24.7 | 7.28 | 7.16 | | | | | | |
| 10-17-93 | 235 | 1030 | 24.8 | 7.40 | 7.16 | | | | | | |
| 10-18-93 | 236 | 1000 | 24.7 | 7.33 | 7.15 | | | | | | |
| 10-19-93 | 237 | 900 | 24.7 | 7.28 | 7.25 | | | | | | |
| 10-20-93 | 238 | 1000 | 24.8 | 7.38 | 7.42 | | | | | | |
| 10-21-93 | 239 | 1000 | 24.8 | 7.39 | 7.29 | 47.6 | 85.5 | 206 | 0.00 | 0.00 | 0.00000 |
| 10-22-93 | 240 | 800 | 24.6 | 7.27 | 7.01 | | | | | | |
| 10-23-93 | 241 | 700 | 24.4 | 7.29 | 7.39 | | | | | | |
| 10-24-93 | 242 | 800 | 24.4 | 7.18 | 7.21 | | | | | | |
| 10-25-93 | 243 | 1000 | 24.4 | 7.27 | 6.73 | | | | | | |
| 10-26-93 | 244 | 1000 | 24.6 | 7.29 | 7.32 | | | | | | |
| 10-27-93 | 245 | 1000 | 24.4 | 7.34 | 7.36 | | | | | | |
| 10-28-93 | 246 | 1000 | 24.9 | 7.43 | 7.09 | 47.6 | 85.5 | 199 | 0.00 | 0.00 | 0.00020 |
| 10-29-93 | 247 | 1230 | 24.7 | 7.34 | 7.16 | | | | | | |
| 10-30-93 | 248 | 700 | 25.1 | 7.41 | 7.21 | | | | | | |
| 10-31-93 | 249 | 800 | 24.7 | 7.49 | 7.14 | | | | | | |
| 11-01-93 | 250 | 930 | 24.5 | 7.31 | 7.12 | | | | | | |
| 11-02-93 | 251 | 930 | 24.8 | 7.37 | 7.13 | | | | | | |
| 11-03-93 | 252 | 1000 | 25.1 | 7.32 | 7.01 | | | | | | |
| 11-04-93 | 253 | 1000 | 25.0 | 7.26 | 7.03 | 47.6 | 85.5 | 206 | 0.00 | 0.00 | 0.00000 |
| 11-05-93 | 254 | 1000 | 25.2 | 7.34 | 6.89 | | | | | | |
| 11-06-93 | 255 | 600 | 25.3 | 7.22 | 7.02 | | | | | | |
| 11-07-93 | 256 | 1000 | 25.0 | 7.28 | 6.86 | | | | | | |
| 11-08-93 | 257 | 1000 | 24.8 | 7.28 | 7.05 | | | | | | |

TANK No. 3
TANK CONCENTRATION: 100% DILUENT WATER
DEN EXPOSURE: 10 mg/L

| Date | Test Day | Time (Military) | Temperature (Celsius) | pH | Dissolved Oxygen (mg/L) | Alkalinity (mg/L as CaCO ₃) | Hardness (mg/L as CaCO ₃) | Conductivity (umohs/cm) | Total Residual Chlorine (mg/L) | Free Available Chlorine (mg/L) | Total Ammonia- Nitrogen (mg/L) | Total Unionized Ammonia- Nitrogen (mg/L) |
|----------|----------|--------------------|--------------------------|------|-------------------------------|---|---|----------------------------|---|---|---|--|
| 02-25-93 | 1 | 1500 | 24.8 | 6.74 | 7.66 | | | | | | | |
| 02-26-93 | 2 | 1000 | 24.1 | 7.16 | 7.68 | | | | | | | |
| 02-27-93 | 3 | 1000 | 24.4 | 7.12 | 7.40 | | | | | | | |
| 02-28-93 | 4 | 1100 | 24.7 | 7.10 | 7.47 | 34.0 | 85.5 | | | | | |
| 03-01-93 | 5 | 930 | 25.0 | 7.09 | 7.24 | | | | | | | |
| 03-02-93 | 6 | 1000 | 25.5 | 7.05 | 7.09 | | | | | | | |
| 03-03-93 | 7 | 1000 | 26.0 | 7.32 | 7.78 | | | | | | | |
| 03-04-93 | 8 | 1000 | 25.2 | 7.45 | 7.68 | | | | | | | |
| 03-05-93 | 9 | 800 | 25.5 | 7.43 | 7.46 | | | | | | | |
| 03-06-93 | 10 | 1000 | 24.9 | 7.29 | 7.78 | 34.0 | 85.5 | 292 | | | | |
| 03-07-93 | 11 | 1100 | 25.0 | 7.10 | 7.79 | | | | | | | |
| 03-08-93 | 12 | 900 | 25.0 | 7.05 | 7.64 | | | | | | | |
| 03-09-93 | 13 | 930 | 25.3 | 7.21 | 7.57 | | | | | | | |
| 03-10-93 | 14 | 900 | 24.9 | 7.08 | 7.65 | | | | | | | |
| 03-11-93 | 15 | 900 | 25.2 | 7.46 | 7.53 | | | | | | | |
| 03-12-93 | 16 | 800 | 24.5 | 7.42 | 8.03 | 54.4 | 85.5 | 194 | 0.00 | 0.00 | | |
| 03-13-93 | 17 | 1600 | 24.8 | 7.17 | 7.78 | | | | | | | |
| 03-14-93 | 18 | 1500 | 24.9 | 7.17 | 7.75 | | | | | | | |
| 03-15-93 | 19 | 900 | 24.7 | 7.12 | 7.90 | | | | | | | |
| 03-16-93 | 20 | 900 | 25.3 | 7.23 | 7.78 | | | | | | | |
| 03-17-93 | 21 | 900 | 25.5 | 7.23 | 7.40 | 34.0 | 85.5 | 409 | 0.00 | 0.00 | 0.021 | 0.00021 |
| 03-18-93 | 22 | 1030 | 25.3 | 7.05 | 7.67 | | | | | | | |
| 03-19-93 | 23 | 1000 | 25.2 | 7.07 | 7.75 | | | | | | | |
| 03-20-93 | 24 | 1100 | 27.0 | 7.16 | 7.59 | | | | | | | |
| 03-21-93 | 25 | 1230 | 25.5 | 7.15 | 7.98 | | | | | | | |
| 03-22-93 | 26 | 900 | 25.0 | 7.22 | 7.93 | | | | | | | |
| 03-23-93 | 27 | 900 | 25.0 | 7.16 | 7.78 | | | | | | | |
| 03-24-93 | 28 | 900 | 25.3 | 7.05 | 7.73 | | | | | | | |
| 03-25-93 | 29 | 900 | 25.2 | 7.00 | 7.72 | 34.0 | 68.4 | 194 | 0.00 | 0.00 | 0.000 | 0.00000 |
| 03-26-93 | 30 | 800 | 24.9 | 7.21 | 8.04 | | | | | | | |
| 03-27-93 | 31 | 1000 | 25.4 | 7.24 | 7.71 | | | | | | | |
| 03-28-93 | 32 | 1000 | 25.7 | 7.36 | 7.60 | | | | | | | |
| 03-29-93 | 33 | 1100 | 25.9 | 7.37 | 7.64 | | | | | | | |
| 03-30-93 | 34 | 1100 | 25.6 | 7.30 | 7.72 | | | | | | | |
| 03-31-93 | 35 | 1000 | 25.6 | 7.36 | 6.08 | | | | | | | |
| 04-01-93 | 36 | 1000 | 24.8 | 7.45 | 7.97 | 40.8 | 68.4 | 188 | 0.00 | 0.00 | 0.000 | 0.00000 |
| 04-02-93 | 37 | 1000 | 24.9 | 7.41 | 7.97 | | | | | | | |
| 04-03-93 | 38 | 1100 | 24.9 | 7.44 | 8.01 | | | | | | | |
| 04-04-93 | 39 | 1300 | 24.8 | 7.42 | 8.18 | | | | | | | |
| 04-05-93 | 40 | 900 | 24.8 | 7.42 | 8.25 | | | | | | | |
| 04-06-93 | 41 | 1100 | 24.4 | 7.44 | 8.11 | | | | | | | |

| | | | | | | | | | | | |
|----------|----|------|------|------|------|------|------|-----|------|-------|---------|
| 04-07-93 | 42 | 900 | 24.9 | 7.50 | 8.35 | 40.8 | 85.5 | 189 | 0.00 | 0.00 | 0.00000 |
| 04-08-93 | 43 | 900 | 25.3 | 7.50 | 8.16 | | | | | | |
| 04-09-93 | 44 | 900 | 25.3 | 7.52 | 8.01 | | | | | | |
| 04-10-93 | 45 | 1000 | 25.7 | 7.47 | 7.85 | | | | | | |
| 04-11-93 | 46 | 800 | 24.8 | 7.50 | 8.33 | | | | | | |
| 04-12-93 | 47 | 1000 | 25.2 | 7.45 | 7.94 | | | | | | |
| 04-13-93 | 48 | 900 | 25.0 | 7.50 | 7.83 | | | | | | |
| 04-14-93 | 49 | 600 | 25.1 | 7.49 | 7.68 | | | | | | |
| 04-15-93 | 50 | 900 | 25.4 | 7.47 | 7.51 | 40.8 | 85.5 | 188 | 0.00 | 0.024 | 0.00045 |
| 04-16-93 | 51 | 1000 | 25.7 | 7.48 | 8.42 | | | | | | |
| 04-17-93 | 52 | 1000 | 25.3 | 7.47 | 7.55 | | | | | | |
| 04-18-93 | 53 | 1000 | 24.8 | 7.51 | 7.71 | | | | | | |
| 04-19-93 | 54 | 900 | 25.1 | 7.52 | 7.77 | | | | | | |
| 04-20-93 | 55 | 900 | 25.6 | 7.46 | 7.63 | | | | | | |
| 04-21-93 | 56 | 1000 | 25.7 | 7.48 | 7.48 | | | | | | |
| 04-22-93 | 57 | 900 | 25.0 | 7.53 | 7.48 | 40.8 | 85.5 | 187 | 0.00 | 0.000 | 0.00000 |
| 04-23-93 | 58 | 900 | 24.9 | 7.51 | 7.29 | | | | | | |
| 04-24-93 | 59 | 1000 | 25.1 | 7.48 | 7.79 | | | | | | |
| 04-25-93 | 60 | 800 | 25.4 | 7.48 | 7.99 | | | | | | |
| 04-26-93 | 61 | 900 | 25.4 | 7.43 | 7.40 | | | | | | |
| 04-27-93 | 62 | 1000 | 24.4 | 7.54 | 7.97 | | | | | | |
| 04-28-93 | 63 | 900 | 25.0 | 7.36 | 7.55 | | | | | | |
| 04-29-93 | 64 | 1000 | 25.0 | 7.46 | 7.72 | 40.8 | 85.5 | 193 | 0.00 | 0.031 | 0.00050 |
| 04-30-93 | 65 | 1000 | 25.3 | 7.42 | 7.80 | | | | | | |
| 05-01-93 | 66 | 1000 | 25.5 | 7.39 | 7.72 | | | | | | |
| 05-02-93 | 67 | 1000 | 25.6 | 7.40 | 7.76 | | | | | | |
| 05-03-93 | 68 | 1000 | 25.6 | 7.38 | 7.88 | | | | | | |
| 05-04-93 | 69 | 1000 | 25.8 | 7.40 | 7.36 | | | | | | |
| 05-05-93 | 70 | 900 | 25.7 | 7.33 | 7.12 | | | | | | |
| 05-06-93 | 71 | 1000 | 25.2 | 7.33 | 7.03 | 40.8 | 85.5 | 198 | 0.00 | 0.032 | 0.00039 |
| 05-07-93 | 72 | 900 | 24.9 | 7.32 | 6.83 | | | | | | |
| 05-08-93 | 73 | 1640 | 25.2 | 7.18 | 5.45 | | | | | | |
| 05-09-93 | 74 | 1440 | 24.4 | 6.90 | 5.36 | | | | | | |
| 05-10-93 | 75 | 1000 | 24.6 | 6.15 | 6.37 | | | | | | |
| 05-11-93 | 76 | 900 | 25.2 | 7.44 | 7.43 | | | | | | |
| 05-12-93 | 77 | 1000 | 25.3 | 7.41 | 6.96 | | | | | | |
| 05-13-93 | 78 | 1000 | 25.2 | 7.44 | 6.98 | 40.8 | 85.5 | 194 | 0.00 | 0.000 | 0.00000 |
| 05-14-93 | 79 | 800 | 24.9 | 7.51 | 7.01 | | | | | | |
| 05-15-93 | 80 | 800 | 25.1 | 7.48 | 7.36 | | | | | | |
| 05-16-93 | 81 | 1100 | 25.3 | 7.47 | 7.58 | | | | | | |
| 05-17-93 | 82 | 900 | 25.1 | 7.34 | 7.48 | | | | | | |
| 05-18-93 | 83 | 1000 | 25.1 | 7.37 | 7.55 | | | | | | |
| 05-19-93 | 84 | 1000 | 25.0 | 7.49 | 7.23 | | | | | | |
| 05-20-93 | 85 | 900 | 25.2 | 7.43 | 6.94 | 40.8 | 85.5 | 203 | 0.00 | 0.058 | 0.00093 |
| 05-21-93 | 86 | 900 | 25.0 | 7.40 | 7.07 | | | | | | |
| 05-22-93 | 87 | 800 | 24.7 | 7.53 | 7.26 | | | | | | |
| 05-23-93 | 88 | 1000 | 24.6 | 7.45 | 7.20 | | | | | | |
| 05-24-93 | 89 | 800 | 24.8 | 7.42 | 7.01 | | | | | | |
| 05-25-93 | 90 | 1000 | 24.9 | 7.51 | 6.83 | | | | | | |
| 05-26-93 | 91 | 900 | 24.9 | 7.54 | 7.31 | | | | | | |
| 05-27-93 | 92 | 900 | 24.8 | 7.44 | 7.13 | 40.8 | 85.5 | 190 | 0.00 | 0.064 | 0.00097 |
| 05-28-93 | 93 | 1000 | 24.9 | 7.49 | 7.38 | | | | | | |
| 05-29-93 | 94 | 1200 | 25.1 | 7.42 | 7.43 | | | | | | |
| 05-30-93 | 95 | 900 | 24.9 | 7.45 | 6.94 | | | | | | |

| | | | | | |
|-----|----------|------|------|------|------|
| 150 | 07-24-93 | 900 | 25.1 | 7.12 | 6.42 |
| 151 | 07-25-93 | 1200 | 25.1 | 7.22 | 6.44 |
| 152 | 07-26-93 | 1000 | 25.0 | 7.23 | 6.85 |
| 153 | 07-27-93 | 800 | 25.0 | 7.25 | 6.47 |
| 154 | 07-28-93 | 1000 | 25.0 | 7.28 | 6.53 |
| 155 | 07-29-93 | 900 | 25.0 | 7.27 | 6.75 |
| 156 | 07-30-93 | 700 | 25.5 | 7.30 | 6.19 |
| 157 | 07-31-93 | 800 | 25.0 | 7.31 | 6.39 |
| 158 | 08-01-93 | 1500 | 25.1 | 7.34 | 6.88 |
| 159 | 08-02-93 | 900 | 25.4 | 7.30 | 6.45 |
| 160 | 08-03-93 | 900 | 25.4 | 7.30 | 6.76 |
| 161 | 08-04-93 | 1130 | 24.6 | 7.31 | 6.81 |
| 162 | 08-05-93 | 900 | 25.0 | 7.28 | 6.57 |
| 163 | 08-06-93 | 900 | 24.9 | 7.35 | 6.63 |
| 164 | 08-07-93 | 1300 | 24.7 | 7.32 | 6.80 |
| 165 | 08-08-93 | 700 | 24.9 | 7.35 | 6.63 |
| 166 | 08-09-93 | 900 | 24.9 | 7.16 | 6.54 |
| 167 | 08-10-93 | 900 | 24.9 | 7.24 | 6.48 |
| 168 | 08-11-93 | 1100 | 24.9 | 7.14 | 5.91 |
| 169 | 08-12-93 | 1100 | 24.8 | 7.28 | 6.59 |
| 170 | 08-13-93 | 1100 | 24.8 | 7.37 | 6.75 |
| 171 | 08-14-93 | 1000 | 25.0 | 7.12 | 6.16 |
| 172 | 08-15-93 | 900 | 25.0 | 7.14 | 6.44 |
| 173 | 08-16-93 | 1000 | 25.1 | 7.22 | 7.48 |
| 174 | 08-17-93 | 1300 | 25.1 | 7.08 | 5.72 |
| 175 | 08-18-93 | 1130 | 24.6 | 7.10 | 6.10 |
| 176 | 08-19-93 | 900 | 24.7 | 7.33 | 6.82 |
| 177 | 08-20-93 | 900 | 24.8 | 7.21 | 6.43 |
| 178 | 08-21-93 | 1200 | 24.8 | 7.15 | 6.92 |
| 179 | 08-22-93 | 830 | 24.8 | 7.15 | 6.49 |
| 180 | 08-23-93 | 500 | 24.9 | 7.28 | 6.94 |
| 181 | 08-24-93 | 830 | 24.9 | 7.34 | 7.42 |
| 182 | 08-25-93 | 830 | 25.0 | 7.40 | 6.74 |
| 183 | 08-26-93 | 730 | 25.0 | 7.32 | 6.75 |
| 184 | 08-27-93 | 830 | 25.0 | 7.38 | 6.86 |
| 185 | 08-28-93 | 915 | 25.0 | 7.36 | 6.91 |
| 186 | 08-29-93 | 905 | 24.9 | 7.45 | 6.79 |
| 187 | 08-30-93 | 900 | 24.9 | 7.42 | 7.07 |
| 188 | 08-31-93 | 900 | 25.0 | 7.44 | 6.94 |
| 189 | 09-01-93 | 930 | 25.0 | 7.45 | 6.81 |
| 190 | 09-02-93 | 900 | 25.0 | 7.46 | 7.01 |
| 191 | 09-03-93 | 1000 | 25.2 | 7.54 | 6.71 |
| 192 | 09-04-93 | 1100 | 25.1 | 7.43 | 6.75 |
| 193 | 09-05-93 | 830 | 25.0 | 7.45 | 6.70 |
| 194 | 09-06-93 | 1115 | 24.9 | 7.42 | 6.50 |
| 195 | 09-07-93 | 930 | 25.1 | 7.47 | 6.93 |
| 196 | 09-08-93 | 900 | 25.1 | 7.48 | 6.65 |
| 197 | 09-09-93 | 930 | 25.0 | 7.50 | 6.98 |
| 198 | 09-10-93 | 930 | 24.9 | 7.32 | 5.63 |
| 199 | 09-11-93 | 1600 | 24.7 | 7.45 | 7.31 |
| 200 | 09-12-93 | 730 | 24.5 | 7.45 | 7.15 |
| 201 | 09-13-93 | 930 | 25.4 | 7.41 | 6.85 |
| 202 | 09-14-93 | 930 | 25.3 | 7.51 | 6.82 |
| 203 | 09-15-93 | 930 | 25.2 | 7.40 | 6.84 |

| | | | | | | | | | | | |
|----------|-----|------|------|------|------|------|------|-----|------|------|---------|
| 09-16-93 | 204 | 930 | 25.2 | 7.40 | 6.73 | 40.8 | 85.5 | 203 | 0.00 | 0.00 | 0.0000 |
| 09-17-93 | 205 | 930 | 24.6 | 7.30 | 6.96 | | | | | | |
| 09-18-93 | 206 | 800 | 24.7 | 7.44 | 7.09 | | | | | | |
| 09-19-93 | 207 | 1300 | 24.5 | 7.39 | 7.47 | | | | | | |
| 09-20-93 | 208 | 930 | 24.4 | 7.36 | 7.36 | | | | | | |
| 09-21-93 | 209 | 930 | 26.0 | 7.45 | 6.86 | | | | | | |
| 09-22-93 | 210 | 900 | 25.3 | 7.27 | 6.95 | | | | | | |
| 09-23-93 | 211 | 930 | 25.1 | 7.35 | 7.14 | 40.8 | 85.5 | 208 | 0.00 | 0.00 | 0.00000 |
| 09-24-93 | 212 | 1000 | 25.3 | 7.39 | 7.00 | | | | | | |
| 09-25-93 | 213 | 900 | 25.1 | 7.39 | 7.30 | | | | | | |
| 09-26-93 | 214 | 2100 | 25.7 | 7.52 | 7.38 | | | | | | |
| 09-27-93 | 215 | 1000 | 25.4 | 7.44 | 7.08 | | | | | | |
| 09-28-93 | 216 | 900 | 24.8 | 7.40 | 7.24 | | | | | | |
| 09-29-93 | 217 | 930 | 24.8 | 7.41 | 7.28 | 40.8 | 85.5 | 209 | 0.00 | 0.00 | 0.00000 |
| 09-30-93 | 218 | 1300 | 24.6 | 7.29 | 7.23 | | | | | | |
| 10-01-93 | 219 | 1100 | 24.1 | 7.31 | 8.77 | | | | | | |
| 10-02-93 | 220 | 830 | 24.7 | 7.34 | 7.21 | | | | | | |
| 10-03-93 | 221 | 830 | 24.9 | 7.32 | 6.70 | | | | | | |
| 10-04-93 | 222 | 1000 | 24.4 | 7.35 | 7.36 | | | | | | |
| 10-05-93 | 223 | 1000 | 24.9 | 7.36 | 7.17 | | | | | | |
| 10-06-93 | 224 | 930 | 24.5 | 7.44 | 7.41 | | | | | | |
| 10-07-93 | 225 | 930 | 24.9 | 7.48 | 7.33 | 47.6 | 85.5 | 206 | 0.00 | 0.00 | 0.00000 |
| 10-08-93 | 226 | 1000 | 25.0 | 7.40 | 7.36 | | | | | | |
| 10-09-93 | 227 | 1130 | 24.9 | 7.46 | 7.41 | | | | | | |
| 10-10-93 | 228 | 1130 | 24.7 | 7.46 | 7.51 | | | | | | |
| 10-11-93 | 229 | 1600 | 24.6 | 7.47 | 7.52 | | | | | | |
| 10-12-93 | 230 | 1000 | 24.5 | 7.36 | 7.37 | | | | | | |
| 10-13-93 | 231 | 930 | 24.7 | 7.41 | 7.29 | | | | | | |
| 10-14-93 | 232 | 1000 | 24.6 | 7.38 | 7.31 | 47.6 | 85.5 | 205 | 0.00 | 0.00 | 0.00000 |
| 10-15-93 | 233 | 1000 | 24.7 | 7.27 | 7.37 | | | | | | |
| 10-16-93 | 234 | 1230 | 24.7 | 7.31 | 7.37 | | | | | | |
| 10-17-93 | 235 | 1030 | 24.8 | 7.40 | 7.21 | | | | | | |
| 10-18-93 | 236 | 1000 | 24.7 | 7.33 | 7.24 | | | | | | |
| 10-19-93 | 237 | 900 | 24.8 | 7.30 | 7.31 | | | | | | |
| 10-20-93 | 238 | 1000 | 24.8 | 7.38 | 7.30 | | | | | | |
| 10-21-93 | 239 | 1000 | 24.9 | 7.39 | 7.19 | | | | | | |
| 10-22-93 | 240 | 800 | 24.7 | 7.28 | 6.93 | | | | | | |
| 10-23-93 | 241 | 700 | 24.5 | 7.29 | 7.41 | | | | | | |
| 10-24-93 | 242 | 800 | 24.5 | 7.20 | 7.30 | | | | | | |
| 10-25-93 | 243 | 1000 | 24.5 | 7.27 | 6.54 | | | | | | |
| 10-26-93 | 244 | 1000 | 24.7 | 7.29 | 7.10 | | | | | | |
| 10-27-93 | 245 | 1000 | 24.5 | 7.34 | 7.21 | | | | | | |
| 10-28-93 | 246 | 1000 | 25.0 | 7.43 | 6.99 | 47.6 | 85.5 | 199 | 0.00 | 0.00 | 0.00030 |
| 10-29-93 | 247 | 1230 | 24.8 | 7.34 | 6.93 | | | | | | |
| 10-30-93 | 248 | 700 | 25.2 | 7.38 | 6.81 | | | | | | |
| 10-31-93 | 249 | 800 | 24.8 | 7.49 | 7.00 | | | | | | |
| 11-01-93 | 250 | 930 | 24.6 | 7.29 | 6.96 | | | | | | |
| 11-02-93 | 251 | 930 | 24.8 | 7.37 | 7.32 | | | | | | |
| 11-03-93 | 252 | 1000 | 25.1 | 7.34 | 7.17 | | | | | | |
| 11-04-93 | 253 | 1000 | 24.9 | 7.26 | 7.14 | 47.6 | 85.5 | 205 | 0.00 | 0.00 | 0.00000 |
| 11-05-93 | 254 | 1000 | 25.2 | 7.34 | 7.06 | | | | | | |
| 11-06-93 | 255 | 600 | 25.3 | 7.22 | 6.87 | | | | | | |
| 11-07-93 | 256 | 1000 | 25.0 | 7.30 | 6.92 | | | | | | |
| 11-08-93 | 257 | 1000 | 24.8 | 7.28 | 7.10 | | | | | | |

TANK No. 4
TANK CONCENTRATION: 100% DILUENT WATER
DEN EXPOSURE: 10 mg/L

| Date | Test Day | Time (Military) | Temperature (Celsius) | pH | Dissolved Oxygen (mg/L) | Alkalinity (mg/L as CaCO3) | Hardness (mg/L as CaCO3) | Conductivity (umohs/cm) | Total Residual Chlorine (mg/L) | Free Available Chlorine (mg/L) | Total Ammonia- Nitrogen (mg/L) | Total Unionized Ammonia- Nitrogen (mg/L) |
|----------|----------|--------------------|--------------------------|------|-------------------------------|----------------------------------|--------------------------------|----------------------------|---|---|---|--|
| 02-25-93 | 1 | 1500 | 24.8 | 6.74 | 7.63 | | | | | | | |
| 02-26-93 | 2 | 1000 | 24.2 | 7.16 | 7.58 | | | | | | | |
| 02-27-93 | 3 | 1000 | 24.5 | 7.13 | 7.33 | | | | | | | |
| 02-28-93 | 4 | 1100 | 24.9 | 7.10 | 7.37 | 34.0 | 85.5 | | | | | |
| 03-01-93 | 5 | 930 | 25.0 | 7.09 | 7.12 | | | | | | | |
| 03-02-93 | 6 | 1000 | 25.6 | 7.05 | 7.06 | | | | | | | |
| 03-03-93 | 7 | 1000 | 26.0 | 7.32 | 7.78 | | | | | | | |
| 03-04-93 | 8 | 1000 | 25.3 | 7.45 | 7.68 | | | | | | | |
| 03-05-93 | 9 | 800 | 25.6 | 7.44 | 7.48 | | | | | | | |
| 03-06-93 | 10 | 1000 | 24.9 | 7.30 | 7.74 | 34.0 | 85.5 | 292 | | | | |
| 03-07-93 | 11 | 1100 | 25.0 | 7.09 | 7.78 | | | | | | | |
| 03-08-93 | 12 | 900 | 25.0 | 7.06 | 7.67 | | | | | | | |
| 03-09-93 | 13 | 930 | 25.3 | 7.21 | 7.58 | | | | | | | |
| 03-10-93 | 14 | 900 | 25.0 | 7.08 | 7.65 | | | | | | | |
| 03-11-93 | 15 | 800 | 25.3 | 7.46 | 7.53 | | | | 0.00 | 0.00 | | |
| 03-12-93 | 16 | 800 | 24.6 | 7.41 | 8.02 | | | | | | | |
| 03-13-93 | 17 | 1600 | 24.8 | 7.18 | 7.77 | 54.4 | 85.5 | 193 | | | | |
| 03-14-93 | 18 | 1500 | 24.9 | 7.18 | 7.77 | | | | | | | |
| 03-15-93 | 19 | 900 | 24.7 | 7.12 | 7.94 | | | | | | | |
| 03-16-93 | 20 | 900 | 25.3 | 7.23 | 7.83 | 34.0 | 85.5 | 409 | 0.00 | 0.00 | 0.020 | 0.00020 |
| 03-17-93 | 21 | 900 | 25.5 | 7.23 | 7.46 | | | | | | | |
| 03-18-93 | 22 | 1030 | 25.5 | 7.05 | 7.59 | | | | | | | |
| 03-19-93 | 23 | 1000 | 25.3 | 7.07 | 7.33 | | | | | | | |
| 03-20-93 | 24 | 1100 | 27.1 | 7.16 | 7.61 | | | | | | | |
| 03-21-93 | 25 | 1230 | 25.5 | 7.15 | 7.96 | | | | | | | |
| 03-22-93 | 26 | 900 | 25.0 | 7.22 | 7.94 | | | | | | | |
| 03-23-93 | 27 | 900 | 25.0 | 7.16 | 7.86 | | | | | | | |
| 03-24-93 | 28 | 900 | 25.3 | 7.05 | 7.79 | | | | | | | |
| 03-25-93 | 29 | 900 | 25.2 | 7.00 | 7.76 | 34.0 | 68.4 | 195 | 0.00 | 0.00 | 0.000 | 0.00000 |
| 03-26-93 | 30 | 800 | 24.9 | 7.21 | 8.03 | | | | | | | |
| 03-27-93 | 31 | 1000 | 25.4 | 7.24 | 7.70 | | | | | | | |
| 03-28-93 | 32 | 1000 | 25.7 | 7.36 | 7.61 | | | | | | | |
| 03-29-93 | 33 | 1100 | 25.9 | 7.37 | 7.63 | | | | | | | |
| 03-30-93 | 34 | 1100 | 25.6 | 7.24 | 7.72 | | | | | | | |
| 03-31-93 | 35 | 1000 | 25.6 | 7.36 | 6.06 | | | | | | | |
| 04-01-93 | 36 | 1000 | 24.8 | 7.47 | 7.88 | 40.8 | 68.4 | 188 | 0.00 | 0.00 | 0.000 | 0.00000 |
| 04-02-93 | 37 | 1000 | 24.9 | 7.41 | 7.92 | | | | | | | |
| 04-03-93 | 38 | 1100 | 24.9 | 7.46 | 8.08 | | | | | | | |
| 04-04-93 | 39 | 1300 | 24.8 | 7.41 | 8.18 | | | | | | | |
| 04-05-93 | 40 | 900 | 24.8 | 7.44 | 8.31 | | | | | | | |
| 04-06-93 | 41 | 1100 | 24.4 | 7.48 | 8.23 | | | | | | | |

| | | | | | | | | | | | |
|----------|----|------|------|------|------|------|------|-----|------|---------|---------|
| 04-07-93 | 42 | 900 | 25.0 | 7.54 | 8.36 | 40.8 | 85.5 | 189 | 0.00 | 0.00 | 0.00000 |
| 04-08-93 | 43 | 900 | 25.3 | 7.51 | 8.20 | | | | | | |
| 04-09-93 | 44 | 900 | 25.3 | 7.53 | 8.04 | | | | | | |
| 04-10-93 | 45 | 1000 | 25.7 | 7.50 | 7.85 | | | | | | |
| 04-11-93 | 46 | 800 | 24.8 | 7.51 | 8.33 | | | | | | |
| 04-12-93 | 47 | 1000 | 25.2 | 7.47 | 8.07 | | | | | | |
| 04-13-93 | 48 | 900 | 25.0 | 7.51 | 7.85 | | | | | | |
| 04-14-93 | 49 | 600 | 25.1 | 7.51 | 7.52 | | | | | | |
| 04-15-93 | 50 | 900 | 25.4 | 7.49 | 7.49 | | | | | | |
| 04-16-93 | 51 | 1000 | 25.7 | 7.49 | 8.38 | | | | | | |
| 04-17-93 | 52 | 1000 | 25.3 | 7.50 | 7.59 | | | | | | |
| 04-18-93 | 53 | 1000 | 24.9 | 7.51 | 7.69 | | | | | | |
| 04-19-93 | 54 | 900 | 25.1 | 7.52 | 7.76 | | | | | | |
| 04-20-93 | 55 | 900 | 25.6 | 7.47 | 7.65 | | | | | | |
| 04-21-93 | 56 | 1000 | 25.7 | 7.49 | 7.44 | | | | | | |
| 04-22-93 | 57 | 900 | 25.0 | 7.53 | 7.09 | 40.8 | 85.5 | 187 | 0.00 | 0.00000 | |
| 04-23-93 | 58 | 900 | 24.9 | 7.51 | 7.78 | | | | | | |
| 04-24-93 | 59 | 1000 | 25.1 | 7.48 | 7.78 | | | | | | |
| 04-25-93 | 60 | 800 | 25.4 | 7.48 | 7.88 | | | | | | |
| 04-26-93 | 61 | 900 | 25.4 | 7.43 | 7.30 | | | | | | |
| 04-27-93 | 62 | 1000 | 24.4 | 7.54 | 7.96 | | | | | | |
| 04-28-93 | 63 | 900 | 25.1 | 7.35 | 7.43 | | | | | | |
| 04-29-93 | 64 | 1000 | 25.0 | 7.46 | 7.68 | 40.8 | 85.5 | 192 | 0.00 | 0.00043 | |
| 04-30-93 | 65 | 1000 | 25.3 | 7.42 | 7.59 | | | | | | |
| 05-01-93 | 66 | 1000 | 25.5 | 7.39 | 8.13 | | | | | | |
| 05-02-93 | 67 | 1000 | 25.6 | 7.40 | 7.87 | | | | | | |
| 05-03-93 | 68 | 1000 | 25.6 | 7.41 | 8.03 | | | | | | |
| 05-04-93 | 69 | 1000 | 25.8 | 7.44 | 7.47 | | | | | | |
| 05-05-93 | 70 | 900 | 25.7 | 7.36 | 7.38 | | | | | | |
| 05-06-93 | 71 | 1000 | 25.2 | 7.34 | 7.24 | 40.8 | 85.5 | 197 | 0.00 | 0.0032 | |
| 05-07-93 | 72 | 900 | 24.9 | 7.37 | 7.24 | | | | | | |
| 05-08-93 | 73 | 1640 | 25.2 | 7.20 | 5.28 | | | | | | |
| 05-09-93 | 74 | 1440 | 25.1 | 7.26 | 5.24 | | | | | | |
| 05-10-93 | 75 | 1000 | 24.8 | 7.35 | 6.94 | | | | | | |
| 05-11-93 | 76 | 900 | 25.2 | 7.44 | 7.58 | | | | | | |
| 05-12-93 | 77 | 1000 | 25.3 | 7.41 | 6.99 | | | | | | |
| 05-13-93 | 78 | 1000 | 25.3 | 7.44 | 7.08 | 40.8 | 85.5 | 194 | 0.00 | 0.00000 | |
| 05-14-93 | 79 | 800 | 25.0 | 7.51 | 7.15 | | | | | | |
| 05-15-93 | 80 | 800 | 25.1 | 7.50 | 7.49 | | | | | | |
| 05-16-93 | 81 | 1100 | 25.3 | 7.47 | 7.62 | | | | | | |
| 05-17-93 | 82 | 900 | 25.1 | 7.36 | 7.56 | | | | | | |
| 05-18-93 | 83 | 1000 | 25.1 | 7.37 | 7.66 | | | | | | |
| 05-19-93 | 84 | 1000 | 25.0 | 7.49 | 7.36 | | | | | | |
| 05-20-93 | 85 | 900 | 25.2 | 7.44 | 7.08 | 40.8 | 85.5 | 203 | 0.00 | 0.0046 | |
| 05-21-93 | 86 | 900 | 25.0 | 7.40 | 7.20 | | | | | | |
| 05-22-93 | 87 | 800 | 24.7 | 7.54 | 7.41 | | | | | | |
| 05-23-93 | 88 | 1000 | 24.6 | 7.45 | 7.32 | | | | | | |
| 05-24-93 | 89 | 800 | 24.8 | 7.45 | 7.25 | | | | | | |
| 05-25-93 | 90 | 1000 | 24.9 | 7.51 | 6.96 | | | | | | |
| 05-26-93 | 91 | 900 | 24.9 | 7.54 | 7.36 | | | | | | |
| 05-27-93 | 92 | 900 | 24.9 | 7.44 | 7.16 | | | | | | |
| 05-28-93 | 93 | 1000 | 25.0 | 7.49 | 7.32 | 40.8 | 85.5 | 190 | 0.00 | 0.0058 | |
| 05-29-93 | 94 | 1200 | 25.1 | 7.42 | 7.50 | | | | | | |
| 05-30-93 | 95 | 900 | 24.9 | 7.47 | 6.93 | | | | | | |

| | | | | | | | | | | | |
|----------|-----|------|------|------|------|------|------|-----|------|------|---------|
| 09-16-93 | 204 | 930 | 25.2 | 7.40 | 6.67 | 40.8 | 85.5 | 203 | 0.00 | 0.00 | 0.0000 |
| 09-17-93 | 205 | 930 | 24.6 | 7.30 | 6.86 | | | | | | |
| 09-18-93 | 206 | 800 | 24.7 | 7.44 | 6.99 | | | | | | |
| 09-19-93 | 207 | 1300 | 24.5 | 7.39 | 7.24 | | | | | | |
| 09-20-93 | 208 | 930 | 24.5 | 7.36 | 7.22 | | | | | | |
| 09-21-93 | 209 | 930 | 26.0 | 7.45 | 6.50 | | | | | | |
| 09-22-93 | 210 | 900 | 25.3 | 7.27 | 6.73 | | | | | | |
| 09-23-93 | 211 | 930 | 25.2 | 7.35 | 6.97 | 40.8 | 85.5 | 208 | 0.00 | 0.00 | 0.00000 |
| 09-24-93 | 212 | 1000 | 25.3 | 7.39 | 6.76 | | | | | | |
| 09-25-93 | 213 | 900 | 25.1 | 7.39 | 7.13 | | | | | | |
| 09-26-93 | 214 | 2100 | 25.7 | 7.52 | 7.12 | | | | | | |
| 09-27-93 | 215 | 1000 | 25.4 | 7.44 | 6.77 | | | | | | |
| 09-28-93 | 216 | 900 | 24.8 | 7.40 | 7.12 | | | | | | |
| 09-29-93 | 217 | 930 | 24.8 | 7.41 | 7.09 | 40.8 | 85.5 | 209 | 0.00 | 0.00 | 0.00011 |
| 09-30-93 | 218 | 1300 | 24.6 | 7.24 | 7.13 | | | | | | |
| 10-01-93 | 219 | 1100 | 24.2 | 7.31 | 8.41 | | | | | | |
| 10-02-93 | 220 | 830 | 24.7 | 7.35 | 7.16 | | | | | | |
| 10-03-93 | 221 | 830 | 24.9 | 7.33 | 6.88 | | | | | | |
| 10-04-93 | 222 | 1000 | 24.4 | 7.31 | 7.03 | | | | | | |
| 10-05-93 | 223 | 1000 | 24.9 | 7.32 | 6.69 | | | | | | |
| 10-06-93 | 224 | 930 | 24.5 | 7.42 | 7.26 | | | | | | |
| 10-07-93 | 225 | 930 | 24.9 | 7.46 | 7.09 | 47.6 | 85.5 | 206 | 0.00 | 0.00 | 0.00000 |
| 10-08-93 | 226 | 1000 | 25.0 | 7.36 | 7.06 | | | | | | |
| 10-09-93 | 227 | 1130 | 25.0 | 7.42 | 7.13 | | | | | | |
| 10-10-93 | 228 | 1130 | 24.6 | 7.42 | 7.33 | | | | | | |
| 10-11-93 | 229 | 1600 | 24.6 | 7.45 | 7.36 | | | | | | |
| 10-12-93 | 230 | 1000 | 24.5 | 7.33 | 7.25 | | | | | | |
| 10-13-93 | 231 | 930 | 24.7 | 7.39 | 7.11 | | | | | | |
| 10-14-93 | 232 | 1000 | 24.7 | 7.36 | 7.11 | 47.6 | 85.5 | 205 | 0.00 | 0.00 | 0.00000 |
| 10-15-93 | 233 | 1000 | 24.8 | 7.25 | 7.06 | | | | | | |
| 10-16-93 | 234 | 1230 | 24.7 | 7.27 | 7.08 | | | | | | |
| 10-17-93 | 235 | 1030 | 24.8 | 7.37 | 6.97 | | | | | | |
| 10-18-93 | 236 | 1000 | 24.8 | 7.30 | 6.92 | | | | | | |
| 10-19-93 | 237 | 900 | 24.8 | 7.26 | 6.97 | | | | | | |
| 10-20-93 | 238 | 1000 | 24.8 | 7.38 | 7.35 | | | | | | |
| 10-21-93 | 239 | 1000 | 24.8 | 7.39 | 7.15 | 47.6 | 85.5 | 206 | 0.00 | 0.00 | 0.00000 |
| 10-22-93 | 240 | 800 | 24.6 | 7.28 | 6.99 | | | | | | |
| 10-23-93 | 241 | 700 | 24.4 | 7.35 | 7.47 | | | | | | |
| 10-24-93 | 242 | 800 | 24.4 | 7.23 | 7.11 | | | | | | |
| 10-25-93 | 243 | 1000 | 24.4 | 7.28 | 6.74 | | | | | | |
| 10-26-93 | 244 | 1000 | 24.6 | 7.29 | 7.22 | | | | | | |
| 10-27-93 | 245 | 1000 | 24.4 | 7.34 | 7.24 | | | | | | |
| 10-28-93 | 246 | 1000 | 24.9 | 7.43 | 7.16 | 47.6 | 85.5 | 199 | 0.00 | 0.00 | 0.00012 |
| 10-29-93 | 247 | 1230 | 24.7 | 7.34 | 7.20 | | | | | | |
| 10-30-93 | 248 | 700 | 25.1 | 7.41 | 7.25 | | | | | | |
| 10-31-93 | 249 | 800 | 24.7 | 7.49 | 7.17 | | | | | | |
| 11-01-93 | 250 | 930 | 24.5 | 7.31 | 7.17 | | | | | | |
| 11-02-93 | 251 | 930 | 24.9 | 7.35 | 7.01 | | | | | | |
| 11-03-93 | 252 | 1000 | 25.1 | 7.32 | 6.80 | | | | | | |
| 11-04-93 | 253 | 1000 | 25.0 | 7.24 | 6.82 | 47.6 | 85.5 | 206 | 0.00 | 0.00 | 0.00000 |
| 11-05-93 | 254 | 1000 | 25.2 | 7.33 | 6.77 | | | | | | |
| 11-06-93 | 255 | 600 | 25.3 | 7.22 | 6.86 | | | | | | |
| 11-07-93 | 256 | 1000 | 25.0 | 7.28 | 6.74 | | | | | | |
| 11-08-93 | 257 | 1000 | 24.8 | 7.28 | 6.92 | | | | | | |

TANK No. 5
TANK CONCENTRATION: 1% GROUNDWATER (WELL CC-33B)
NO DEN EXPOSURE

| Date | Test Day | Time (Military) | Temperature (Celsius) | pH | Dissolved Oxygen (mg/L) | Alkalinity (mg/L as CaCO3) | Hardness (mg/L as CaCO3) | Conductivity (umohs/cm) | Total Residual Chlorine (mg/L) | Free Available Chlorine (mg/L) | Total Ammonia- Nitrogen (mg/L) | Total Unionized Ammonia- Nitrogen (mg/L) |
|----------|----------|--------------------|--------------------------|------|-------------------------------|----------------------------------|--------------------------------|----------------------------|---|---|---|--|
| 02-25-93 | 1 | 1500 | 24.7 | 6.74 | 7.72 | | | | | | | |
| 02-26-93 | 2 | 1000 | 24.1 | 7.15 | 7.67 | | | | | | | |
| 02-27-93 | 3 | 1000 | 24.4 | 7.12 | 7.46 | | | | | | | |
| 02-28-93 | 4 | 1100 | 24.7 | 7.08 | 7.38 | 34.0 | 85.5 | | | | | |
| 03-01-93 | 5 | 930 | 24.7 | 7.07 | 7.23 | | | | | | | |
| 03-02-93 | 6 | 1000 | 25.4 | 7.03 | 7.06 | | | | | | | |
| 03-03-93 | 7 | 1000 | 25.9 | 7.28 | 7.79 | | | | | | | |
| 03-04-93 | 8 | 1000 | 25.2 | 7.35 | 7.77 | | | | | | | |
| 03-05-93 | 9 | 800 | 25.6 | 7.35 | 7.52 | | | | | | | |
| 03-06-93 | 10 | 1000 | 24.9 | 7.20 | 7.78 | 34.0 | 85.5 | 320 | | | | |
| 03-07-93 | 11 | 1100 | 24.9 | 7.04 | 7.88 | | | | | | | |
| 03-08-93 | 12 | 900 | 25.0 | 7.00 | 7.74 | | | | | | | |
| 03-09-93 | 13 | 930 | 25.3 | 7.17 | 7.58 | | | | | | | |
| 03-10-93 | 14 | 900 | 24.9 | 7.02 | 7.69 | | | | | | | |
| 03-11-93 | 15 | 900 | 25.3 | 7.39 | 7.54 | 54.4 | 85.5 | 222 | 0.00 | 0.00 | | |
| 03-12-93 | 16 | 800 | 24.5 | 7.29 | 8.00 | | | | | | | |
| 03-13-93 | 17 | 1600 | 24.8 | 7.05 | 7.82 | | | | | | | |
| 03-14-93 | 18 | 1500 | 24.9 | 7.14 | 7.77 | | | | | | | |
| 03-15-93 | 19 | 900 | 24.7 | 7.05 | 7.93 | | | | | | | |
| 03-16-93 | 20 | 900 | 25.3 | 7.16 | 7.90 | | | | | | | |
| 03-17-93 | 21 | 900 | 25.5 | 7.17 | 7.43 | 34.0 | 85.5 | 433 | 0.00 | 0.00 | 0.019 | 0.00016 |
| 03-18-93 | 22 | 1030 | 25.3 | 7.02 | 7.72 | | | | | | | |
| 03-19-93 | 23 | 1000 | 25.3 | 7.08 | 7.60 | | | | | | | |
| 03-20-93 | 24 | 1100 | 26.8 | 7.13 | 7.69 | | | | | | | |
| 03-21-93 | 25 | 1230 | 25.5 | 7.12 | 8.09 | | | | | | | |
| 03-22-93 | 26 | 900 | 25.0 | 7.18 | 8.03 | | | | | | | |
| 03-23-93 | 27 | 900 | 24.9 | 7.12 | 7.96 | | | | | | | |
| 03-24-93 | 28 | 900 | 25.3 | 7.04 | 7.80 | | | | | | | |
| 03-25-93 | 29 | 900 | 25.1 | 6.96 | 7.78 | 34.0 | 68.4 | 216 | 0.00 | 0.00 | 0.000 | 0.00000 |
| 03-26-93 | 30 | 800 | 24.9 | 7.14 | 8.02 | | | | | | | |
| 03-27-93 | 31 | 1000 | 25.3 | 7.16 | 7.78 | | | | | | | |
| 03-28-93 | 32 | 1000 | 25.6 | 7.27 | 7.70 | | | | | | | |
| 03-29-93 | 33 | 1100 | 25.8 | 7.32 | 7.76 | | | | | | | |
| 03-30-93 | 34 | 1100 | 25.5 | 7.17 | 7.62 | | | | | | | |
| 03-31-93 | 35 | 1000 | 25.6 | 7.33 | 6.23 | | | | | | | |
| 04-01-93 | 36 | 1000 | 24.9 | 7.45 | 8.09 | 40.8 | 85.5 | 204 | 0.00 | 0.00 | 0.000 | 0.00000 |
| 04-02-93 | 37 | 1000 | 24.9 | 7.40 | 8.21 | | | | | | | |
| 04-03-93 | 38 | 1100 | 24.8 | 7.46 | 8.37 | | | | | | | |
| 04-04-93 | 39 | 1300 | 24.8 | 7.44 | 8.54 | | | | | | | |
| 04-05-93 | 40 | 900 | 24.8 | 7.38 | 8.45 | | | | | | | |
| 04-06-93 | 41 | 1100 | 24.5 | 7.40 | 8.26 | | | | | | | |

| | | | | | | | | | | | |
|----------|----|------|------|------|------|------|------|-----|------|------|---------|
| 04-07-93 | 42 | 900 | 25.1 | 7.48 | 8.29 | 40.8 | 85.5 | 207 | 0.00 | 0.00 | 0.00000 |
| 04-08-93 | 43 | 900 | 25.3 | 7.45 | 8.13 | | | | | | |
| 04-09-93 | 44 | 900 | 25.4 | 7.47 | 7.98 | | | | | | |
| 04-10-93 | 45 | 1000 | 25.7 | 7.40 | 7.73 | | | | | | |
| 04-11-93 | 46 | 800 | 24.9 | 7.42 | 8.24 | | | | | | |
| 04-12-93 | 47 | 1000 | 25.4 | 7.42 | 8.16 | | | | | | |
| 04-13-93 | 48 | 800 | 25.1 | 7.44 | 7.89 | | | | | | |
| 04-14-93 | 49 | 600 | 25.2 | 7.44 | 7.57 | | | | | | |
| 04-15-93 | 50 | 900 | 25.5 | 7.44 | 7.60 | | | | | | |
| 04-16-93 | 51 | 1000 | 25.7 | 7.48 | 8.65 | | | | | | |
| 04-17-93 | 52 | 1000 | 25.5 | 7.47 | 7.75 | | | | | | |
| 04-18-93 | 53 | 1000 | 24.9 | 7.54 | 7.87 | | | | | | |
| 04-19-93 | 54 | 900 | 25.2 | 7.51 | 7.98 | | | | | | |
| 04-20-93 | 55 | 900 | 25.6 | 7.43 | 7.82 | | | | | | |
| 04-21-93 | 56 | 1000 | 25.6 | 7.44 | 7.63 | | | | | | |
| 04-22-93 | 57 | 900 | 25.0 | 7.49 | | | | | | | |
| 04-23-93 | 58 | 900 | 24.9 | 7.46 | 7.38 | | | | | | |
| 04-24-93 | 59 | 1000 | 25.2 | 7.43 | 8.11 | | | | | | |
| 04-25-93 | 60 | 800 | 25.4 | 7.43 | 8.09 | | | | | | |
| 04-26-93 | 61 | 900 | 25.4 | 7.41 | 7.43 | | | | | | |
| 04-27-93 | 62 | 1000 | 24.4 | 7.55 | 8.13 | | | | | | |
| 04-28-93 | 63 | 900 | 25.1 | 7.34 | 7.74 | | | | | | |
| 04-29-93 | 64 | 1000 | 25.1 | 7.45 | 7.75 | | | | | | |
| 04-30-93 | 65 | 1000 | 25.3 | 7.45 | 8.16 | | | | | | |
| 05-01-93 | 66 | 1000 | 25.5 | 7.39 | 8.24 | | | | | | |
| 05-02-93 | 67 | 1000 | 25.6 | 7.36 | 7.97 | | | | | | |
| 05-03-93 | 68 | 1000 | 25.6 | 7.37 | 8.10 | | | | | | |
| 05-04-93 | 69 | 1000 | 25.8 | 7.39 | 7.51 | | | | | | |
| 05-05-93 | 70 | 900 | 25.7 | 7.39 | 7.39 | | | | | | |
| 05-06-93 | 71 | 1000 | 25.2 | 7.35 | 7.42 | | | | | | |
| 05-07-93 | 72 | 900 | 24.9 | 7.35 | 7.56 | | | | | | |
| 05-08-93 | 73 | 1640 | 25.2 | 7.25 | 5.54 | | | | | | |
| 05-09-93 | 74 | 1440 | 25.0 | 7.31 | 6.22 | | | | | | |
| 05-10-93 | 75 | 1000 | 25.0 | 7.32 | 7.24 | | | | | | |
| 05-11-93 | 76 | 900 | 25.2 | 7.39 | 7.81 | | | | | | |
| 05-12-93 | 77 | 1000 | 25.3 | 7.34 | 7.06 | | | | | | |
| 05-13-93 | 78 | 1000 | 25.2 | 7.37 | 7.22 | | | | | | |
| 05-14-93 | 79 | 800 | 24.8 | 7.43 | 7.31 | | | | | | |
| 05-15-93 | 80 | 800 | 25.0 | 7.40 | 7.65 | | | | | | |
| 05-16-93 | 81 | 1100 | 25.2 | 7.39 | 7.83 | | | | | | |
| 05-17-93 | 82 | 900 | 25.0 | 7.28 | 7.80 | | | | | | |
| 05-18-93 | 83 | 1000 | 25.0 | 7.29 | 7.88 | | | | | | |
| 05-19-93 | 84 | 1000 | 25.1 | 7.37 | 7.42 | | | | | | |
| 05-20-93 | 85 | 900 | 25.3 | 7.31 | 7.18 | | | | | | |
| 05-21-93 | 86 | 900 | 24.9 | 7.32 | 7.34 | | | | | | |
| 05-22-93 | 87 | 800 | 24.6 | 7.44 | 7.47 | | | | | | |
| 05-23-93 | 88 | 1000 | 24.5 | 7.37 | 7.48 | | | | | | |
| 05-24-93 | 89 | 800 | 24.7 | 7.33 | 7.35 | | | | | | |
| 05-25-93 | 90 | 1000 | 24.8 | 7.41 | 7.02 | | | | | | |
| 05-26-93 | 91 | 900 | 24.9 | 7.43 | 7.27 | | | | | | |
| 05-27-93 | 92 | 900 | 24.8 | 7.32 | 7.18 | | | | | | |
| 05-28-93 | 93 | 1000 | 24.9 | 7.35 | 7.19 | | | | | | |
| 05-29-93 | 94 | 1200 | 25.1 | 7.31 | 7.45 | | | | | | |
| 05-30-93 | 95 | 900 | 24.8 | 7.38 | 6.91 | | | | | | |

| | | | | | | | | | | | |
|----------|-----|------|------|------|------|------|------|-----|------|------|---------|
| 09-16-93 | 204 | 930 | 25.2 | 7.30 | 6.50 | 40.8 | 85.5 | 223 | 0.00 | 0.00 | 0.00000 |
| 09-17-93 | 205 | 930 | 24.5 | 7.22 | 6.71 | | | | | | |
| 09-18-93 | 206 | 800 | 24.7 | 7.34 | 6.90 | | | | | | |
| 09-19-93 | 207 | 1300 | 24.5 | 7.31 | 7.21 | | | | | | |
| 09-20-93 | 208 | 930 | 24.4 | 7.24 | 7.03 | | | | | | |
| 09-21-93 | 209 | 930 | 26.0 | 7.36 | 6.34 | | | | | | |
| 09-22-93 | 210 | 900 | 25.3 | 7.17 | 6.52 | | | | | | |
| 09-23-93 | 211 | 930 | 25.1 | 7.27 | 6.91 | 40.8 | 85.5 | 231 | 0.00 | 0.00 | 0.00000 |
| 09-24-93 | 212 | 1000 | 25.3 | 7.28 | 6.70 | | | | | | |
| 09-25-93 | 213 | 900 | 25.1 | 7.29 | 7.00 | | | | | | |
| 09-26-93 | 214 | 2100 | 25.7 | 7.40 | 7.12 | | | | | | |
| 09-27-93 | 215 | 1000 | 25.4 | 7.34 | 6.78 | | | | | | |
| 09-28-93 | 216 | 900 | 24.8 | 7.32 | 7.05 | | | | | | |
| 09-29-93 | 217 | 930 | 24.7 | 7.29 | 7.05 | 40.8 | 85.5 | 230 | 0.00 | 0.00 | 0.00008 |
| 09-30-93 | 218 | 1300 | 24.6 | 7.13 | 7.07 | | | | | | |
| 10-01-93 | 219 | 1100 | 24.0 | 7.23 | 8.33 | | | | | | |
| 10-02-93 | 220 | 830 | 24.7 | 7.24 | 6.96 | | | | | | |
| 10-03-93 | 221 | 830 | 24.9 | 7.29 | 6.80 | | | | | | |
| 10-04-93 | 222 | 1000 | 24.5 | 7.21 | 7.00 | | | | | | |
| 10-05-93 | 223 | 1000 | 24.9 | 7.22 | 6.75 | | | | | | |
| 10-06-93 | 224 | 930 | 24.4 | 7.33 | 7.19 | | | | | | |
| 10-07-93 | 225 | 930 | 24.9 | 7.37 | 7.03 | 47.6 | 85.5 | 226 | 0.00 | 0.00 | 0.00000 |
| 10-08-93 | 226 | 1000 | 24.9 | 7.29 | 7.06 | | | | | | |
| 10-09-93 | 227 | 1130 | 24.9 | 7.35 | 7.12 | | | | | | |
| 10-10-93 | 228 | 1130 | 24.8 | 7.34 | 7.31 | | | | | | |
| 10-11-93 | 229 | 1600 | 24.5 | 7.38 | 7.37 | | | | | | |
| 10-12-93 | 230 | 1000 | 24.4 | 7.27 | 7.25 | | | | | | |
| 10-13-93 | 231 | 930 | 24.7 | 7.32 | 7.18 | | | | | | |
| 10-14-93 | 232 | 1000 | 24.6 | 7.30 | 7.15 | 40.8 | 85.5 | 225 | 0.00 | 0.00 | 0.00000 |
| 10-15-93 | 233 | 1000 | 24.7 | 7.22 | 7.14 | | | | | | |
| 10-16-93 | 234 | 1230 | 24.6 | 7.22 | 7.14 | | | | | | |
| 10-17-93 | 235 | 1030 | 24.7 | 7.29 | 7.09 | | | | | | |
| 10-18-93 | 236 | 1000 | 24.7 | 7.25 | 7.02 | | | | | | |
| 10-19-93 | 237 | 900 | 24.7 | 7.21 | 7.07 | | | | | | |
| 10-20-93 | 238 | 1000 | 24.8 | 7.29 | 7.40 | | | | | | |
| 10-21-93 | 239 | 1000 | 24.8 | 7.28 | 7.15 | 40.8 | 85.5 | 232 | 0.00 | 0.00 | 0.00000 |
| 10-22-93 | 240 | 800 | 24.6 | 7.25 | 6.95 | | | | | | |
| 10-23-93 | 241 | 700 | 24.4 | 7.29 | 7.63 | | | | | | |
| 10-24-93 | 242 | 800 | 24.4 | 7.19 | 7.24 | | | | | | |
| 10-25-93 | 243 | 1000 | 24.4 | 7.19 | 6.68 | | | | | | |
| 10-26-93 | 244 | 1000 | 24.6 | 7.22 | 7.26 | | | | | | |
| 10-27-93 | 245 | 1000 | 24.4 | 7.27 | 7.26 | | | | | | |
| 10-28-93 | 246 | 1000 | 25.0 | 7.34 | 7.00 | 40.8 | 85.5 | 220 | 0.00 | 0.00 | 0.00023 |
| 10-29-93 | 247 | 1230 | 24.7 | 7.28 | 7.11 | | | | | | |
| 10-30-93 | 248 | 700 | 25.1 | 7.33 | 7.02 | | | | | | |
| 10-31-93 | 249 | 800 | 24.7 | 7.43 | 7.07 | | | | | | |
| 11-01-93 | 250 | 930 | 24.5 | 7.28 | 7.05 | | | | | | |
| 11-02-93 | 251 | 930 | 24.8 | 7.29 | 7.03 | | | | | | |
| 11-03-93 | 252 | 1000 | 24.9 | 7.22 | 6.91 | | | | | | |
| 11-04-93 | 253 | 1000 | 24.9 | 7.20 | 6.89 | 40.8 | 85.5 | 227 | 0.00 | 0.00 | 0.00000 |
| 11-05-93 | 254 | 1000 | 25.2 | 7.27 | 6.83 | | | | | | |
| 11-06-93 | 255 | 600 | 25.3 | 7.17 | 6.74 | | | | | | |
| 11-07-93 | 256 | 1000 | 24.9 | 7.22 | 6.79 | | | | | | |
| 11-08-93 | 257 | 1000 | 24.8 | 7.23 | 7.00 | | | | | | |

[illegible]

TANK No. 6
TANK CONCENTRATION: 1% GROUNDWATER (WELL CC-33B)
NO DEN EXPOSURE

| Date | Test Day | Time (Military) | Temperature (Celsius) | pH | Dissolved Oxygen (mg/L) | Alkalinity (mg/L as CaCO3) | Hardness (mg/L as CaCO3) | Conductivity (umohs/cm) | Total Residual Chlorine (mg/L) | Free Available Chlorine (mg/L) | Total Ammonia- Nitrogen (mg/L) | Total Un-ionized Ammonia- Nitrogen (mg/L) |
|----------|----------|--------------------|--------------------------|------|-------------------------------|----------------------------------|--------------------------------|----------------------------|---|---|---|---|
| 02-25-93 | 1 | 1500 | 24.8 | 6.76 | 7.67 | | | | | | | |
| 02-26-93 | 2 | 1000 | 24.1 | 7.15 | 7.69 | | | | | | | |
| 02-27-93 | 3 | 1000 | 24.4 | 7.12 | 7.36 | | | | | | | |
| 02-28-93 | 4 | 1100 | 24.7 | 7.08 | 7.36 | 34.0 | 85.5 | | | | | |
| 03-01-93 | 5 | 930 | 24.7 | 7.07 | 7.26 | | | | | | | |
| 03-02-93 | 6 | 1000 | 25.4 | 7.03 | 7.14 | | | | | | | |
| 03-03-93 | 7 | 1000 | 25.9 | 7.29 | 7.78 | | | | | | | |
| 03-04-93 | 8 | 1000 | 25.2 | 7.36 | 7.79 | | | | | | | |
| 03-05-93 | 9 | 800 | 25.8 | 7.37 | 7.54 | | | | | | | |
| 03-06-93 | 10 | 1000 | 24.9 | 7.19 | 7.75 | | | | | | | |
| 03-07-93 | 11 | 1100 | 24.9 | 7.03 | 7.90 | | | | | | | |
| 03-08-93 | 12 | 900 | 25.0 | 7.00 | 7.78 | | | | | | | |
| 03-09-93 | 13 | 930 | 25.3 | 7.17 | 7.62 | | | | | | | |
| 03-10-93 | 14 | 900 | 24.8 | 7.02 | 7.77 | | | | | | | |
| 03-11-93 | 15 | 900 | 25.3 | 7.40 | 7.60 | 54.4 | 85.5 | 229 | 0.00 | 0.00 | | |
| 03-12-93 | 16 | 800 | 24.5 | 7.29 | 8.08 | | | | | | | |
| 03-13-93 | 17 | 1600 | 24.8 | 7.04 | 7.83 | | | | | | | |
| 03-14-93 | 18 | 1500 | 24.9 | 7.14 | 7.80 | | | | | | | |
| 03-15-93 | 19 | 900 | 24.6 | 7.05 | 7.90 | | | | | | | |
| 03-16-93 | 20 | 900 | 25.3 | 7.16 | 7.90 | | | | | | | |
| 03-17-93 | 21 | 900 | 25.5 | 7.17 | 7.50 | 34.0 | 85.5 | 439 | 0.00 | 0.00 | 0.020 | 0.00017 |
| 03-18-93 | 22 | 1030 | 25.3 | 7.01 | 7.75 | | | | | | | |
| 03-19-93 | 23 | 1000 | 25.4 | 7.11 | 7.66 | | | | | | | |
| 03-20-93 | 24 | 1100 | 26.8 | 7.13 | 7.64 | | | | | | | |
| 03-21-93 | 25 | 1230 | 25.5 | 7.12 | 8.04 | | | | | | | |
| 03-22-93 | 26 | 900 | 25.0 | 7.18 | 7.99 | | | | | | | |
| 03-23-93 | 27 | 900 | 24.9 | 7.12 | 7.93 | | | | | | | |
| 03-24-93 | 28 | 900 | 25.3 | 7.04 | 7.80 | | | | | | | |
| 03-25-93 | 29 | 900 | 25.1 | 6.96 | 7.78 | 34.0 | 68.4 | 223 | 0.00 | 0.00 | 0.0000 | 0.00000 |
| 03-26-93 | 30 | 800 | 24.8 | 7.14 | 8.01 | | | | | | | |
| 03-27-93 | 31 | 1000 | 25.3 | 7.15 | 7.75 | | | | | | | |
| 03-28-93 | 32 | 1000 | 25.7 | 7.27 | 7.69 | | | | | | | |
| 03-29-93 | 33 | 1100 | 25.9 | 7.32 | 7.75 | | | | | | | |
| 03-30-93 | 34 | 1100 | 25.6 | 7.11 | 7.60 | | | | | | | |
| 03-31-93 | 35 | 1000 | 25.6 | 7.33 | 6.22 | | | | | | | |
| 04-01-93 | 36 | 1000 | 24.9 | 7.44 | 8.12 | 40.8 | 85.5 | 210 | 0.00 | 0.00 | 0.0000 | 0.00000 |
| 04-02-93 | 37 | 1000 | 24.9 | 7.39 | 8.22 | | | | | | | |
| 04-03-93 | 38 | 1100 | 24.9 | 7.45 | 8.44 | | | | | | | |
| 04-04-93 | 39 | 1300 | 24.8 | 7.42 | 8.58 | | | | | | | |
| 04-05-93 | 40 | 900 | 24.8 | 7.35 | 8.48 | | | | | | | |
| 04-06-93 | 41 | 1100 | 24.5 | 7.40 | 8.34 | | | | | | | |

| | | | | | | | | | | | |
|----------|----|------|------|------|------|------|------|-----|------|-------|---------|
| 04-07-93 | 42 | 900 | 25.0 | 7.47 | 8.40 | 40.8 | 85.5 | 211 | 0.00 | 0.00 | 0.0000 |
| 04-08-93 | 43 | 900 | 25.3 | 7.44 | 8.18 | | | | | | |
| 04-09-93 | 44 | 900 | 25.4 | 7.46 | 8.01 | | | | | | |
| 04-10-93 | 45 | 1000 | 25.7 | 7.39 | 7.77 | | | | | | |
| 04-11-93 | 46 | 800 | 24.8 | 7.41 | 8.20 | | | | | | |
| 04-12-93 | 47 | 1000 | 25.3 | 7.41 | 8.18 | | | | | | |
| 04-13-93 | 48 | 900 | 25.0 | 7.43 | 7.92 | | | | | | |
| 04-14-93 | 49 | 600 | 25.1 | 7.43 | 7.53 | | | | | | |
| 04-15-93 | 50 | 900 | 25.4 | 7.43 | 7.65 | | | | | | |
| 04-16-93 | 51 | 1000 | 25.7 | 7.47 | 8.69 | | | | | | |
| 04-17-93 | 52 | 1000 | 25.4 | 7.46 | 7.83 | 40.8 | 85.5 | 209 | 0.00 | 0.009 | 0.00014 |
| 04-18-93 | 53 | 1000 | 24.8 | 7.54 | 7.90 | | | | | | |
| 04-19-93 | 54 | 900 | 25.2 | 7.50 | 7.96 | | | | | | |
| 04-20-93 | 55 | 900 | 25.6 | 7.43 | 7.80 | | | | | | |
| 04-21-93 | 56 | 1000 | 25.7 | 7.42 | 7.62 | | | | | | |
| 04-22-93 | 57 | 900 | 25.0 | 7.48 | 7.62 | 40.8 | 85.5 | 206 | 0.00 | 0.000 | 0.00000 |
| 04-23-93 | 58 | 900 | 24.9 | 7.44 | 7.35 | | | | | | |
| 04-24-93 | 59 | 1000 | 25.2 | 7.42 | 8.09 | | | | | | |
| 04-25-93 | 60 | 800 | 25.4 | 7.42 | 8.09 | | | | | | |
| 04-26-93 | 61 | 900 | 25.4 | 7.41 | 7.43 | | | | | | |
| 04-27-93 | 62 | 1000 | 24.4 | 7.54 | 8.14 | | | | | | |
| 04-28-93 | 63 | 900 | 25.1 | 7.34 | 7.73 | | | | | | |
| 04-29-93 | 64 | 1000 | 25.0 | 7.44 | 7.76 | 40.8 | 85.5 | 217 | 0.00 | 0.024 | 0.00037 |
| 04-30-93 | 65 | 1000 | 25.3 | 7.42 | 7.93 | | | | | | |
| 05-01-93 | 66 | 1000 | 25.5 | 7.38 | 8.28 | | | | | | |
| 05-02-93 | 67 | 1000 | 25.6 | 7.35 | 7.94 | | | | | | |
| 05-03-93 | 68 | 1000 | 25.6 | 7.34 | 7.90 | | | | | | |
| 05-04-93 | 69 | 1000 | 25.8 | 7.37 | 7.38 | | | | | | |
| 05-05-93 | 70 | 900 | 25.7 | 7.35 | 7.23 | | | | | | |
| 05-06-93 | 71 | 1000 | 25.2 | 7.32 | 7.25 | | | | | | |
| 05-07-93 | 72 | 900 | 24.9 | 7.32 | 7.22 | 40.8 | 85.5 | 222 | 0.00 | 0.026 | 0.00031 |
| 05-08-93 | 73 | 1640 | 25.2 | 7.20 | 5.70 | | | | | | |
| 05-09-93 | 74 | 1440 | 25.1 | 7.29 | 5.35 | | | | | | |
| 05-10-93 | 75 | 1000 | 24.9 | 7.30 | 7.03 | | | | | | |
| 05-11-93 | 76 | 900 | 25.2 | 7.36 | 7.61 | | | | | | |
| 05-12-93 | 77 | 1000 | 25.3 | 7.34 | 7.00 | | | | | | |
| 05-13-93 | 78 | 1000 | 25.2 | 7.34 | 7.10 | 40.8 | 85.5 | 216 | 0.00 | 0.000 | 0.00000 |
| 05-14-93 | 79 | 800 | 24.9 | 7.40 | 7.22 | | | | | | |
| 05-15-93 | 80 | 800 | 25.0 | 7.39 | 7.55 | | | | | | |
| 05-16-93 | 81 | 1100 | 25.2 | 7.37 | 7.71 | | | | | | |
| 05-17-93 | 82 | 900 | 25.0 | 7.28 | 7.65 | | | | | | |
| 05-18-93 | 83 | 1000 | 25.0 | 7.27 | 7.78 | | | | | | |
| 05-19-93 | 84 | 1000 | 25.0 | 7.37 | 7.42 | | | | | | |
| 05-20-93 | 85 | 900 | 25.2 | 7.31 | 7.20 | 40.8 | 85.5 | 220 | 0.00 | 0.051 | 0.00059 |
| 05-21-93 | 86 | 900 | 24.9 | 7.32 | 7.35 | | | | | | |
| 05-22-93 | 87 | 800 | 24.6 | 7.43 | 7.50 | | | | | | |
| 05-23-93 | 88 | 1000 | 24.5 | 7.36 | 7.44 | | | | | | |
| 05-24-93 | 89 | 800 | 24.7 | 7.33 | 7.37 | | | | | | |
| 05-25-93 | 90 | 1000 | 24.8 | 7.41 | 7.05 | | | | | | |
| 05-26-93 | 91 | 900 | 24.9 | 7.41 | 7.29 | | | | | | |
| 05-27-93 | 92 | 900 | 24.8 | 7.31 | 7.11 | | | | | | |
| 05-28-93 | 93 | 1000 | 24.9 | 7.35 | 7.20 | 40.8 | 85.5 | 204 | 0.00 | 0.050 | 0.00057 |
| 05-29-93 | 94 | 1200 | 25.1 | 7.30 | 7.44 | | | | | | |
| 05-30-93 | 95 | 900 | 24.8 | 7.34 | 6.80 | | | | | | |

| | | | | | |
|----------|-----|------|------|------|------|
| 05-31-93 | 96 | 1100 | 24.9 | 7.35 | 6.98 |
| 06-01-93 | 97 | 1000 | 24.9 | 7.31 | 7.06 |
| 06-02-93 | 98 | 700 | 24.9 | 7.37 | 7.06 |
| 06-03-93 | 99 | 1000 | 25.0 | 7.12 | 6.72 |
| 06-04-93 | 100 | 900 | 25.0 | 7.20 | 6.75 |
| 06-05-93 | 101 | 1000 | 25.0 | 7.14 | 6.69 |
| 06-06-93 | 102 | 930 | 24.9 | 7.27 | 6.75 |
| 06-07-93 | 103 | 900 | 24.8 | 7.28 | 7.07 |
| 06-08-93 | 104 | 900 | 25.2 | 7.24 | 6.77 |
| 06-09-93 | 105 | 1000 | 25.2 | 7.28 | 6.86 |
| 06-10-93 | 106 | 900 | 25.3 | 7.23 | 7.05 |
| 06-11-93 | 107 | 800 | 25.4 | 7.24 | 6.89 |
| 06-12-93 | 108 | 1000 | 24.5 | 7.19 | 6.91 |
| 06-13-93 | 109 | 1030 | 24.8 | 7.17 | 6.80 |
| 06-14-93 | 110 | 1000 | 25.1 | 7.24 | 6.79 |
| 06-15-93 | 111 | 700 | 25.3 | 7.28 | 7.10 |
| 06-16-93 | 112 | 900 | 25.4 | 7.20 | 7.19 |
| 06-17-93 | 113 | 1000 | 25.1 | 7.24 | 7.15 |
| 06-18-93 | 114 | 800 | 25.4 | 7.26 | 6.82 |
| 06-19-93 | 115 | 800 | 25.5 | 7.21 | 6.68 |
| 06-20-93 | 116 | 900 | 25.4 | 7.22 | 7.00 |
| 06-21-93 | 117 | 900 | 24.8 | 7.24 | 6.76 |
| 06-22-93 | 118 | 900 | 24.7 | 7.23 | 6.76 |
| 06-23-93 | 119 | 900 | 24.7 | 7.25 | 6.76 |
| 06-24-93 | 120 | 900 | 24.5 | 7.26 | 6.84 |
| 06-25-93 | 121 | 900 | 24.8 | 7.12 | 6.60 |
| 06-26-93 | 122 | 1000 | 24.8 | 7.03 | 6.31 |
| 06-27-93 | 123 | 1100 | 24.8 | 7.29 | 6.78 |
| 06-28-93 | 124 | 800 | 24.9 | 7.31 | 6.66 |
| 06-29-93 | 125 | 900 | 25.0 | 7.21 | 6.47 |
| 06-30-93 | 126 | 1000 | 25.6 | 7.33 | 6.81 |
| 07-01-93 | 127 | 1000 | 25.7 | 7.33 | 6.79 |
| 07-02-93 | 128 | 900 | 25.6 | 7.22 | 6.49 |
| 07-03-93 | 129 | 1100 | 25.5 | 7.31 | 6.43 |
| 07-04-93 | 130 | 800 | 25.4 | 7.26 | 6.43 |
| 07-05-93 | 131 | 1000 | 24.6 | 7.19 | 6.48 |
| 07-06-93 | 132 | 1030 | 24.6 | 7.22 | 6.53 |
| 07-07-93 | 133 | 930 | 24.7 | 7.25 | 6.27 |
| 07-08-93 | 134 | 1030 | 24.8 | 7.23 | 6.34 |
| 07-09-93 | 135 | 1000 | 24.7 | 7.28 | 6.19 |
| 07-10-93 | 136 | 1100 | 24.7 | 7.25 | 6.31 |
| 07-11-93 | 137 | 1330 | 24.9 | 7.25 | 6.77 |
| 07-12-93 | 138 | 800 | 24.9 | 7.25 | 6.38 |
| 07-13-93 | 139 | 1000 | 25.1 | 7.31 | 6.49 |
| 07-14-93 | 140 | 800 | 25.1 | 7.36 | 6.51 |
| 07-15-93 | 141 | 900 | 25.0 | 7.25 | 6.47 |
| 07-16-93 | 142 | 800 | 25.0 | 7.29 | 6.28 |
| 07-17-93 | 143 | 900 | 25.0 | 7.18 | 6.54 |
| 07-18-93 | 144 | 1530 | 25.1 | 7.22 | 6.91 |
| 07-19-93 | 145 | 900 | 25.0 | 7.16 | 6.82 |
| 07-20-93 | 146 | 1000 | 25.0 | 7.24 | 6.92 |
| 07-21-93 | 147 | 1000 | 24.9 | 7.25 | 7.73 |
| 07-22-93 | 148 | 900 | 24.8 | 7.13 | 7.11 |
| 07-23-93 | 149 | 800 | 24.7 | 6.78 | 6.85 |

| | | | | | | | | | | | |
|----------|-----|------|------|------|------|------|------|-----|------|------|---------|
| 09-16-93 | 204 | 930 | 25.2 | 7.30 | 6.60 | 40.8 | 85.5 | 223 | 0.00 | 0.00 | 0.00000 |
| 09-17-93 | 205 | 930 | 24.5 | 7.22 | 6.82 | | | | | | |
| 09-18-93 | 206 | 800 | 24.7 | 7.34 | 7.04 | | | | | | |
| 09-19-93 | 207 | 1300 | 24.5 | 7.31 | 7.21 | | | | | | |
| 09-20-93 | 208 | 930 | 24.4 | 7.24 | 7.20 | | | | | | |
| 09-21-93 | 209 | 930 | 26.0 | 7.36 | 6.45 | | | | | | |
| 09-22-93 | 210 | 900 | 25.3 | 7.17 | 6.75 | | | | | | |
| 09-23-93 | 211 | 930 | 25.1 | 7.27 | 6.91 | 40.8 | 85.5 | 230 | 0.00 | 0.00 | 0.00000 |
| 09-24-93 | 212 | 1000 | 25.3 | 7.28 | 6.75 | | | | | | |
| 09-25-93 | 213 | 900 | 25.1 | 7.29 | 7.04 | | | | | | |
| 09-26-93 | 214 | 2100 | 25.7 | 7.40 | 7.21 | | | | | | |
| 09-27-93 | 215 | 1000 | 25.4 | 7.34 | 6.90 | | | | | | |
| 09-28-93 | 216 | 900 | 24.8 | 7.32 | 7.19 | | | | | | |
| 09-29-93 | 217 | 930 | 24.7 | 7.29 | 7.05 | 40.8 | 85.5 | 228 | 0.00 | 0.00 | 0.00010 |
| 09-30-93 | 218 | 1300 | 24.5 | 7.17 | 7.13 | | | | | | |
| 10-01-93 | 219 | 1100 | 24.1 | 7.22 | 8.32 | | | | | | |
| 10-02-93 | 220 | 830 | 24.7 | 7.27 | 6.84 | | | | | | |
| 10-03-93 | 221 | 830 | 24.9 | 7.27 | 6.82 | | | | | | |
| 10-04-93 | 222 | 1000 | 24.4 | 7.21 | 7.01 | | | | | | |
| 10-05-93 | 223 | 1000 | 24.8 | 7.22 | 6.84 | | | | | | |
| 10-06-93 | 224 | 930 | 24.4 | 7.33 | 7.21 | | | | | | |
| 10-07-93 | 225 | 930 | 24.8 | 7.37 | 7.07 | 47.6 | 85.5 | 226 | 0.00 | 0.00 | 0.00000 |
| 10-08-93 | 226 | 1000 | 24.9 | 7.29 | 7.10 | | | | | | |
| 10-09-93 | 227 | 1130 | 24.9 | 7.35 | 7.18 | | | | | | |
| 10-10-93 | 228 | 1130 | 24.6 | 7.35 | 7.43 | | | | | | |
| 10-11-93 | 229 | 1600 | 24.5 | 7.39 | 7.52 | | | | | | |
| 10-12-93 | 230 | 1000 | 24.4 | 7.28 | 7.34 | | | | | | |
| 10-13-93 | 231 | 930 | 24.7 | 7.33 | 7.26 | | | | | | |
| 10-14-93 | 232 | 1000 | 24.6 | 7.30 | 7.22 | 40.8 | 85.5 | 225 | 0.00 | 0.00 | 0.00000 |
| 10-15-93 | 233 | 1000 | 24.7 | 7.22 | 7.21 | | | | | | |
| 10-16-93 | 234 | 1230 | 24.6 | 7.23 | 7.19 | | | | | | |
| 10-17-93 | 235 | 1030 | 24.7 | 7.29 | 7.12 | | | | | | |
| 10-18-93 | 236 | 1000 | 24.7 | 7.25 | 7.08 | | | | | | |
| 10-19-93 | 237 | 900 | 24.7 | 7.21 | 7.22 | | | | | | |
| 10-20-93 | 238 | 1000 | 24.8 | 7.31 | 7.40 | | | | | | |
| 10-21-93 | 239 | 1000 | 24.8 | 7.31 | 7.23 | 40.8 | 85.5 | 221 | 0.00 | 0.00 | 0.00000 |
| 10-22-93 | 240 | 800 | 24.6 | 7.28 | 6.97 | | | | | | |
| 10-23-93 | 241 | 700 | 24.3 | 7.30 | 7.50 | | | | | | |
| 10-24-93 | 242 | 800 | 24.3 | 7.20 | 7.23 | | | | | | |
| 10-25-93 | 243 | 1000 | 24.4 | 7.19 | 6.69 | | | | | | |
| 10-26-93 | 244 | 1000 | 24.6 | 7.22 | 7.27 | | | | | | |
| 10-27-93 | 245 | 1000 | 24.4 | 7.27 | 7.30 | | | | | | |
| 10-28-93 | 246 | 1000 | 24.9 | 7.34 | 7.11 | 40.8 | 85.5 | 220 | 0.00 | 0.00 | 0.00022 |
| 10-29-93 | 247 | 1230 | 24.6 | 7.28 | 7.19 | | | | | | |
| 10-30-93 | 248 | 700 | 25.1 | 7.33 | 7.12 | | | | | | |
| 10-31-93 | 249 | 800 | 24.7 | 7.43 | 7.12 | | | | | | |
| 11-01-93 | 250 | 930 | 24.4 | 7.28 | 7.12 | | | | | | |
| 11-02-93 | 251 | 900 | 24.8 | 7.29 | 7.10 | | | | | | |
| 11-03-93 | 252 | 1000 | 24.9 | 7.21 | 6.92 | | | | | | |
| 11-04-93 | 253 | 1000 | 24.9 | 7.20 | 6.95 | 40.8 | 85.5 | 228 | 0.00 | 0.00 | 0.00000 |
| 11-05-93 | 254 | 1000 | 25.1 | 7.27 | 6.91 | | | | | | |
| 11-06-93 | 255 | 600 | 25.3 | 7.17 | 6.88 | | | | | | |
| 11-07-93 | 256 | 1000 | 24.9 | 7.22 | 6.79 | | | | | | |
| 11-08-93 | 257 | 1000 | 24.8 | 7.23 | 6.98 | | | | | | |

TANK No. 7
TANK CONCENTRATION: 1% GROUNDWATER (WELL CC-33B)
DEN EXPOSURE: 10 mg/L

| Date | Test Day | Time (Military) | Temperature (Celsius) | pH | Dissolved Oxygen (mg/L) | Alkalinity (mg/L as CaCO3) | Hardness (mg/L as CaCO3) | Conductivity (umohs/cm) | Total Residual Chlorine (mg/L) | Free Available Chlorine (mg/L) | Total Ammonia- Nitrogen (mg/L) | Total Unionized Ammonia- Nitrogen (mg/L) |
|----------|----------|--------------------|--------------------------|------|-------------------------------|----------------------------------|--------------------------------|----------------------------|---|---|---|--|
| 02-25-93 | 1 | 1500 | 24.7 | 6.76 | 7.67 | | | | | | | |
| 02-26-93 | 2 | 1000 | 24.1 | 7.15 | 7.66 | | | | | | | |
| 02-27-93 | 3 | 1000 | 24.4 | 7.14 | 7.40 | | | | | | | |
| 02-28-93 | 4 | 1100 | 24.7 | 7.09 | 7.38 | 34.0 | 85.5 | | | | | |
| 03-01-93 | 5 | 930 | 24.8 | 7.08 | 7.31 | | | | | | | |
| 03-02-93 | 6 | 1000 | 25.4 | 7.05 | 7.08 | | | | | | | |
| 03-03-93 | 7 | 1000 | 25.9 | 7.30 | 7.80 | | | | | | | |
| 03-04-93 | 8 | 1000 | 25.2 | 7.38 | 7.76 | | | | | | | |
| 03-05-93 | 9 | 800 | 25.6 | 7.36 | 7.52 | | | | | | | |
| 03-06-93 | 10 | 1000 | 24.9 | 7.20 | 7.79 | 34.0 | 85.5 | 316 | | | | |
| 03-07-93 | 11 | 1100 | 24.9 | 7.02 | 7.87 | | | | | | | |
| 03-08-93 | 12 | 900 | 25.0 | 7.00 | 7.78 | | | | | | | |
| 03-09-93 | 13 | 930 | 25.3 | 7.18 | 7.58 | | | | | | | |
| 03-10-93 | 14 | 900 | 24.9 | 7.03 | 7.79 | | | | | | | |
| 03-11-93 | 15 | 900 | 25.3 | 7.40 | 7.56 | | | | | | | |
| 03-12-93 | 16 | 800 | 24.5 | 7.29 | 8.07 | | | | | | | |
| 03-13-93 | 17 | 1600 | 24.8 | 7.05 | 7.77 | | | | | | | |
| 03-14-93 | 18 | 1500 | 24.9 | 7.14 | 7.77 | | | | | | | |
| 03-15-93 | 19 | 900 | 24.7 | 7.05 | 7.93 | | | | | | | |
| 03-16-93 | 20 | 900 | 25.3 | 7.16 | 7.88 | | | | | | | |
| 03-17-93 | 21 | 900 | 25.5 | 7.17 | 7.50 | | | | | | | |
| 03-18-93 | 22 | 1030 | 25.3 | 7.01 | 7.56 | | | | | | | |
| 03-19-93 | 23 | 1000 | 25.4 | 7.11 | 7.06 | | | | | | | |
| 03-20-93 | 24 | 1100 | 27.0 | 7.14 | 7.58 | | | | | | | |
| 03-21-93 | 25 | 1230 | 25.5 | 7.12 | 8.00 | | | | | | | |
| 03-22-93 | 26 | 900 | 25.0 | 7.18 | 7.95 | | | | | | | |
| 03-23-93 | 27 | 900 | 25.0 | 7.12 | 7.89 | | | | | | | |
| 03-24-93 | 28 | 900 | 25.3 | 7.04 | 7.79 | | | | | | | |
| 03-25-93 | 29 | 900 | 25.1 | 6.96 | 7.75 | | | | | | | |
| 03-26-93 | 30 | 800 | 24.8 | 7.14 | 8.00 | | | | | | | |
| 03-27-93 | 31 | 1000 | 25.3 | 7.15 | 7.72 | | | | | | | |
| 03-28-93 | 32 | 1000 | 25.7 | 7.27 | 7.62 | | | | | | | |
| 03-29-93 | 33 | 1100 | 25.9 | 7.32 | 7.60 | | | | | | | |
| 03-30-93 | 34 | 1100 | 25.6 | 7.12 | 7.68 | | | | | | | |
| 03-31-93 | 35 | 1000 | 25.6 | 7.29 | 6.07 | | | | | | | |
| 04-01-93 | 36 | 1000 | 25.0 | 7.41 | 7.95 | 40.8 | 68.4 | 206 | 0.00 | 0.00 | 0.0000 | 0.00000 |
| 04-02-93 | 37 | 1000 | 24.9 | 7.35 | 8.06 | | | | | | | |
| 04-03-93 | 38 | 1100 | 24.9 | 7.40 | 8.20 | | | | | | | |
| 04-04-93 | 39 | 1300 | 24.8 | 7.38 | 8.34 | | | | | | | |
| 04-05-93 | 40 | 900 | 24.8 | 7.33 | 8.40 | | | | | | | |
| 04-06-93 | 41 | 1100 | 24.5 | 7.37 | 8.27 | | | | | | | |

| | | | | | | | | | | | |
|----------|----|------|------|------|------|------|------|-----|------|------|---------|
| 04-07-93 | 42 | 900 | 25.1 | 7.44 | 8.31 | 40.8 | 85.5 | 206 | 0.00 | 0.00 | 0.00000 |
| 04-08-93 | 43 | 900 | 25.3 | 7.43 | 8.16 | | | | | | |
| 04-09-93 | 44 | 900 | 25.5 | 7.45 | 8.00 | | | | | | |
| 04-10-93 | 45 | 1000 | 25.8 | 7.38 | 7.69 | | | | | | |
| 04-11-93 | 46 | 800 | 24.9 | 7.41 | 8.20 | | | | | | |
| 04-12-93 | 47 | 1000 | 25.4 | 7.41 | 8.00 | | | | | | |
| 04-13-93 | 48 | 900 | 25.2 | 7.43 | 7.85 | | | | | | |
| 04-14-93 | 49 | 600 | 25.2 | 7.43 | 7.53 | | | | | | |
| 04-15-93 | 50 | 900 | 25.5 | 7.43 | 7.53 | | | | | | |
| 04-16-93 | 51 | 1000 | 25.7 | 7.45 | 8.49 | | | | | | |
| 04-17-93 | 52 | 1000 | 25.5 | 7.45 | 7.60 | | | | | | |
| 04-18-93 | 53 | 1000 | 25.0 | 7.54 | 7.77 | | | | | | |
| 04-19-93 | 54 | 900 | 25.3 | 7.50 | 7.83 | 40.8 | 85.5 | 205 | 0.00 | 0.00 | 0.00039 |
| 04-20-93 | 55 | 900 | 25.7 | 7.44 | 7.95 | | | | | | |
| 04-21-93 | 56 | 1000 | 25.7 | 7.44 | 7.59 | | | | | | |
| 04-22-93 | 57 | 900 | 25.1 | 7.49 | 7.37 | 40.8 | 85.5 | 202 | 0.00 | 0.00 | 0.00000 |
| 04-23-93 | 58 | 900 | 25.0 | 7.46 | 7.37 | | | | | | |
| 04-24-93 | 59 | 1000 | 25.3 | 7.42 | 7.97 | | | | | | |
| 04-25-93 | 60 | 800 | 25.5 | 7.43 | 7.98 | | | | | | |
| 04-26-93 | 61 | 900 | 25.5 | 7.41 | 7.43 | | | | | | |
| 04-27-93 | 62 | 1000 | 24.5 | 7.55 | 8.06 | | | | | | |
| 04-28-93 | 63 | 900 | 25.2 | 7.35 | 7.63 | | | | | | |
| 04-29-93 | 64 | 1000 | 25.2 | 7.44 | 7.72 | 40.8 | 85.5 | 210 | 0.00 | 0.00 | 0.00027 |
| 04-30-93 | 65 | 1000 | 25.4 | 7.43 | 7.94 | | | | | | |
| 05-01-93 | 66 | 1000 | 25.6 | 7.34 | 8.18 | | | | | | |
| 05-02-93 | 67 | 1000 | 25.7 | 7.33 | 7.90 | | | | | | |
| 05-03-93 | 68 | 1000 | 25.7 | 7.33 | 7.91 | | | | | | |
| 05-04-93 | 69 | 1000 | 25.9 | 7.37 | 7.37 | | | | | | |
| 05-05-93 | 70 | 900 | 25.8 | 7.31 | 7.09 | | | | | | |
| 05-06-93 | 71 | 1000 | 25.3 | 7.30 | 7.15 | 40.8 | 85.5 | 217 | 0.00 | 0.00 | 0.00033 |
| 05-07-93 | 72 | 900 | 25.0 | 7.29 | 7.27 | | | | | | |
| 05-08-93 | 73 | 1640 | 25.4 | 7.19 | 5.28 | | | | | | |
| 05-09-93 | 74 | 1440 | 25.2 | 7.29 | 5.14 | | | | | | |
| 05-10-93 | 75 | 1000 | 25.1 | 7.30 | 5.14 | | | | | | |
| 05-11-93 | 76 | 900 | 25.2 | 7.36 | 7.56 | | | | | | |
| 05-12-93 | 77 | 1000 | 25.3 | 7.34 | 6.96 | | | | | | |
| 05-13-93 | 78 | 1000 | 25.3 | 7.36 | 7.14 | 40.8 | 85.5 | 211 | 0.00 | 0.00 | 0.00000 |
| 05-14-93 | 79 | 800 | 24.9 | 7.41 | 7.10 | | | | | | |
| 05-15-93 | 80 | 800 | 25.1 | 7.40 | 7.51 | | | | | | |
| 05-16-93 | 81 | 1100 | 25.3 | 7.38 | 7.61 | | | | | | |
| 05-17-93 | 82 | 900 | 25.1 | 7.28 | 7.56 | | | | | | |
| 05-18-93 | 83 | 1000 | 25.1 | 7.27 | 7.58 | | | | | | |
| 05-19-93 | 84 | 1000 | 25.1 | 7.37 | 7.26 | 40.8 | 85.5 | 218 | 0.00 | 0.00 | 0.00070 |
| 05-20-93 | 85 | 900 | 25.3 | 7.32 | 6.98 | | | | | | |
| 05-21-93 | 86 | 900 | 25.0 | 7.32 | 7.14 | | | | | | |
| 05-22-93 | 87 | 800 | 24.7 | 7.44 | 7.25 | | | | | | |
| 05-23-93 | 88 | 1000 | 24.6 | 7.36 | 7.15 | | | | | | |
| 05-24-93 | 89 | 800 | 24.8 | 7.33 | 7.11 | | | | | | |
| 05-25-93 | 90 | 1000 | 24.9 | 7.42 | 6.74 | | | | | | |
| 05-26-93 | 91 | 900 | 24.9 | 7.44 | 7.18 | | | | | | |
| 05-27-93 | 92 | 900 | 24.8 | 7.33 | 7.04 | | | | | | |
| 05-28-93 | 93 | 1000 | 25.0 | 7.35 | 7.02 | 40.8 | 85.5 | 194 | 0.00 | 0.00 | 0.00074 |
| 05-29-93 | 94 | 1200 | 25.2 | 7.30 | 7.18 | | | | | | |
| 05-30-93 | 95 | 900 | 24.9 | 7.35 | 6.76 | | | | | | |

| | | | | | | | | | | | |
|----------|-----|------|------|------|------|------|------|-----|------|------|---------|
| 09-16-93 | 204 | 930 | 25.3 | 7.30 | 6.66 | 40.8 | 85.5 | 222 | 0.00 | 0.00 | 0.00000 |
| 09-17-93 | 205 | 930 | 24.6 | 7.22 | 6.81 | | | | | | |
| 09-18-93 | 206 | 800 | 24.7 | 7.34 | 7.00 | | | | | | |
| 09-19-93 | 207 | 1300 | 24.5 | 7.31 | 7.24 | | | | | | |
| 09-20-93 | 208 | 930 | 24.4 | 7.24 | 7.26 | | | | | | |
| 09-21-93 | 209 | 930 | 26.1 | 7.36 | 6.48 | | | | | | |
| 09-22-93 | 210 | 900 | 25.3 | 7.17 | 6.73 | | | | | | |
| 09-23-93 | 211 | 930 | 25.1 | 7.27 | 7.00 | 40.8 | 85.5 | 228 | 0.00 | 0.00 | 0.00000 |
| 09-24-93 | 212 | 1000 | 25.3 | 7.28 | 6.76 | | | | | | |
| 09-25-93 | 213 | 900 | 25.1 | 7.29 | 7.16 | | | | | | |
| 09-26-93 | 214 | 2100 | 25.7 | 7.40 | 7.15 | | | | | | |
| 09-27-93 | 215 | 1000 | 25.4 | 7.34 | 6.88 | | | | | | |
| 09-28-93 | 216 | 900 | 24.8 | 7.32 | 7.20 | | | | | | |
| 09-29-93 | 217 | 930 | 24.8 | 7.29 | 7.11 | 40.8 | 85.5 | 227 | 0.00 | 0.00 | 0.00009 |
| 09-30-93 | 218 | 1300 | 24.5 | 7.17 | 7.14 | | | | | | |
| 10-01-93 | 219 | 1100 | 24.0 | 7.24 | 8.21 | | | | | | |
| 10-02-93 | 220 | 830 | 24.7 | 7.27 | 6.96 | | | | | | |
| 10-03-93 | 221 | 830 | 25.0 | 7.26 | 6.78 | | | | | | |
| 10-04-93 | 222 | 1000 | 24.4 | 7.21 | 7.08 | | | | | | |
| 10-05-93 | 223 | 1000 | 24.8 | 7.22 | 6.89 | | | | | | |
| 10-06-93 | 224 | 930 | 24.4 | 7.33 | 7.33 | 47.6 | 85.5 | 225 | 0.00 | 0.00 | 0.00000 |
| 10-07-93 | 225 | 930 | 24.9 | 7.37 | 7.21 | | | | | | |
| 10-08-93 | 226 | 1000 | 24.9 | 7.29 | 7.11 | | | | | | |
| 10-09-93 | 227 | 1130 | 24.9 | 7.35 | 7.22 | | | | | | |
| 10-10-93 | 228 | 1130 | 24.6 | 7.34 | 7.42 | | | | | | |
| 10-11-93 | 229 | 1600 | 24.5 | 7.36 | 7.46 | | | | | | |
| 10-12-93 | 230 | 1000 | 24.4 | 7.27 | 7.30 | | | | | | |
| 10-13-93 | 231 | 930 | 24.7 | 7.32 | 7.25 | | | | | | |
| 10-14-93 | 232 | 1000 | 24.6 | 7.30 | 7.23 | 40.8 | 85.5 | 224 | 0.00 | 0.00 | 0.00000 |
| 10-15-93 | 233 | 1000 | 24.7 | 7.18 | 7.18 | | | | | | |
| 10-16-93 | 234 | 1230 | 24.6 | 7.22 | 7.22 | | | | | | |
| 10-17-93 | 235 | 1030 | 24.7 | 7.29 | 7.17 | | | | | | |
| 10-18-93 | 236 | 1000 | 24.7 | 7.25 | 7.08 | | | | | | |
| 10-19-93 | 237 | 900 | 24.7 | 7.21 | 7.24 | | | | | | |
| 10-20-93 | 238 | 1000 | 24.7 | 7.29 | 7.38 | | | | | | |
| 10-21-93 | 239 | 1000 | 24.8 | 7.31 | 7.16 | 40.8 | 85.5 | 224 | 0.00 | 0.00 | 0.00003 |
| 10-22-93 | 240 | 800 | 24.6 | 7.28 | 6.93 | | | | | | |
| 10-23-93 | 241 | 700 | 24.3 | 7.35 | 7.54 | | | | | | |
| 10-24-93 | 242 | 800 | 24.3 | 7.21 | 7.17 | | | | | | |
| 10-25-93 | 243 | 1000 | 24.3 | 7.19 | 6.78 | | | | | | |
| 10-26-93 | 244 | 1000 | 24.6 | 7.22 | 7.24 | | | | | | |
| 10-27-93 | 245 | 1000 | 24.3 | 7.27 | 7.19 | | | | | | |
| 10-28-93 | 246 | 1000 | 24.9 | 7.33 | 7.07 | 40.8 | 85.5 | 218 | 0.00 | 0.00 | 0.00024 |
| 10-29-93 | 247 | 1230 | 24.7 | 7.27 | 7.11 | | | | | | |
| 10-30-93 | 248 | 700 | 25.1 | 7.33 | 7.11 | | | | | | |
| 10-31-93 | 249 | 800 | 24.7 | 7.43 | 7.02 | | | | | | |
| 11-01-93 | 250 | 930 | 24.4 | 7.27 | 7.05 | | | | | | |
| 11-02-93 | 251 | 930 | 24.8 | 7.29 | 7.09 | | | | | | |
| 11-03-93 | 252 | 1000 | 24.9 | 7.22 | 6.94 | | | | | | |
| 11-04-93 | 253 | 1000 | 24.9 | 7.20 | 6.96 | 40.8 | 85.5 | 225 | 0.00 | 0.00 | 0.00000 |
| 11-05-93 | 254 | 1000 | 25.2 | 7.28 | 6.99 | | | | | | |
| 11-06-93 | 255 | 600 | 25.3 | 7.18 | 6.92 | | | | | | |
| 11-07-93 | 256 | 1000 | 24.9 | 7.22 | 6.81 | | | | | | |
| 11-08-93 | 257 | 1000 | 24.7 | 7.23 | 6.99 | | | | | | |

TANK No. 8
TANK CONCENTRATION: 1% GROUNDWATER (WELL CC-33B)
DEN EXPOSURE: 10 mg/L

| Date | Test Day | Time (Military) | Temperature (Celsius) | pH | Dissolved Oxygen (mg/L) | Alkalinity (mg/L as CaCO3) | Hardness (mg/L as CaCO3) | Conductivity (umohs/cm) | Total Residual Chlorine (mg/L) | Free Available Chlorine (mg/L) | Total Ammonia- Nitrogen (mg/L) | Total Unionized Ammonia- Nitrogen (mg/L) |
|----------|----------|--------------------|--------------------------|------|-------------------------------|----------------------------------|--------------------------------|----------------------------|---|---|---|--|
| 02-25-93 | 1 | 1500 | 24.7 | 6.78 | 7.48 | | | | | | | |
| 02-26-93 | 2 | 1000 | 24.2 | 7.15 | 7.59 | | | | | | | |
| 02-27-93 | 3 | 1000 | 24.5 | 7.13 | 7.41 | | | | | | | |
| 02-28-93 | 4 | 1100 | 24.8 | 7.09 | 7.36 | 34.0 | 85.5 | | | | | |
| 03-01-93 | 5 | 930 | 24.8 | 7.08 | 7.31 | | | | | | | |
| 03-02-93 | 6 | 1000 | 25.5 | 7.05 | 6.94 | | | | | | | |
| 03-03-93 | 7 | 1000 | 26.0 | 7.30 | 7.79 | | | | | | | |
| 03-04-93 | 8 | 1000 | 25.3 | 7.38 | 7.77 | | | | | | | |
| 03-05-93 | 9 | 800 | 25.6 | 7.37 | 7.47 | | | | | | | |
| 03-06-93 | 10 | 1000 | 25.0 | 7.21 | 7.79 | 34.0 | 85.5 | 314 | | | | |
| 03-07-93 | 11 | 1100 | 24.9 | 7.03 | 7.86 | | | | | | | |
| 03-08-93 | 12 | 900 | 25.0 | 7.01 | 7.77 | | | | | | | |
| 03-09-93 | 13 | 930 | 25.3 | 7.17 | 7.57 | | | | | | | |
| 03-10-93 | 14 | 900 | 24.8 | 7.03 | 7.75 | | | | | | | |
| 03-11-93 | 15 | 900 | 25.3 | 7.41 | 7.58 | | | | | | | |
| 03-12-93 | 16 | 800 | 24.5 | 7.31 | 8.04 | 54.4 | 85.5 | 215 | 0.00 | 0.00 | | |
| 03-13-93 | 17 | 1600 | 24.9 | 7.05 | 7.77 | | | | | | | |
| 03-14-93 | 18 | 1500 | 24.9 | 7.14 | 7.75 | | | | | | | |
| 03-15-93 | 19 | 900 | 24.8 | 7.06 | 7.92 | | | | | | | |
| 03-16-93 | 20 | 900 | 25.3 | 7.16 | 7.87 | | | | | | | |
| 03-17-93 | 21 | 900 | 25.6 | 7.17 | 7.46 | 34.0 | 85.5 | 425 | 0.00 | 0.00 | 0.019 | 0.00017 |
| 03-18-93 | 22 | 1030 | 25.4 | 7.02 | 7.61 | | | | | | | |
| 03-19-93 | 23 | 1000 | 25.3 | 7.11 | 7.54 | | | | | | | |
| 03-20-93 | 24 | 1100 | 27.0 | 7.13 | 7.61 | | | | | | | |
| 03-21-93 | 25 | 1230 | 25.5 | 7.12 | 7.98 | | | | | | | |
| 03-22-93 | 26 | 900 | 25.1 | 7.18 | 7.96 | | | | | | | |
| 03-23-93 | 27 | 900 | 24.9 | 7.12 | 7.92 | | | | | | | |
| 03-24-93 | 28 | 900 | 25.3 | 7.04 | 7.78 | | | | | | | |
| 03-25-93 | 29 | 900 | 25.2 | 6.97 | 7.74 | 34.0 | 68.4 | 213 | 0.00 | 0.00 | 0.000 | 0.00000 |
| 03-26-93 | 30 | 800 | 24.8 | 7.14 | 8.00 | | | | | | | |
| 03-27-93 | 31 | 1000 | 25.3 | 7.15 | 7.69 | | | | | | | |
| 03-28-93 | 32 | 1000 | 25.7 | 7.27 | 7.61 | | | | | | | |
| 03-29-93 | 33 | 1100 | 25.9 | 7.32 | 7.60 | | | | | | | |
| 03-30-93 | 34 | 1100 | 25.6 | 7.17 | 7.50 | | | | | | | |
| 03-31-93 | 35 | 1000 | 25.6 | 7.29 | 6.08 | | | | | | | |
| 04-01-93 | 36 | 1000 | 25.0 | 7.43 | 8.00 | 40.8 | 68.4 | 206 | 0.00 | 0.00 | 0.000 | 0.00000 |
| 04-02-93 | 37 | 1000 | 24.9 | 7.37 | 8.10 | | | | | | | |
| 04-03-93 | 38 | 1100 | 24.9 | 7.41 | 8.20 | | | | | | | |
| 04-04-93 | 39 | 1300 | 24.8 | 7.40 | 8.27 | | | | | | | |
| 04-05-93 | 40 | 900 | 24.8 | 7.34 | 8.38 | | | | | | | |
| 04-06-93 | 41 | 1100 | 24.4 | 7.38 | 8.27 | | | | | | | |

| | | | | | | | | | | | |
|----------|----|------|------|------|------|------|------|-----|------|------|---------|
| 04-07-93 | 42 | 900 | 25.0 | 7.45 | 8.31 | 40.8 | 85.5 | 208 | 0.00 | 0.00 | 0.00000 |
| 04-08-93 | 43 | 900 | 25.3 | 7.45 | 8.13 | | | | | | |
| 04-09-93 | 44 | 900 | 25.4 | 7.47 | 8.03 | | | | | | |
| 04-10-93 | 45 | 1000 | 25.8 | 7.39 | 7.70 | | | | | | |
| 04-11-93 | 46 | 800 | 24.9 | 7.44 | 8.30 | | | | | | |
| 04-12-93 | 47 | 1000 | 25.3 | 7.42 | 8.14 | | | | | | |
| 04-13-93 | 48 | 900 | 25.1 | 7.45 | 7.90 | | | | | | |
| 04-14-93 | 49 | 600 | 25.1 | 7.45 | 7.62 | | | | | | |
| 04-15-93 | 50 | 900 | 25.5 | 7.45 | 7.62 | 40.8 | 85.5 | 204 | 0.00 | 0.00 | 0.00015 |
| 04-16-93 | 51 | 1000 | 25.7 | 7.47 | 8.60 | | | | | | |
| 04-17-93 | 52 | 1000 | 25.4 | 7.46 | 7.67 | | | | | | |
| 04-18-93 | 53 | 1000 | 24.9 | 7.54 | 7.79 | | | | | | |
| 04-19-93 | 54 | 900 | 25.2 | 7.51 | 7.85 | | | | | | |
| 04-20-93 | 55 | 900 | 25.7 | 7.45 | 7.93 | | | | | | |
| 04-21-93 | 56 | 1000 | 25.8 | 7.46 | 7.60 | | | | | | |
| 04-22-93 | 57 | 900 | 25.0 | 7.50 | 7.30 | 40.8 | 85.5 | 201 | 0.00 | 0.00 | 0.00000 |
| 04-23-93 | 58 | 900 | 25.0 | 7.47 | 7.93 | | | | | | |
| 04-24-93 | 59 | 1000 | 25.3 | 7.43 | 7.36 | | | | | | |
| 04-25-93 | 60 | 800 | 25.5 | 7.43 | 8.01 | | | | | | |
| 04-26-93 | 61 | 900 | 25.5 | 7.41 | 7.36 | | | | | | |
| 04-27-93 | 62 | 1000 | 24.5 | 7.55 | 7.98 | | | | | | |
| 04-28-93 | 63 | 900 | 25.2 | 7.35 | 7.54 | | | | | | |
| 04-29-93 | 64 | 1000 | 25.1 | 7.44 | 7.58 | 40.8 | 85.5 | 210 | 0.00 | 0.00 | 0.00050 |
| 04-30-93 | 65 | 1000 | 25.4 | 7.43 | 7.90 | | | | | | |
| 05-01-93 | 66 | 1000 | 25.6 | 7.36 | 8.18 | | | | | | |
| 05-02-93 | 67 | 1000 | 25.7 | 7.36 | 7.95 | | | | | | |
| 05-03-93 | 68 | 1000 | 25.7 | 7.34 | 7.96 | | | | | | |
| 05-04-93 | 69 | 1000 | 25.9 | 7.38 | 7.31 | | | | | | |
| 05-05-93 | 70 | 900 | 25.8 | 7.33 | 7.08 | | | | | | |
| 05-06-93 | 71 | 1000 | 25.4 | 7.32 | 7.13 | 40.8 | 85.5 | 216 | 0.00 | 0.00 | 0.00038 |
| 05-07-93 | 72 | 900 | 25.0 | 7.28 | 6.57 | | | | | | |
| 05-08-93 | 73 | 1640 | 25.4 | 7.16 | 4.92 | | | | | | |
| 05-09-93 | 74 | 1440 | 25.3 | 7.26 | 5.36 | | | | | | |
| 05-10-93 | 75 | 1000 | 25.0 | 7.30 | 6.73 | | | | | | |
| 05-11-93 | 76 | 900 | 25.2 | 7.36 | 7.41 | | | | | | |
| 05-12-93 | 77 | 1000 | 25.3 | 7.34 | 7.03 | | | | | | |
| 05-13-93 | 78 | 1000 | 25.3 | 7.37 | 7.14 | 40.8 | 85.5 | 211 | 0.00 | 0.00 | 0.00000 |
| 05-14-93 | 79 | 800 | 24.9 | 7.42 | 7.14 | | | | | | |
| 05-15-93 | 80 | 800 | 25.1 | 7.41 | 7.48 | | | | | | |
| 05-16-93 | 81 | 1100 | 25.3 | 7.39 | 7.67 | | | | | | |
| 05-17-93 | 82 | 900 | 25.1 | 7.28 | 7.56 | | | | | | |
| 05-18-93 | 83 | 1000 | 25.1 | 7.29 | 7.69 | | | | | | |
| 05-19-93 | 84 | 1000 | 25.0 | 7.38 | 7.36 | | | | | | |
| 05-20-93 | 85 | 900 | 25.2 | 7.33 | 7.08 | 40.8 | 85.5 | 218 | 0.00 | 0.00 | 0.00062 |
| 05-21-93 | 86 | 900 | 25.0 | 7.32 | 7.17 | | | | | | |
| 05-22-93 | 87 | 800 | 24.6 | 7.44 | 7.34 | | | | | | |
| 05-23-93 | 88 | 1000 | 24.6 | 7.37 | 7.21 | | | | | | |
| 05-24-93 | 89 | 800 | 24.8 | 7.33 | 7.13 | | | | | | |
| 05-25-93 | 90 | 1000 | 24.9 | 7.43 | 6.74 | | | | | | |
| 05-26-93 | 91 | 900 | 24.9 | 7.44 | 7.37 | | | | | | |
| 05-27-93 | 92 | 900 | 24.8 | 7.35 | 7.16 | | | | | | |
| 05-28-93 | 93 | 1000 | 25.0 | 7.35 | 7.22 | 40.8 | 85.5 | 200 | 0.00 | 0.00 | 0.00068 |
| 05-29-93 | 94 | 1200 | 25.2 | 7.31 | 7.44 | | | | | | |
| 05-30-93 | 95 | 900 | 24.9 | 7.34 | 6.83 | | | | | | |

| | | | | | | | | | | | |
|----------|-----|------|------|------|------|------|------|-----|------|------|---------|
| 09-16-93 | 204 | 930 | 25.3 | 7.30 | 6.89 | 40.8 | 85.5 | 222 | 0.00 | 0.00 | 0.00000 |
| 09-17-93 | 205 | 930 | 24.5 | 7.22 | 6.92 | | | | | | |
| 09-18-93 | 206 | 800 | 24.7 | 7.34 | 7.04 | | | | | | |
| 09-19-93 | 207 | 1300 | 24.5 | 7.31 | 7.34 | | | | | | |
| 09-20-93 | 208 | 930 | 24.4 | 7.24 | 7.30 | | | | | | |
| 09-21-93 | 209 | 930 | 26.1 | 7.36 | 6.57 | | | | | | |
| 09-22-93 | 210 | 900 | 25.3 | 7.17 | 6.80 | | | | | | |
| 09-23-93 | 211 | 930 | 25.1 | 7.27 | 7.05 | 40.8 | 85.5 | 229 | 0.00 | 0.00 | 0.00000 |
| 09-24-93 | 212 | 1000 | 25.3 | 7.28 | 6.95 | | | | | | |
| 09-25-93 | 213 | 900 | 25.1 | 7.29 | 7.23 | | | | | | |
| 09-26-93 | 214 | 2100 | 25.7 | 7.40 | 7.18 | | | | | | |
| 09-27-93 | 215 | 1000 | 25.4 | 7.34 | 6.98 | | | | | | |
| 09-28-93 | 216 | 900 | 24.8 | 7.32 | 7.26 | | | | | | |
| 09-29-93 | 217 | 930 | 24.7 | 7.29 | 7.27 | | | | | | |
| 09-30-93 | 218 | 1300 | 24.5 | 7.16 | 7.21 | 40.8 | 85.5 | 228 | 0.00 | 0.00 | 0.00005 |
| 10-01-93 | 219 | 1100 | 24.1 | 7.25 | 8.50 | | | | | | |
| 10-02-93 | 220 | 830 | 24.7 | 7.29 | 6.92 | | | | | | |
| 10-03-93 | 221 | 830 | 25.0 | 7.28 | 6.74 | | | | | | |
| 10-04-93 | 222 | 1000 | 24.4 | 7.21 | 7.22 | | | | | | |
| 10-05-93 | 223 | 1000 | 24.8 | 7.22 | 7.13 | | | | | | |
| 10-06-93 | 224 | 930 | 24.3 | 7.33 | 7.50 | | | | | | |
| 10-07-93 | 225 | 930 | 24.8 | 7.37 | 7.24 | 47.6 | 85.5 | 226 | 0.00 | 0.00 | 0.00000 |
| 10-08-93 | 226 | 1000 | 24.9 | 7.29 | 7.19 | | | | | | |
| 10-09-93 | 227 | 1130 | 24.9 | 7.35 | 7.36 | | | | | | |
| 10-10-93 | 228 | 1130 | 24.6 | 7.34 | 7.43 | | | | | | |
| 10-11-93 | 229 | 1600 | 24.5 | 7.36 | 7.46 | | | | | | |
| 10-12-93 | 230 | 1000 | 24.4 | 7.28 | 7.45 | | | | | | |
| 10-13-93 | 231 | 930 | 24.7 | 7.32 | 7.36 | | | | | | |
| 10-14-93 | 232 | 1000 | 24.5 | 7.30 | 7.37 | 40.8 | 85.5 | 224 | 0.00 | 0.00 | 0.00000 |
| 10-15-93 | 233 | 1000 | 24.7 | 7.18 | 7.28 | | | | | | |
| 10-16-93 | 234 | 1230 | 24.6 | 7.22 | 7.28 | | | | | | |
| 10-17-93 | 235 | 1030 | 24.7 | 7.29 | 7.23 | | | | | | |
| 10-18-93 | 236 | 1000 | 24.7 | 7.25 | 7.18 | | | | | | |
| 10-19-93 | 237 | 900 | 24.7 | 7.21 | 7.29 | | | | | | |
| 10-20-93 | 238 | 1000 | 24.7 | 7.29 | 7.49 | | | | | | |
| 10-21-93 | 239 | 1000 | 24.8 | 7.30 | 7.27 | 40.8 | 85.5 | 225 | 0.00 | 0.00 | 0.00000 |
| 10-22-93 | 240 | 800 | 24.8 | 7.28 | 6.87 | | | | | | |
| 10-23-93 | 241 | 700 | 24.3 | 7.33 | 7.50 | | | | | | |
| 10-24-93 | 242 | 1000 | 24.4 | 7.20 | 7.21 | | | | | | |
| 10-25-93 | 243 | 1000 | 24.4 | 7.19 | 6.73 | | | | | | |
| 10-26-93 | 244 | 1000 | 24.6 | 7.21 | 7.15 | | | | | | |
| 10-27-93 | 245 | 1000 | 24.4 | 7.27 | 7.21 | | | | | | |
| 10-28-93 | 246 | 1000 | 24.9 | 7.33 | 7.10 | 40.8 | 85.5 | 218 | 0.00 | 0.00 | 0.00034 |
| 10-29-93 | 247 | 1230 | 24.7 | 7.27 | 7.06 | | | | | | |
| 10-30-93 | 248 | 700 | 25.0 | 7.33 | 7.21 | | | | | | |
| 10-31-93 | 249 | 800 | 24.7 | 7.42 | 7.14 | | | | | | |
| 11-01-93 | 250 | 930 | 24.4 | 7.27 | 7.08 | | | | | | |
| 11-02-93 | 251 | 930 | 24.8 | 7.29 | 7.05 | | | | | | |
| 11-03-93 | 252 | 1000 | 24.9 | 7.21 | 6.90 | | | | | | |
| 11-04-93 | 253 | 1000 | 24.9 | 7.20 | 6.97 | 40.8 | 85.5 | 225 | 0.00 | 0.00 | 0.00000 |
| 11-05-93 | 254 | 1000 | 25.2 | 7.28 | 6.92 | | | | | | |
| 11-06-93 | 255 | 1000 | 25.3 | 7.18 | 6.90 | | | | | | |
| 11-07-93 | 256 | 1000 | 24.9 | 7.22 | 6.76 | | | | | | |
| 11-08-93 | 257 | 1000 | 24.7 | 7.23 | 7.01 | | | | | | |

TANK No. 9
TANK CONCENTRATION: 10% GROUNDWATER (WELL CC-33B)
NO DEN EXPOSURE

| Date | Test Day | Time (Military) | Temperature (Celsius) | pH | Dissolved Oxygen (mg/L) | Alkalinity (mg/L as CaCO3) | Hardness (mg/L as CaCO3) | Conductivity (umohs/cm) | Total Residual Chlorine (mg/L) | Free Available Chlorine (mg/L) | Total Ammonia- Nitrogen (mg/L) | Total Unionized Ammonia- Nitrogen (mg/L) |
|----------|----------|--------------------|--------------------------|------|-------------------------------|----------------------------------|--------------------------------|----------------------------|---|---|---|--|
| 02-25-93 | 1 | 1500 | 24.6 | 6.53 | 7.57 | | | | | | | |
| 02-26-93 | 2 | 1000 | 24.0 | 6.70 | 7.62 | | | | | | | |
| 02-27-93 | 3 | 1000 | 24.4 | 6.88 | 7.44 | | | | | | | |
| 02-28-93 | 4 | 1100 | 24.8 | 6.95 | 7.36 | 34.0 | 102.6 | | | | | |
| 03-01-93 | 5 | 930 | 24.8 | 6.66 | 7.27 | | | | | | | |
| 03-02-93 | 6 | 1000 | 25.5 | 6.66 | 7.00 | | | | | | | |
| 03-03-93 | 7 | 1000 | 25.9 | 6.70 | 7.60 | | | | | | | |
| 03-04-93 | 8 | 1000 | 25.1 | 6.70 | 7.61 | | | | | | | |
| 03-05-93 | 9 | 800 | 25.4 | 6.70 | 7.36 | | | | | | | |
| 03-06-93 | 10 | 1000 | 24.9 | 6.52 | 7.63 | 34.0 | 102.6 | 459 | | | | |
| 03-07-93 | 11 | 1100 | 24.9 | 6.31 | 7.74 | | | | | | | |
| 03-08-93 | 12 | 900 | 25.0 | 6.40 | 7.64 | | | | | | | |
| 03-09-93 | 13 | 930 | 25.2 | 6.57 | 7.40 | | | | | | | |
| 03-10-93 | 14 | 900 | 24.8 | 6.49 | 7.72 | | | | | | | |
| 03-11-93 | 15 | 900 | 25.3 | 6.88 | 7.41 | 47.6 | 102.6 | 369 | 0.00 | 0.00 | | |
| 03-12-93 | 16 | 800 | 24.5 | 6.75 | 7.88 | | | | | | | |
| 03-13-93 | 17 | 1600 | 24.8 | 6.67 | 7.57 | | | | | | | |
| 03-14-93 | 18 | 1500 | 24.9 | 6.73 | 7.61 | | | | | | | |
| 03-15-93 | 19 | 900 | 24.7 | 6.74 | 7.72 | | | | | | | |
| 03-16-93 | 20 | 900 | 25.3 | 6.78 | 7.69 | | | | | | | |
| 03-17-93 | 21 | 900 | 25.5 | 6.73 | 7.29 | 34.0 | 102.6 | 568 | 0.00 | 0.00 | 0.030 | 0.00009 |
| 03-18-93 | 22 | 1030 | 25.4 | 6.78 | 7.45 | | | | | | | |
| 03-19-93 | 23 | 1000 | 25.8 | 7.10 | 7.47 | | | | | | | |
| 03-20-93 | 24 | 1100 | 26.9 | 6.73 | 7.47 | | | | | | | |
| 03-21-93 | 25 | 1230 | 25.6 | 6.72 | 7.83 | | | | | | | |
| 03-22-93 | 26 | 900 | 25.1 | 6.73 | 7.76 | | | | | | | |
| 03-23-93 | 27 | 900 | 25.0 | 6.71 | 7.73 | | | | | | | |
| 03-24-93 | 28 | 900 | 25.4 | 6.65 | 7.55 | 27.2 | 85.5 | 371 | 0.00 | 0.00 | 0.000 | 0.00000 |
| 03-25-93 | 29 | 900 | 25.1 | 6.65 | 7.63 | | | | | | | |
| 03-26-93 | 30 | 800 | 24.9 | 6.26 | 7.81 | | | | | | | |
| 03-27-93 | 31 | 1000 | 25.3 | 6.27 | 7.54 | | | | | | | |
| 03-28-93 | 32 | 1000 | 25.6 | 6.40 | 7.53 | | | | | | | |
| 03-29-93 | 33 | 1100 | 25.8 | 6.45 | 7.42 | | | | | | | |
| 03-30-93 | 34 | 1100 | 25.5 | 6.38 | 7.48 | | | | | | | |
| 03-31-93 | 35 | 1000 | 25.5 | 6.48 | 5.85 | 34.0 | 85.5 | 359 | 0.00 | 0.00 | 0.000 | 0.00000 |
| 04-01-93 | 36 | 1000 | 24.8 | 6.53 | 7.70 | | | | | | | |
| 04-02-93 | 37 | 1000 | 24.8 | 6.53 | 7.86 | | | | | | | |
| 04-03-93 | 38 | 1100 | 24.8 | 6.53 | 7.98 | | | | | | | |
| 04-04-93 | 39 | 1300 | 24.8 | 6.48 | 8.12 | | | | | | | |
| 04-05-93 | 40 | 900 | 24.8 | 6.47 | 8.23 | | | | | | | |
| 04-06-93 | 41 | 1100 | 24.7 | 6.49 | 8.01 | | | | | | | |

| | | | | | | | | | | | |
|----------|----|------|------|------|------|------|-------|-----|------|-------|---------|
| 04-07-93 | 42 | 900 | 25.0 | 6.58 | 8.11 | 34.0 | 85.5 | 361 | 0.00 | 0.00 | 0.00000 |
| 04-08-93 | 43 | 900 | 25.2 | 6.58 | 7.93 | | | | | | |
| 04-09-93 | 44 | 900 | 25.2 | 6.60 | 7.85 | | | | | | |
| 04-10-93 | 45 | 1000 | 25.4 | 6.54 | 7.58 | | | | | | |
| 04-11-93 | 46 | 800 | 24.8 | 6.58 | 8.07 | | | | | | |
| 04-12-93 | 47 | 1000 | 25.2 | 6.61 | 7.93 | | | | | | |
| 04-13-93 | 48 | 900 | 25.0 | 6.58 | 7.69 | | | | | | |
| 04-14-93 | 49 | 600 | 25.0 | 6.61 | 7.51 | | | | | | |
| 04-15-93 | 50 | 900 | 25.3 | 6.61 | 7.52 | | | | | | |
| 04-16-93 | 51 | 1000 | 25.5 | 8.60 | 8.43 | 34.0 | 102.6 | 350 | 0.00 | 0.023 | 0.00005 |
| 04-17-93 | 52 | 1000 | 25.3 | 6.59 | 7.55 | | | | | | |
| 04-18-93 | 53 | 1000 | 24.8 | 6.67 | 7.64 | | | | | | |
| 04-19-93 | 54 | 900 | 25.1 | 8.60 | 7.72 | | | | | | |
| 04-20-93 | 55 | 900 | 25.4 | 8.61 | 7.82 | | | | | | |
| 04-21-93 | 56 | 1000 | 25.4 | 6.64 | 7.54 | | | | | | |
| 04-22-93 | 57 | 900 | 24.8 | 6.64 | | 34.0 | 102.6 | 344 | 0.00 | 0.000 | 0.00000 |
| 04-23-93 | 58 | 900 | 24.8 | 6.66 | 7.00 | | | | | | |
| 04-24-93 | 59 | 1000 | 25.1 | 6.62 | 7.80 | | | | | | |
| 04-25-93 | 60 | 800 | 25.2 | 6.60 | 7.86 | | | | | | |
| 04-26-93 | 61 | 900 | 25.3 | 6.62 | 7.13 | | | | | | |
| 04-27-93 | 62 | 1000 | 24.4 | 6.71 | 7.83 | | | | | | |
| 04-28-93 | 63 | 900 | 25.0 | 6.64 | 7.46 | | | | | | |
| 04-29-93 | 64 | 1000 | 25.0 | 6.66 | 7.27 | 34.0 | 102.6 | 348 | 0.00 | 0.021 | 0.00005 |
| 04-30-93 | 65 | 1000 | 25.2 | 6.67 | 7.81 | | | | | | |
| 05-01-93 | 66 | 1000 | 25.4 | 6.65 | 8.10 | | | | | | |
| 05-02-93 | 67 | 1000 | 25.4 | 6.65 | 7.81 | | | | | | |
| 05-03-93 | 68 | 1000 | 25.4 | 6.65 | 7.90 | | | | | | |
| 05-04-93 | 69 | 1000 | 25.5 | 6.69 | 7.21 | | | | | | |
| 05-05-93 | 70 | 900 | 25.5 | 6.68 | 7.16 | | | | | | |
| 05-06-93 | 71 | 1000 | 25.1 | 6.66 | 7.13 | 34.0 | 85.5 | 358 | 0.00 | 0.022 | 0.00006 |
| 05-07-93 | 72 | 900 | 24.9 | 6.71 | 7.22 | | | | | | |
| 05-08-93 | 73 | 1640 | 25.2 | 6.63 | 4.96 | | | | | | |
| 05-09-93 | 74 | 1440 | 25.2 | 6.78 | 5.71 | | | | | | |
| 05-10-93 | 75 | 1000 | 24.9 | 6.63 | 6.92 | | | | | | |
| 05-11-93 | 76 | 900 | 25.0 | 6.65 | 7.42 | | | | | | |
| 05-12-93 | 77 | 1000 | 25.1 | 6.69 | 6.90 | | | | | | |
| 05-13-93 | 78 | 1000 | 25.1 | 6.70 | 6.91 | 34.0 | 102.6 | 350 | 0.00 | 0.000 | 0.00000 |
| 05-14-93 | 79 | 800 | 24.8 | 6.74 | 6.87 | | | | | | |
| 05-15-93 | 80 | 800 | 24.9 | 6.74 | 7.31 | | | | | | |
| 05-16-93 | 81 | 1100 | 25.1 | 6.72 | 7.52 | | | | | | |
| 05-17-93 | 82 | 900 | 24.9 | 6.64 | 7.40 | | | | | | |
| 05-18-93 | 83 | 1000 | 24.9 | 6.66 | 7.47 | | | | | | |
| 05-19-93 | 84 | 1000 | 25.2 | 6.76 | 7.15 | | | | | | |
| 05-20-93 | 85 | 900 | 25.3 | 6.73 | 6.82 | 34.0 | 102.6 | 357 | 0.00 | 0.062 | 0.00019 |
| 05-21-93 | 86 | 900 | 24.9 | 6.71 | 6.94 | | | | | | |
| 05-22-93 | 87 | 800 | 24.7 | 6.83 | 7.22 | | | | | | |
| 05-23-93 | 88 | 1000 | 24.6 | 6.77 | 7.06 | | | | | | |
| 05-24-93 | 89 | 800 | 24.7 | 6.70 | 7.01 | | | | | | |
| 05-25-93 | 90 | 1000 | 24.9 | 6.85 | 6.59 | | | | | | |
| 05-26-93 | 91 | 900 | 24.9 | 6.76 | 7.06 | | | | | | |
| 05-27-93 | 92 | 900 | 24.8 | 6.62 | 7.07 | | | | | | |
| 05-28-93 | 93 | 1000 | 24.9 | 6.65 | 6.91 | 34.0 | 102.6 | 361 | 0.00 | 0.062 | 0.00014 |
| 05-29-93 | 94 | 1200 | 25.1 | 6.64 | 7.18 | | | | | | |
| 05-30-93 | 95 | 900 | 24.9 | 6.75 | 6.63 | | | | | | |

| | | | | | | | | | | | |
|----------|-----|------|------|------|------|------|-------|-----|------|------|---------|
| 09-16-93 | 204 | 930 | 25.2 | 6.70 | 6.32 | 34.0 | 102.6 | 382 | 0.00 | 0.00 | 0.00000 |
| 09-17-93 | 205 | 930 | 24.6 | 6.66 | 6.48 | | | | | | |
| 09-18-93 | 206 | 800 | 24.8 | 6.77 | 6.78 | | | | | | |
| 09-19-93 | 207 | 1300 | 24.6 | 6.72 | 7.07 | | | | | | |
| 09-20-93 | 208 | 930 | 24.5 | 6.68 | 6.95 | | | | | | |
| 09-21-93 | 209 | 930 | 25.9 | 6.80 | 6.20 | | | | | | |
| 09-22-93 | 210 | 900 | 25.2 | 6.61 | 6.40 | | | | | | |
| 09-23-93 | 211 | 930 | 25.1 | 6.69 | 6.73 | 34.0 | 102.6 | 406 | 0.00 | 0.00 | 0.00000 |
| 09-24-93 | 212 | 1000 | 25.2 | 6.70 | 6.51 | | | | | | |
| 09-25-93 | 213 | 900 | 25.1 | 8.72 | 6.88 | | | | | | |
| 09-26-93 | 214 | 2100 | 25.6 | 8.78 | 6.87 | | | | | | |
| 09-27-93 | 215 | 1000 | 25.3 | 6.77 | 6.70 | | | | | | |
| 09-28-93 | 216 | 900 | 24.8 | 6.75 | 7.02 | 34.0 | 102.6 | 402 | 0.00 | 0.00 | 0.00001 |
| 09-29-93 | 217 | 930 | 24.8 | 6.72 | 6.89 | | | | | | |
| 09-30-93 | 218 | 1300 | 24.6 | 6.58 | 6.83 | | | | | | |
| 10-01-93 | 219 | 1100 | 24.1 | 6.67 | 8.12 | | | | | | |
| 10-02-93 | 220 | 830 | 24.8 | 6.73 | 6.87 | | | | | | |
| 10-03-93 | 221 | 830 | 25.0 | 8.78 | 6.70 | | | | | | |
| 10-04-93 | 222 | 1000 | 24.6 | 6.69 | 6.96 | | | | | | |
| 10-05-93 | 223 | 1000 | 24.9 | 8.65 | 6.72 | | | | | | |
| 10-06-93 | 224 | 930 | 24.5 | 8.77 | 7.08 | | | | | | |
| 10-07-93 | 225 | 930 | 24.9 | 8.82 | 6.92 | 40.8 | 102.6 | 406 | 0.00 | 0.00 | 0.00000 |
| 10-08-93 | 226 | 1000 | 25.0 | 6.77 | 6.92 | | | | | | |
| 10-09-93 | 227 | 1130 | 24.9 | 6.77 | 6.96 | | | | | | |
| 10-10-93 | 228 | 1130 | 24.6 | 6.76 | 7.17 | | | | | | |
| 10-11-93 | 229 | 1600 | 24.7 | 6.77 | 7.17 | | | | | | |
| 10-12-93 | 230 | 1000 | 24.5 | 6.73 | 7.15 | | | | | | |
| 10-13-93 | 231 | 930 | 24.8 | 6.74 | 6.97 | | | | | | |
| 10-14-93 | 232 | 1000 | 24.7 | 6.74 | 8.95 | 34.0 | 102.6 | 399 | 0.00 | 0.00 | 0.00000 |
| 10-15-93 | 233 | 1000 | 24.8 | 8.65 | 6.90 | | | | | | |
| 10-16-93 | 234 | 1230 | 24.7 | 6.68 | 6.81 | | | | | | |
| 10-17-93 | 235 | 1030 | 24.8 | 6.73 | 6.86 | | | | | | |
| 10-18-93 | 236 | 1000 | 24.7 | 6.69 | 6.86 | | | | | | |
| 10-19-93 | 237 | 900 | 24.7 | 6.65 | 8.93 | | | | | | |
| 10-20-93 | 238 | 1000 | 24.8 | 6.75 | 7.14 | | | | | | |
| 10-21-93 | 239 | 1000 | 24.8 | 6.78 | 6.90 | 34.0 | 102.6 | 397 | 0.00 | 0.00 | 0.00001 |
| 10-22-93 | 240 | 800 | 24.7 | 8.85 | 8.88 | | | | | | |
| 10-23-93 | 241 | 700 | 24.4 | 6.84 | 7.43 | | | | | | |
| 10-24-93 | 242 | 800 | 24.4 | 8.83 | 7.11 | | | | | | |
| 10-25-93 | 243 | 1000 | 24.5 | 6.64 | 6.57 | | | | | | |
| 10-26-93 | 244 | 1000 | 24.7 | 6.67 | 7.06 | | | | | | |
| 10-27-93 | 245 | 1000 | 24.4 | 6.76 | 7.07 | | | | | | |
| 10-28-93 | 246 | 1000 | 24.9 | 6.77 | 6.74 | 34.0 | 102.6 | 380 | 0.00 | 0.00 | 0.00011 |
| 10-29-93 | 247 | 1230 | 24.7 | 6.78 | 6.70 | | | | | | |
| 10-30-93 | 248 | 700 | 25.0 | 8.82 | 6.88 | | | | | | |
| 10-31-93 | 249 | 800 | 24.7 | 6.87 | 6.83 | | | | | | |
| 11-01-93 | 250 | 930 | 24.4 | 6.76 | 6.78 | | | | | | |
| 11-02-93 | 251 | 930 | 24.8 | 6.79 | 6.83 | | | | | | |
| 11-03-93 | 252 | 1000 | 25.0 | 6.74 | 6.66 | | | | | | |
| 11-04-93 | 253 | 1000 | 24.9 | 6.70 | 6.70 | 34.0 | 102.6 | 392 | 0.00 | 0.00 | 0.00000 |
| 11-05-93 | 254 | 1000 | 25.1 | 6.78 | 6.65 | | | | | | |
| 11-06-93 | 255 | 600 | 25.2 | 6.70 | 6.69 | | | | | | |
| 11-07-93 | 256 | 1000 | 25.0 | 6.74 | 6.59 | | | | | | |
| 11-08-93 | 257 | 1000 | 24.8 | 8.76 | 6.77 | | | | | | |

| | | | | | | | | | | | |
|----------|-----|------|------|-------|-------|------|-------|------|-------|--------|----------|
| 11-09-93 | 258 | 1000 | 24.7 | 6.74 | 6.76 | 34.0 | 102.6 | 380 | 0.00 | 0.005 | 0.00002 |
| 11-10-93 | 259 | 1000 | 24.7 | 6.70 | 6.71 | | | | | | |
| 11-11-93 | 260 | 1100 | 24.7 | 6.74 | 6.63 | | | | | | |
| 11-12-93 | 261 | 1000 | 24.9 | 6.72 | 6.47 | | | | | | |
| 11-13-93 | 262 | 1230 | 25.0 | 6.75 | 6.51 | | | | | | |
| 11-14-93 | 263 | 900 | 25.0 | 6.73 | 6.53 | | | | | | |
| 11-15-93 | 264 | 1000 | 25.2 | 6.71 | 6.35 | | | | | | |
| 11-16-93 | 265 | 1000 | 25.0 | 6.66 | 6.46 | | | | | | |
| 11-17-93 | 266 | 1200 | 25.2 | 6.66 | 5.79 | | | | | | |
| 11-18-93 | 267 | 1000 | 24.9 | 6.73 | 6.50 | 34.0 | 102.6 | 375 | 0.00 | 0.025 | 0.00007 |
| 11-19-93 | 268 | 1000 | 24.9 | 6.75 | 6.44 | | | | | | |
| 11-20-93 | 269 | 830 | 25.0 | 6.72 | 6.37 | | | | | | |
| 11-21-93 | 270 | 1300 | 24.7 | 6.71 | 6.57 | | | | | | |
| 11-22-93 | 271 | 930 | 24.6 | 6.71 | 6.60 | | | | | | |
| 11-23-93 | 272 | 1000 | 24.8 | 6.69 | 6.74 | | | | | | |
| 11-24-93 | 273 | 1300 | 24.7 | 6.81 | 6.14 | 34.0 | 102.6 | 386 | 0.00 | 0.048 | 0.00017 |
| 11-25-93 | 274 | 900 | 24.7 | 6.77 | 6.71 | | | | | | |
| 11-26-93 | 275 | 1200 | 24.6 | 6.70 | 6.76 | | | | | | |
| 11-27-93 | 276 | 1100 | 24.8 | 6.74 | 6.64 | | | | | | |
| 11-28-93 | 277 | 1100 | 25.3 | 6.72 | 6.29 | | | | | | |
| 11-29-93 | 278 | 600 | 24.1 | 7.40 | 7.71 | | | | | | |
| MEAN | | | 25.0 | 6.70 | 6.90 | 35.0 | 100.9 | 373 | 0.00 | 0.016 | 0.00004 |
| MINIMUM | | | 24.0 | 6.26 | 4.96 | 27.2 | 85.5 | 329 | 0.00 | 0.000 | 0.00000 |
| MAXIMUM | | | 26.9 | 7.40 | 8.43 | 47.6 | 102.6 | 568 | 0.00 | 0.092 | 0.00019 |
| Std Dev | | | 0.32 | 0.119 | 0.566 | 3.58 | 5.13 | 40.1 | 0.000 | 0.0215 | 0.000056 |
| N | | | 278 | 278 | 277 | 40 | 40 | 39 | 38 | 37 | 37 |

TANK No. 10
TANK CONCENTRATION: 10% GROUNDWATER (WELL CC-33B)
NO DEN EXPOSURE

| Date | Test Day | Time (Military) | Temperature (Celsius) | pH | Dissolved Oxygen (mg/L) | Alkalinity (mg/L as CaCO ₃) | Hardness (mg/L as CaCO ₃) | Conductivity (umohs/cm) | Total Residual Chlorine (mg/L) | Free Available Chlorine (mg/L) | Total Ammonia- Nitrogen (mg/L) | Total Un-ionized Ammonia- Nitrogen (mg/L) |
|----------|----------|--------------------|--------------------------|------|-------------------------------|---|---|----------------------------|---|---|---|---|
| 02-25-93 | 1 | 1500 | 24.6 | 6.49 | 7.57 | | | | | | | |
| 02-26-93 | 2 | 1000 | 24.0 | 8.70 | 7.57 | | | | | | | |
| 02-27-93 | 3 | 1000 | 24.4 | 6.88 | 7.37 | | | | | | | |
| 02-28-93 | 4 | 1100 | 24.9 | 6.95 | 7.23 | 34.0 | 102.6 | | | | | |
| 03-01-93 | 5 | 930 | 24.8 | 6.65 | 7.17 | | | | | | | |
| 03-02-93 | 6 | 1000 | 25.5 | 6.66 | 6.97 | | | | | | | |
| 03-03-93 | 7 | 1000 | 25.9 | 6.69 | 7.64 | | | | | | | |
| 03-04-93 | 8 | 1000 | 25.1 | 6.70 | 7.64 | | | | | | | |
| 03-05-93 | 9 | 800 | 25.4 | 6.70 | 7.40 | | | | | | | |
| 03-06-93 | 10 | 1000 | 24.9 | 6.52 | 7.67 | 27.2 | 102.6 | 459 | | | | |
| 03-07-93 | 11 | 1100 | 24.9 | 6.30 | 7.76 | | | | | | | |
| 03-08-93 | 12 | 900 | 25.0 | 6.39 | 7.66 | | | | | | | |
| 03-09-93 | 13 | 930 | 25.2 | 6.57 | 7.44 | | | | | | | |
| 03-10-93 | 14 | 900 | 24.8 | 6.49 | 7.74 | | | | | | | |
| 03-11-93 | 15 | 900 | 25.3 | 6.89 | 7.38 | 47.6 | 102.6 | 373 | 0.00 | 0.00 | | |
| 03-12-93 | 16 | 800 | 24.5 | 6.75 | 7.87 | | | | | | | |
| 03-13-93 | 17 | 1600 | 24.9 | 6.67 | 7.60 | | | | | | | |
| 03-14-93 | 18 | 1500 | 24.9 | 6.73 | 7.64 | | | | | | | |
| 03-15-93 | 19 | 900 | 24.7 | 6.74 | 7.75 | | | | | | | |
| 03-16-93 | 20 | 900 | 25.2 | 6.78 | 7.73 | | | | | | | |
| 03-17-93 | 21 | 900 | 25.5 | 6.73 | 7.34 | 34.0 | 102.6 | 573 | 0.00 | 0.00 | 0.018 | 0.00005 |
| 03-18-93 | 22 | 1030 | 25.4 | 6.77 | 7.50 | | | | | | | |
| 03-19-93 | 23 | 1000 | 25.7 | 7.10 | 7.46 | | | | | | | |
| 03-20-93 | 24 | 1100 | 26.9 | 6.73 | 7.45 | | | | | | | |
| 03-21-93 | 25 | 1230 | 25.6 | 6.72 | 7.84 | | | | | | | |
| 03-22-93 | 26 | 900 | 25.1 | 6.73 | 7.79 | | | | | | | |
| 03-23-93 | 27 | 900 | 25.0 | 6.71 | 7.75 | | | | | | | |
| 03-24-93 | 28 | 900 | 25.4 | 6.65 | 7.60 | | | | | | | |
| 03-25-93 | 29 | 900 | 25.2 | 6.65 | 7.64 | 27.2 | 85.5 | 371 | 0.00 | 0.00 | 0.000 | 0.00000 |
| 03-26-93 | 30 | 800 | 24.9 | 6.26 | 7.80 | | | | | | | |
| 03-27-93 | 31 | 1000 | 25.3 | 6.27 | 7.58 | | | | | | | |
| 03-28-93 | 32 | 1000 | 25.6 | 6.40 | 7.57 | | | | | | | |
| 03-29-93 | 33 | 1100 | 25.8 | 6.45 | 7.54 | | | | | | | |
| 03-30-93 | 34 | 1100 | 25.6 | 6.40 | 7.53 | | | | | | | |
| 03-31-93 | 35 | 1000 | 25.5 | 6.51 | 5.95 | 34.0 | 85.5 | 358 | 0.00 | 0.00 | 0.000 | 0.00000 |
| 04-01-93 | 36 | 1000 | 24.8 | 6.55 | 7.89 | | | | | | | |
| 04-02-93 | 37 | 1000 | 24.8 | 6.55 | 8.04 | | | | | | | |
| 04-03-93 | 38 | 1100 | 24.8 | 6.54 | 8.13 | | | | | | | |
| 04-04-93 | 39 | 1300 | 24.8 | 6.52 | 8.32 | | | | | | | |
| 04-05-93 | 40 | 900 | 24.8 | 6.49 | 8.28 | | | | | | | |
| 04-06-93 | 41 | 1100 | 24.7 | 6.49 | 8.09 | | | | | | | |

| | | | | | | | | | | | |
|----------|----|------|------|------|------|------|-------|-----|------|------|---------|
| 04-07-93 | 42 | 900 | 25.1 | 6.59 | 8.11 | 34.0 | 85.5 | 360 | 0.00 | 0.00 | 0.00000 |
| 04-08-93 | 43 | 900 | 25.2 | 6.60 | 7.97 | | | | | | |
| 04-09-93 | 44 | 900 | 25.3 | 6.61 | 7.80 | | | | | | |
| 04-10-93 | 45 | 1000 | 25.4 | 6.54 | 7.58 | | | | | | |
| 04-11-93 | 46 | 800 | 24.8 | 6.59 | 8.02 | | | | | | |
| 04-12-93 | 47 | 1000 | 25.2 | 6.61 | 7.93 | | | | | | |
| 04-13-93 | 48 | 900 | 25.0 | 6.59 | 7.68 | | | | | | |
| 04-14-93 | 49 | 600 | 25.0 | 6.61 | 7.37 | | | | | | |
| 04-15-93 | 50 | 900 | 25.3 | 6.61 | 7.47 | 34.0 | 85.5 | 351 | 0.00 | 0.00 | 0.00007 |
| 04-16-93 | 51 | 1000 | 25.5 | 6.62 | 8.42 | | | | | | |
| 04-17-93 | 52 | 1000 | 25.3 | 6.62 | 7.55 | | | | | | |
| 04-18-93 | 53 | 1000 | 24.8 | 6.69 | 7.67 | | | | | | |
| 04-19-93 | 54 | 900 | 25.1 | 6.62 | 7.72 | | | | | | |
| 04-20-93 | 55 | 900 | 25.4 | 6.62 | 7.82 | | | | | | |
| 04-21-93 | 56 | 1000 | 25.4 | 6.64 | 7.28 | | | | | | |
| 04-22-93 | 57 | 900 | 24.8 | 6.65 | | 34.0 | 102.6 | 343 | 0.00 | 0.00 | 0.00000 |
| 04-23-93 | 58 | 900 | 24.8 | 6.69 | 7.03 | | | | | | |
| 04-24-93 | 59 | 1000 | 25.1 | 6.64 | 7.71 | | | | | | |
| 04-25-93 | 60 | 800 | 25.2 | 6.61 | 7.76 | | | | | | |
| 04-26-93 | 61 | 900 | 25.3 | 6.62 | 7.05 | | | | | | |
| 04-27-93 | 62 | 1000 | 24.4 | 6.71 | 7.65 | | | | | | |
| 04-28-93 | 63 | 900 | 25.0 | 6.64 | 7.21 | | | | | | |
| 04-29-93 | 64 | 1000 | 25.0 | 6.66 | 7.44 | 34.0 | 102.6 | 349 | 0.00 | 0.00 | 0.00006 |
| 04-30-93 | 65 | 1000 | 25.2 | 6.67 | 7.45 | | | | | | |
| 05-01-93 | 66 | 1000 | 25.4 | 6.66 | 7.76 | | | | | | |
| 05-02-93 | 67 | 1000 | 25.4 | 6.66 | 7.55 | | | | | | |
| 05-03-93 | 68 | 1000 | 25.4 | 6.67 | 7.67 | | | | | | |
| 05-04-93 | 69 | 1000 | 25.5 | 6.69 | 7.22 | | | | | | |
| 05-05-93 | 70 | 900 | 25.5 | 6.68 | 6.85 | | | | | | |
| 05-06-93 | 71 | 1000 | 25.1 | 6.66 | 6.88 | 34.0 | 85.5 | 359 | 0.00 | 0.00 | 0.00010 |
| 05-07-93 | 72 | 900 | 24.9 | 6.71 | 6.71 | | | | | | |
| 05-08-93 | 73 | 1640 | 25.2 | 6.59 | 4.50 | | | | | | |
| 05-09-93 | 74 | 1440 | 25.1 | 6.72 | 5.22 | | | | | | |
| 05-10-93 | 75 | 1000 | 24.9 | 6.63 | 6.48 | | | | | | |
| 05-11-93 | 76 | 900 | 25.0 | 6.65 | 6.96 | | | | | | |
| 05-12-93 | 77 | 1000 | 25.1 | 6.69 | 6.53 | | | | | | |
| 05-13-93 | 78 | 1000 | 25.0 | 6.70 | 6.87 | 34.0 | 85.5 | 351 | 0.00 | 0.00 | 0.00000 |
| 05-14-93 | 79 | 800 | 24.8 | 6.74 | 6.77 | | | | | | |
| 05-15-93 | 80 | 800 | 24.9 | 6.74 | 7.20 | | | | | | |
| 05-16-93 | 81 | 1100 | 25.1 | 6.72 | 7.29 | | | | | | |
| 05-17-93 | 82 | 900 | 24.9 | 6.64 | 7.23 | | | | | | |
| 05-18-93 | 83 | 1000 | 24.9 | 6.66 | 7.26 | | | | | | |
| 05-19-93 | 84 | 1000 | 25.2 | 6.76 | 6.82 | | | | | | |
| 05-20-93 | 85 | 900 | 25.3 | 6.73 | 6.75 | 34.0 | 102.6 | 355 | 0.00 | 0.00 | 0.00021 |
| 05-21-93 | 86 | 900 | 24.9 | 6.71 | 6.73 | | | | | | |
| 05-22-93 | 87 | 800 | 24.7 | 6.83 | 6.92 | | | | | | |
| 05-23-93 | 88 | 1000 | 24.6 | 6.78 | 6.98 | | | | | | |
| 05-24-93 | 89 | 800 | 24.7 | 6.70 | 6.76 | | | | | | |
| 05-25-93 | 90 | 1000 | 24.9 | 6.85 | 6.33 | | | | | | |
| 05-26-93 | 91 | 900 | 24.9 | 6.78 | 6.96 | | | | | | |
| 05-27-93 | 92 | 900 | 24.8 | 6.62 | 6.95 | | | | | | |
| 05-28-93 | 93 | 1000 | 24.9 | 6.65 | 6.81 | 34.0 | 102.6 | 357 | 0.00 | 0.00 | 0.00020 |
| 05-29-93 | 94 | 1200 | 25.1 | 6.64 | 7.06 | | | | | | |
| 05-30-93 | 95 | 900 | 24.9 | 6.72 | 6.49 | | | | | | |

| | | | | | | | | | | | |
|----------|-----|------|------|------|------|------|-------|-----|------|------|---------|
| 09-16-93 | 204 | 930 | 25.1 | 6.70 | 6.46 | 34.0 | 102.6 | 373 | 0.00 | 0.00 | 0.00000 |
| 09-17-93 | 205 | 930 | 24.6 | 8.66 | 6.61 | | | | | | |
| 09-18-93 | 206 | 800 | 24.8 | 6.77 | 6.86 | | | | | | |
| 09-19-93 | 207 | 1300 | 24.6 | 7.05 | 7.05 | | | | | | |
| 09-20-93 | 208 | 930 | 24.5 | 6.68 | 6.93 | | | | | | |
| 09-21-93 | 209 | 930 | 25.9 | 6.80 | 6.19 | | | | | | |
| 09-22-93 | 210 | 900 | 25.2 | 6.61 | 6.49 | | | | | | |
| 09-23-93 | 211 | 930 | 25.1 | 6.69 | 6.83 | 34.0 | 102.6 | 403 | 0.00 | 0.00 | 0.00000 |
| 09-24-93 | 212 | 1000 | 25.2 | 8.70 | 6.58 | | | | | | |
| 09-25-93 | 213 | 900 | 25.0 | 8.72 | 6.91 | | | | | | |
| 09-26-93 | 214 | 2100 | 25.5 | 6.78 | 8.94 | | | | | | |
| 09-27-93 | 215 | 1000 | 25.3 | 6.77 | 6.64 | | | | | | |
| 09-28-93 | 216 | 900 | 24.8 | 6.75 | 6.91 | | | | | | |
| 09-29-93 | 217 | 930 | 24.8 | 6.72 | 6.96 | 34.0 | 102.6 | 400 | 0.00 | 0.00 | 0.00001 |
| 09-30-93 | 218 | 1300 | 24.6 | 6.55 | 6.82 | | | | | | |
| 10-01-93 | 219 | 1100 | 24.2 | 6.65 | 8.01 | | | | | | |
| 10-02-93 | 220 | 830 | 24.7 | 8.71 | 6.95 | | | | | | |
| 10-03-93 | 221 | 830 | 24.9 | 6.73 | 6.76 | | | | | | |
| 10-04-93 | 222 | 1000 | 24.5 | 8.69 | 6.94 | | | | | | |
| 10-05-93 | 223 | 1000 | 24.9 | 8.65 | 6.69 | | | | | | |
| 10-06-93 | 224 | 930 | 24.5 | 8.77 | 7.13 | | | | | | |
| 10-07-93 | 225 | 930 | 24.9 | 6.82 | 6.91 | 40.8 | 102.6 | 389 | 0.00 | 0.00 | 0.00000 |
| 10-08-93 | 226 | 1000 | 24.9 | 6.77 | 7.00 | | | | | | |
| 10-09-93 | 227 | 1130 | 24.9 | 8.77 | 7.04 | | | | | | |
| 10-10-93 | 228 | 1130 | 24.6 | 8.77 | 7.16 | | | | | | |
| 10-11-93 | 229 | 1600 | 24.6 | 6.79 | 7.19 | | | | | | |
| 10-12-93 | 230 | 1000 | 24.5 | 6.74 | 7.06 | | | | | | |
| 10-13-93 | 231 | 930 | 24.8 | 6.76 | 6.99 | | | | | | |
| 10-14-93 | 232 | 1000 | 24.8 | 8.74 | 6.98 | 34.0 | 102.6 | 384 | 0.00 | 0.00 | 0.00000 |
| 10-15-93 | 233 | 1000 | 24.8 | 6.67 | 6.91 | | | | | | |
| 10-16-93 | 234 | 1230 | 24.7 | 6.67 | 6.98 | | | | | | |
| 10-17-93 | 235 | 1030 | 24.8 | 8.73 | 6.96 | | | | | | |
| 10-18-93 | 236 | 1000 | 24.7 | 8.71 | 6.94 | | | | | | |
| 10-19-93 | 237 | 900 | 24.8 | 6.67 | 7.01 | | | | | | |
| 10-20-93 | 238 | 1000 | 24.8 | 8.75 | 7.17 | | | | | | |
| 10-21-93 | 239 | 1000 | 24.8 | 8.76 | 7.01 | 34.0 | 102.6 | 394 | 0.00 | 0.00 | 0.00000 |
| 10-22-93 | 240 | 800 | 24.6 | 8.81 | 6.79 | | | | | | |
| 10-23-93 | 241 | 700 | 24.3 | 8.82 | 7.32 | | | | | | |
| 10-24-93 | 242 | 800 | 24.4 | 6.79 | 7.01 | | | | | | |
| 10-25-93 | 243 | 1000 | 24.5 | 8.65 | 6.67 | | | | | | |
| 10-26-93 | 244 | 1000 | 24.7 | 8.67 | 7.21 | | | | | | |
| 10-27-93 | 245 | 1000 | 24.4 | 8.76 | 7.12 | | | | | | |
| 10-28-93 | 246 | 1000 | 24.9 | 6.77 | 6.90 | 34.0 | 102.6 | 378 | 0.00 | 0.00 | 0.00012 |
| 10-29-93 | 247 | 1230 | 24.7 | 8.78 | 6.74 | | | | | | |
| 10-30-93 | 248 | 700 | 25.0 | 8.82 | 6.91 | | | | | | |
| 10-31-93 | 249 | 800 | 24.7 | 8.87 | 6.89 | | | | | | |
| 11-01-93 | 250 | 930 | 24.4 | 6.76 | 6.90 | | | | | | |
| 11-02-93 | 251 | 930 | 24.8 | 6.79 | 6.81 | | | | | | |
| 11-03-93 | 252 | 1000 | 25.0 | 6.74 | 6.66 | | | | | | |
| 11-04-93 | 253 | 1000 | 24.9 | 8.70 | 6.75 | 34.0 | 102.6 | 389 | 0.00 | 0.00 | 0.00000 |
| 11-05-93 | 254 | 1000 | 25.2 | 6.78 | 6.72 | | | | | | |
| 11-06-93 | 255 | 600 | 25.2 | 6.70 | 6.71 | | | | | | |
| 11-07-93 | 256 | 1000 | 25.0 | 6.74 | 6.67 | | | | | | |
| 11-08-93 | 257 | 1000 | 24.8 | 8.76 | 6.81 | | | | | | |

| | | | | | | | | | | | |
|----------|-----|------|------|-------|-------|------|-------|------|-------|-------|---------|
| 11-09-93 | 258 | 1000 | 24.7 | 6.74 | 6.85 | 34.0 | 102.6 | 377 | 0.00 | 0.00 | 0.00000 |
| 11-10-93 | 259 | 1000 | 24.7 | 6.70 | 6.79 | | | | | | |
| 11-11-93 | 260 | 1100 | 24.7 | 6.74 | 6.72 | | | | | | |
| 11-12-93 | 261 | 1000 | 24.9 | 6.71 | 6.61 | | | | | | |
| 11-13-93 | 262 | 1230 | 24.9 | 6.75 | 6.57 | | | | | | |
| 11-14-93 | 263 | 900 | 25.0 | 6.73 | 6.57 | | | | | | |
| 11-15-93 | 264 | 1000 | 25.1 | 6.71 | 6.47 | | | | | | |
| 11-16-93 | 265 | 1000 | 25.0 | 6.66 | 6.60 | | | | | | |
| 11-17-93 | 266 | 1200 | 25.2 | 6.66 | 5.88 | | | | | | |
| 11-18-93 | 267 | 1000 | 24.9 | 6.73 | 6.61 | 34.0 | 102.6 | 371 | 0.00 | 0.00 | 0.00006 |
| 11-19-93 | 268 | 1000 | 24.9 | 6.75 | 6.56 | | | | | | |
| 11-20-93 | 269 | 830 | 24.9 | 6.72 | 6.59 | | | | | | |
| 11-21-93 | 270 | 1300 | 24.6 | 6.71 | 6.80 | | | | | | |
| 11-22-93 | 271 | 930 | 24.6 | 6.71 | 6.79 | | | | | | |
| 11-23-93 | 272 | 1000 | 24.7 | 6.69 | 6.82 | | | | | | |
| 11-24-93 | 273 | 1300 | 24.7 | 6.80 | 6.18 | 34.0 | 102.6 | 384 | 0.00 | 0.00 | 0.00017 |
| 11-25-93 | 274 | 900 | 24.7 | 6.77 | 6.76 | | | | | | |
| 11-26-93 | 275 | 1200 | 24.6 | 6.70 | 6.82 | | | | | | |
| 11-27-93 | 276 | 1100 | 24.8 | 6.74 | 6.67 | | | | | | |
| 11-28-93 | 277 | 1100 | 25.2 | 6.72 | 6.49 | | | | | | |
| 11-29-93 | 278 | 600 | 24.2 | 7.35 | 7.19 | | | | | | |
| MEAN | | | | | | | | | | | |
| MINIMUM | | | 25.0 | 6.70 | 6.87 | 34.9 | 100.0 | 374 | 0.00 | 0.00 | 0.00005 |
| MAXIMUM | | | 24.0 | 6.26 | 4.50 | 27.2 | 85.5 | 343 | 0.00 | 0.00 | 0.00000 |
| Std Dev | | | 26.9 | 7.35 | 8.42 | 47.6 | 102.6 | 573 | 0.00 | 0.00 | 0.00021 |
| N | | | 0.31 | 0.114 | 0.574 | 3.78 | 6.11 | 38.6 | 0.000 | 0.000 | 0.00064 |
| | | | 278 | 278 | 277 | 40 | 40 | 39 | 38 | 38 | 37 |

TANK No. 11
TANK CONCENTRATION: 10% GROUNDWATER (WELL CC-33B)
DEN EXPOSURE: 10 mg/L

| Date | Test Day | Time (Military) | Temperature (Celsius) | pH | Dissolved Oxygen (mg/L) | Alkalinity (mg/L as CaCO ₃) | Hardness (mg/L as CaCO ₃) | Conductivity (umohs/cm) | Total Residual Chlorine (mg/L) | Free Available Chlorine (mg/L) | Total Ammonia- Nitrogen (mg/L) | Total Unionized Ammonia- Nitrogen (mg/L) |
|----------|----------|--------------------|--------------------------|------|-------------------------------|---|---|----------------------------|---|---|---|--|
| 02-25-93 | 1 | 1500 | 24.5 | 6.48 | 7.50 | | | | | | | |
| 02-26-93 | 2 | 1000 | 23.9 | 6.64 | 7.57 | | | | | | | |
| 02-27-93 | 3 | 1000 | 24.4 | 6.82 | 7.33 | | | | | | | |
| 02-28-93 | 4 | 1100 | 24.8 | 6.88 | 7.31 | 34.0 | 102.6 | | | | | |
| 03-01-93 | 5 | 930 | 24.7 | 6.60 | 7.14 | | | | | | | |
| 03-02-93 | 6 | 1000 | 25.5 | 6.62 | 6.91 | | | | | | | |
| 03-03-93 | 7 | 1000 | 25.8 | 6.66 | 7.67 | | | | | | | |
| 03-04-93 | 8 | 1000 | 25.0 | 6.69 | 7.63 | | | | | | | |
| 03-05-93 | 9 | 800 | 25.4 | 6.69 | 7.40 | | | | | | | |
| 03-06-93 | 10 | 1000 | 24.8 | 6.52 | 7.66 | 34.0 | 102.6 | 461 | | | | |
| 03-07-93 | 11 | 1100 | 24.8 | 6.30 | 7.71 | | | | | | | |
| 03-08-93 | 12 | 900 | 24.9 | 6.40 | 7.65 | | | | | | | |
| 03-09-93 | 13 | 930 | 25.2 | 6.57 | 7.40 | | | | | | | |
| 03-10-93 | 14 | 900 | 24.8 | 6.49 | 7.56 | | | | | | | |
| 03-11-93 | 15 | 900 | 25.3 | 6.90 | 7.40 | 47.6 | 102.6 | 378 | 0.00 | 0.00 | | |
| 03-12-93 | 16 | 800 | 24.5 | 6.75 | 7.86 | | | | | | | |
| 03-13-93 | 17 | 1600 | 24.8 | 6.67 | 7.59 | | | | | | | |
| 03-14-93 | 18 | 1500 | 24.9 | 6.73 | 7.62 | | | | | | | |
| 03-15-93 | 19 | 900 | 24.7 | 6.74 | 7.65 | | | | | | | |
| 03-16-93 | 20 | 900 | 25.2 | 6.79 | 7.71 | | | | | | 0.029 | 0.00009 |
| 03-17-93 | 21 | 900 | 25.5 | 6.73 | 7.29 | 34.0 | 102.6 | 565 | 0.00 | 0.00 | | |
| 03-18-93 | 22 | 1030 | 25.4 | 6.78 | 7.43 | | | | | | | |
| 03-19-93 | 23 | 1000 | 25.7 | 7.11 | 7.49 | | | | | | | |
| 03-20-93 | 24 | 1100 | 26.9 | 6.73 | 7.43 | | | | | | | |
| 03-21-93 | 25 | 1230 | 25.6 | 6.72 | 7.82 | | | | | | | |
| 03-22-93 | 26 | 900 | 25.1 | 6.73 | 7.73 | | | | | | | |
| 03-23-93 | 27 | 900 | 25.0 | 6.71 | 7.75 | | | | | | | |
| 03-24-93 | 28 | 900 | 25.3 | 6.65 | 7.54 | | | | | | | |
| 03-25-93 | 29 | 900 | 25.1 | 6.65 | 7.63 | 27.2 | 85.5 | 373 | 0.00 | 0.00 | 0.000 | 0.00000 |
| 03-26-93 | 30 | 800 | 24.9 | 6.25 | 7.78 | | | | | | | |
| 03-27-93 | 31 | 1000 | 25.3 | 6.27 | 7.55 | | | | | | | |
| 03-28-93 | 32 | 1000 | 25.6 | 6.38 | 7.50 | | | | | | | |
| 03-29-93 | 33 | 1100 | 25.7 | 6.45 | 7.38 | | | | | | | |
| 03-30-93 | 34 | 1100 | 25.5 | 6.40 | 7.58 | | | | | | | |
| 03-31-93 | 35 | 1000 | 25.5 | 6.48 | 5.78 | | | | | | | |
| 04-01-93 | 36 | 1000 | 24.8 | 6.53 | 7.68 | 34.0 | 85.5 | 358 | 0.00 | 0.00 | 0.000 | 0.00000 |
| 04-02-93 | 37 | 1000 | 24.7 | 6.51 | 7.84 | | | | | | | |
| 04-03-93 | 38 | 1100 | 24.8 | 6.53 | 7.90 | | | | | | | |
| 04-04-93 | 39 | 1300 | 24.7 | 6.46 | 8.08 | | | | | | | |
| 04-05-93 | 40 | 900 | 24.7 | 6.44 | 8.16 | | | | | | | |
| 04-06-93 | 41 | 1100 | 24.6 | 6.47 | 8.00 | | | | | | | |

| | | | | | | | | | | |
|----------|------|------|------|------|------|-------|-----|------|-------|---------|
| 04-07-93 | 900 | 25.0 | 6.58 | 8.12 | 34.0 | 85.5 | 361 | 0.00 | 0.00 | 0.00000 |
| 04-08-93 | 900 | 25.2 | 6.60 | 7.95 | | | | | | |
| 04-09-93 | 900 | 25.2 | 6.58 | 7.80 | | | | | | |
| 04-10-93 | 1000 | 25.3 | 6.53 | 7.54 | | | | | | |
| 04-11-93 | 800 | 24.8 | 6.58 | 8.00 | | | | | | |
| 04-12-93 | 1000 | 25.2 | 6.60 | 7.79 | | | | | | |
| 04-13-93 | 900 | 25.0 | 6.57 | 7.61 | | | | | | |
| 04-14-93 | 600 | 25.0 | 6.61 | 7.36 | | | | | | |
| 04-15-93 | 900 | 25.3 | 6.61 | 7.43 | | | | | | |
| 04-16-93 | 1000 | 25.5 | 6.60 | 8.22 | | | | | | |
| 04-17-93 | 1000 | 25.3 | 6.59 | 7.40 | | | | | | |
| 04-18-93 | 1000 | 24.8 | 6.66 | 7.52 | | | | | | |
| 04-19-93 | 900 | 25.1 | 6.58 | 7.63 | | | | | | |
| 04-20-93 | 900 | 25.4 | 6.58 | 7.68 | | | | | | |
| 04-21-93 | 1000 | 25.4 | 6.61 | 7.40 | | | | | | |
| 04-22-93 | 900 | 24.9 | 6.61 | | 34.0 | 102.6 | 345 | 0.00 | 0.00 | 0.00000 |
| 04-23-93 | 900 | 24.8 | 6.63 | 7.16 | | | | | | |
| 04-24-93 | 1000 | 25.1 | 6.62 | 7.64 | | | | | | |
| 04-25-93 | 800 | 25.2 | 6.60 | 7.79 | | | | | | |
| 04-26-93 | 900 | 25.3 | 6.60 | 7.07 | | | | | | |
| 04-27-93 | 1000 | 24.4 | 6.71 | 7.74 | | | | | | |
| 04-28-93 | 900 | 25.0 | 6.64 | 7.34 | | | | | | |
| 04-29-93 | 1000 | 25.0 | 6.65 | 7.34 | | | | | | |
| 04-30-93 | 1000 | 25.2 | 6.66 | 7.59 | 34.0 | 102.6 | 350 | 0.00 | 0.023 | 0.00006 |
| 05-01-93 | 1000 | 25.4 | 6.64 | 8.04 | | | | | | |
| 05-02-93 | 1000 | 25.4 | 6.63 | 7.80 | | | | | | |
| 05-03-93 | 1000 | 25.4 | 6.63 | 7.89 | | | | | | |
| 05-04-93 | 1000 | 25.5 | 6.65 | 7.27 | | | | | | |
| 05-05-93 | 900 | 25.5 | 6.66 | 7.01 | | | | | | |
| 05-06-93 | 1000 | 25.1 | 6.66 | 7.07 | 34.0 | 85.5 | 363 | 0.00 | 0.021 | 0.00005 |
| 05-07-93 | 900 | 24.9 | 6.69 | 7.18 | | | | | | |
| 05-08-93 | 1640 | 25.2 | 6.57 | 5.25 | | | | | | |
| 05-09-93 | 1440 | 25.1 | 6.68 | 5.82 | | | | | | |
| 05-10-93 | 1000 | 24.9 | 6.62 | 6.89 | | | | | | |
| 05-11-93 | 900 | 25.0 | 6.64 | 7.35 | | | | | | |
| 05-12-93 | 1000 | 25.1 | 6.69 | 6.87 | | | | | | |
| 05-13-93 | 1000 | 25.1 | 6.70 | 7.09 | 34.0 | 102.6 | 354 | 0.00 | 0.000 | 0.00000 |
| 05-14-93 | 800 | 24.8 | 6.72 | 7.01 | | | | | | |
| 05-15-93 | 800 | 24.9 | 6.73 | 7.41 | | | | | | |
| 05-16-93 | 1100 | 25.1 | 6.71 | 7.60 | | | | | | |
| 05-17-93 | 900 | 24.9 | 6.62 | 7.49 | | | | | | |
| 05-18-93 | 1000 | 24.9 | 6.64 | 7.53 | | | | | | |
| 05-19-93 | 1000 | 25.2 | 6.74 | 7.15 | | | | | | |
| 05-20-93 | 900 | 25.3 | 6.70 | 6.90 | 34.0 | 102.6 | 356 | 0.00 | 0.057 | 0.00017 |
| 05-21-93 | 900 | 24.9 | 6.68 | 6.92 | | | | | | |
| 05-22-93 | 800 | 24.6 | 6.82 | 7.16 | | | | | | |
| 05-23-93 | 1000 | 24.5 | 6.76 | 7.05 | | | | | | |
| 05-24-93 | 800 | 24.7 | 6.68 | 7.01 | | | | | | |
| 05-25-93 | 1000 | 24.8 | 6.82 | 6.71 | | | | | | |
| 05-26-93 | 900 | 24.9 | 6.78 | 7.05 | | | | | | |
| 05-27-93 | 900 | 24.9 | 6.62 | 6.90 | 34.0 | 102.6 | 336 | 0.00 | 0.083 | 0.00020 |
| 05-28-93 | 1000 | 25.0 | 6.66 | 6.86 | | | | | | |
| 05-29-93 | 1200 | 25.2 | 6.67 | 7.09 | | | | | | |
| 05-30-93 | 900 | 25.0 | 6.73 | 6.41 | | | | | | |

| | | | | | | | | | | | |
|----------|-----|------|------|------|------|------|-------|-----|------|------|---------|
| 09-16-93 | 204 | 930 | 25.2 | 6.70 | 6.48 | 34.0 | 102.6 | 390 | 0.00 | 0.00 | 0.0000 |
| 09-17-93 | 205 | 930 | 24.5 | 6.66 | 6.57 | | | | | | |
| 09-18-93 | 206 | 800 | 24.8 | 6.77 | 6.83 | | | | | | |
| 09-19-93 | 207 | 1300 | 24.6 | 6.72 | 7.04 | | | | | | |
| 09-20-93 | 208 | 930 | 24.5 | 6.68 | 6.95 | | | | | | |
| 09-21-93 | 209 | 930 | 26.0 | 6.80 | 6.30 | | | | | | |
| 09-22-93 | 210 | 900 | 25.3 | 6.62 | 6.47 | | | | | | |
| 09-23-93 | 211 | 930 | 25.1 | 6.69 | 6.80 | 34.0 | 102.6 | 402 | 0.00 | 0.00 | 0.00000 |
| 09-24-93 | 212 | 1000 | 25.2 | 6.70 | 6.56 | | | | | | |
| 09-25-93 | 213 | 900 | 25.1 | 6.72 | 6.90 | | | | | | |
| 09-26-93 | 214 | 2100 | 25.6 | 6.78 | 6.93 | | | | | | |
| 09-27-93 | 215 | 1000 | 25.4 | 6.77 | 6.67 | | | | | | |
| 09-28-93 | 216 | 900 | 24.8 | 6.75 | 6.94 | | | | | | |
| 09-29-93 | 217 | 930 | 24.8 | 6.72 | 6.93 | 34.0 | 102.6 | 399 | 0.00 | 0.00 | 0.00001 |
| 09-30-93 | 218 | 1300 | 24.7 | 6.56 | 6.78 | | | | | | |
| 10-01-93 | 219 | 1100 | 24.2 | 6.67 | 7.89 | | | | | | |
| 10-02-93 | 220 | 830 | 24.3 | 6.71 | 6.90 | | | | | | |
| 10-03-93 | 221 | 830 | 25.0 | 6.72 | 6.69 | | | | | | |
| 10-04-93 | 222 | 1000 | 24.5 | 6.69 | 6.92 | | | | | | |
| 10-05-93 | 223 | 1000 | 24.9 | 6.65 | 6.65 | | | | | | |
| 10-06-93 | 224 | 930 | 24.5 | 6.77 | 7.09 | | | | | | |
| 10-07-93 | 225 | 930 | 24.9 | 6.82 | 6.95 | 40.8 | 102.6 | 398 | 0.00 | 0.00 | 0.00000 |
| 10-08-93 | 226 | 1000 | 24.9 | 6.77 | 6.91 | | | | | | |
| 10-09-93 | 227 | 1130 | 24.9 | 6.77 | 6.97 | | | | | | |
| 10-10-93 | 228 | 1130 | 24.6 | 6.77 | 7.13 | | | | | | |
| 10-11-93 | 229 | 1600 | 24.6 | 6.79 | 7.15 | | | | | | |
| 10-12-93 | 230 | 1000 | 24.5 | 6.74 | 7.05 | | | | | | |
| 10-13-93 | 231 | 930 | 24.7 | 6.76 | 6.96 | 34.0 | 102.6 | 395 | 0.00 | 0.00 | 0.00000 |
| 10-14-93 | 232 | 1000 | 24.6 | 6.74 | 6.95 | | | | | | |
| 10-15-93 | 233 | 1000 | 24.7 | 6.67 | 6.91 | | | | | | |
| 10-16-93 | 234 | 1230 | 24.7 | 6.67 | 6.88 | | | | | | |
| 10-17-93 | 235 | 1030 | 24.8 | 6.73 | 6.82 | | | | | | |
| 10-18-93 | 236 | 1000 | 24.7 | 6.71 | 6.83 | | | | | | |
| 10-19-93 | 237 | 900 | 24.7 | 6.67 | 6.87 | | | | | | |
| 10-20-93 | 238 | 1000 | 24.7 | 6.75 | 7.07 | | | | | | |
| 10-21-93 | 239 | 1000 | 24.8 | 6.76 | 6.95 | 34.0 | 102.6 | 398 | 0.00 | 0.00 | 0.00000 |
| 10-22-93 | 240 | 800 | 24.6 | 6.78 | 6.91 | | | | | | |
| 10-23-93 | 241 | 700 | 24.3 | 6.82 | 7.21 | | | | | | |
| 10-24-93 | 242 | 800 | 24.4 | 6.75 | 7.12 | | | | | | |
| 10-25-93 | 243 | 1000 | 24.4 | 6.66 | 6.47 | | | | | | |
| 10-26-93 | 244 | 1000 | 24.6 | 6.67 | 6.90 | | | | | | |
| 10-27-93 | 245 | 1000 | 24.3 | 6.76 | 6.91 | 34.0 | 102.6 | 382 | 0.00 | 0.00 | 0.00015 |
| 10-28-93 | 246 | 1000 | 24.8 | 6.77 | 6.77 | | | | | | |
| 10-29-93 | 247 | 1230 | 24.6 | 6.78 | 6.70 | | | | | | |
| 10-30-93 | 248 | 700 | 24.9 | 6.82 | 6.89 | | | | | | |
| 10-31-93 | 249 | 800 | 24.6 | 6.87 | 6.84 | | | | | | |
| 11-01-93 | 250 | 930 | 24.3 | 6.76 | 6.80 | | | | | | |
| 11-02-93 | 251 | 930 | 24.7 | 6.79 | 6.91 | | | | | | |
| 11-03-93 | 252 | 1000 | 24.9 | 6.74 | 6.77 | | | | | | |
| 11-04-93 | 253 | 1000 | 24.9 | 6.70 | 6.81 | 34.0 | 102.6 | 396 | 0.00 | 0.00 | 0.00000 |
| 11-05-93 | 254 | 1000 | 25.1 | 6.78 | 6.71 | | | | | | |
| 11-06-93 | 255 | 600 | 25.2 | 6.70 | 6.68 | | | | | | |
| 11-07-93 | 256 | 1000 | 24.9 | 6.74 | 6.63 | | | | | | |
| 11-08-93 | 257 | 1000 | 24.8 | 6.75 | 6.80 | | | | | | |

| | | | | | | | | | | | |
|----------|-----|------|------|-------|-------|------|-------|------|-------|--------|----------|
| 11-09-93 | 258 | 1000 | 24.6 | 6.74 | 6.82 | 34.0 | 102.6 | 387 | 0.00 | 0.003 | 0.00001 |
| 11-10-93 | 259 | 1000 | 24.7 | 6.70 | 6.70 | | | | | | |
| 11-11-93 | 260 | 1100 | 24.6 | 6.74 | 6.74 | | | | | | |
| 11-12-93 | 261 | 1000 | 24.9 | 6.71 | 6.57 | | | | | | |
| 11-13-93 | 262 | 1230 | 24.9 | 6.75 | 6.56 | | | | | | |
| 11-14-93 | 263 | 900 | 25.0 | 6.74 | 6.53 | | | | | | |
| 11-15-93 | 264 | 1000 | 25.1 | 6.71 | 6.47 | | | | | | |
| 11-16-93 | 265 | 1000 | 25.0 | 6.66 | 6.55 | | | | | | |
| 11-17-93 | 266 | 1200 | 25.2 | 6.66 | 5.93 | | | | | | |
| 11-18-93 | 267 | 1000 | 24.8 | 6.73 | 6.63 | 34.0 | 102.6 | 384 | 0.00 | 0.019 | 0.00006 |
| 11-19-93 | 268 | 1000 | 24.8 | 6.75 | 6.67 | | | | | | |
| 11-20-93 | 269 | 830 | 24.8 | 6.72 | 6.62 | | | | | | |
| 11-21-93 | 270 | 1300 | 24.5 | 6.71 | 6.82 | | | | | | |
| 11-22-93 | 271 | 930 | 24.5 | 6.71 | 6.78 | | | | | | |
| 11-23-93 | 272 | 1000 | 24.6 | 6.69 | 6.92 | | | | | | |
| 11-24-93 | 273 | 1300 | 24.5 | 6.80 | 6.28 | 34.0 | 102.6 | 393 | 0.00 | 0.044 | 0.00015 |
| 11-25-93 | 274 | 900 | 24.5 | 6.77 | 6.79 | | | | | | |
| 11-26-93 | 275 | 1200 | 24.4 | 6.70 | 6.88 | | | | | | |
| 11-27-93 | 276 | 1100 | 24.7 | 6.74 | 6.74 | | | | | | |
| 11-28-93 | 277 | 1100 | 25.2 | 6.72 | 6.45 | | | | | | |
| 11-29-93 | 278 | 600 | 24.0 | 7.38 | 7.43 | | | | | | |
| MEAN | | | 25.0 | 6.69 | 6.90 | 35.0 | 100.5 | 378 | 0.00 | 0.018 | 0.00005 |
| MINIMUM | | | 23.9 | 6.25 | 5.25 | 27.2 | 85.5 | 336 | 0.00 | 0.000 | 0.00000 |
| MAXIMUM | | | 26.9 | 7.38 | 8.22 | 47.6 | 102.6 | 565 | 0.00 | 0.111 | 0.00021 |
| Std Dev | | | 0.34 | 0.114 | 0.533 | 3.58 | 5.66 | 37.7 | 0.000 | 0.0256 | 0.000064 |
| N | | | 278 | 278 | 277 | 40 | 40 | 39 | 38 | 37 | 37 |

TANK No. 12
TANK CONCENTRATION: 10% GROUNDWATER (WELL CC-33B)
DEN EXPOSURE: 10 mg/L

| Date | Test Day | Time (Military) | Temperature (Celsius) | pH | Dissolved Oxygen (mg/L) | Alkalinity (mg/L as CaCO3) | Hardness (mg/L as CaCO3) | Conductivity (umohs/cm) | Total Residual Chlorine (mg/L) | Free Available Chlorine (mg/L) | Total Ammonia-Nitrogen (mg/L) | Total Unionized Ammonia-Nitrogen (mg/L) |
|----------|----------|-----------------|-----------------------|------|-------------------------|----------------------------|--------------------------|-------------------------|--------------------------------|--------------------------------|-------------------------------|---|
| 02-25-93 | 1 | 1500 | 24.4 | 6.47 | 7.55 | | | | | | | |
| 02-26-93 | 2 | 1000 | 23.9 | 6.63 | 7.61 | | | | | | | |
| 02-27-93 | 3 | 1000 | 24.4 | 6.83 | 7.39 | | | | | | | |
| 02-28-93 | 4 | 1100 | 24.6 | 6.88 | 7.28 | 34.0 | 102.6 | | | | | |
| 03-01-93 | 5 | 930 | 24.7 | 6.60 | 7.15 | | | | | | | |
| 03-02-93 | 6 | 1000 | 25.5 | 6.62 | 6.97 | | | | | | | |
| 03-03-93 | 7 | 1000 | 25.8 | 6.66 | 7.67 | | | | | | | |
| 03-04-93 | 8 | 1000 | 25.1 | 6.70 | 7.64 | | | | | | | |
| 03-05-93 | 9 | 800 | 25.4 | 6.70 | 7.34 | | | | | | | |
| 03-06-93 | 10 | 1000 | 24.9 | 6.51 | 7.60 | 27.2 | 102.6 | 464 | | | | |
| 03-07-93 | 11 | 1100 | 24.8 | 6.29 | 7.71 | | | | | | | |
| 03-08-93 | 12 | 900 | 24.9 | 6.39 | 7.59 | | | | | | | |
| 03-09-93 | 13 | 930 | 25.2 | 6.56 | 7.43 | | | | | | | |
| 03-10-93 | 14 | 900 | 24.8 | 6.49 | 7.42 | | | | | | | |
| 03-11-93 | 15 | 900 | 25.3 | 6.89 | 7.32 | | | | 0.00 | | | |
| 03-12-93 | 16 | 800 | 24.5 | 6.75 | 7.84 | 47.6 | 102.6 | 380 | | | | |
| 03-13-93 | 17 | 1600 | 24.8 | 6.67 | 7.53 | | | | | | | |
| 03-14-93 | 18 | 1500 | 24.8 | 6.73 | 7.49 | | | | | | | |
| 03-15-93 | 19 | 900 | 24.7 | 6.74 | 7.65 | | | | | | | |
| 03-16-93 | 20 | 900 | 25.2 | 6.79 | 7.67 | | | | | | | |
| 03-17-93 | 21 | 900 | 25.5 | 6.73 | 7.21 | 34.0 | 102.6 | 568 | 0.00 | 0.00 | 0.028 | 0.00008 |
| 03-18-93 | 22 | 1030 | 25.3 | 6.77 | 7.29 | | | | | | | |
| 03-19-93 | 23 | 1000 | 25.6 | 7.11 | 7.44 | | | | | | | |
| 03-20-93 | 24 | 1100 | 26.9 | 6.73 | 7.43 | | | | | | | |
| 03-21-93 | 25 | 1230 | 25.6 | 6.72 | 7.79 | | | | | | | |
| 03-22-93 | 26 | 900 | 25.0 | 6.72 | 7.77 | | | | | | | |
| 03-23-93 | 27 | 900 | 24.9 | 6.71 | 7.71 | | | | | | | |
| 03-24-93 | 28 | 900 | 25.3 | 6.65 | 7.57 | | | | | | | |
| 03-25-93 | 29 | 900 | 25.1 | 6.64 | 7.52 | 27.2 | 85.5 | 372 | 0.00 | 0.00 | 0.000 | 0.00000 |
| 03-26-93 | 30 | 800 | 24.9 | 6.25 | 7.75 | | | | | | | |
| 03-27-93 | 31 | 1000 | 25.2 | 6.27 | 7.51 | | | | | | | |
| 03-28-93 | 32 | 1000 | 25.6 | 6.38 | 7.47 | | | | | | | |
| 03-29-93 | 33 | 1100 | 25.7 | 6.45 | 7.39 | | | | | | | |
| 03-30-93 | 34 | 1100 | 25.5 | 6.39 | 7.59 | | | | | | | |
| 03-31-93 | 35 | 1000 | 25.5 | 6.46 | 5.76 | | | | | | | |
| 04-01-93 | 36 | 1000 | 24.8 | 6.50 | 7.66 | 34.0 | 85.5 | 359 | 0.00 | 0.00 | 0.000 | 0.00000 |
| 04-02-93 | 37 | 1000 | 24.7 | 6.51 | 7.72 | | | | | | | |
| 04-03-93 | 38 | 1100 | 24.8 | 6.53 | 7.90 | | | | | | | |
| 04-04-93 | 39 | 1300 | 24.7 | 6.46 | 7.99 | | | | | | | |
| 04-05-93 | 40 | 900 | 24.7 | 6.44 | 8.13 | | | | | | | |
| 04-06-93 | 41 | 1100 | 24.6 | 6.47 | 7.99 | | | | | | | |

| | | | | | | | | | | | |
|----------|----|------|------|------|------|------|------|-----|------|------|---------|
| 04-07-93 | 42 | 900 | 25.0 | 6.58 | 8.06 | 34.0 | 85.5 | 357 | 0.00 | 0.00 | 0.00000 |
| 04-08-93 | 43 | 900 | 25.2 | 6.58 | 7.94 | | | | | | |
| 04-09-93 | 44 | 900 | 25.3 | 6.57 | 7.80 | | | | | | |
| 04-10-93 | 45 | 1000 | 25.3 | 6.53 | 7.52 | | | | | | |
| 04-11-93 | 46 | 800 | 24.8 | 6.57 | 7.97 | | | | | | |
| 04-12-93 | 47 | 1000 | 25.2 | 6.59 | 7.78 | | | | | | |
| 04-13-93 | 48 | 900 | 25.0 | 6.56 | 7.59 | | | | | | |
| 04-14-93 | 49 | 600 | 25.1 | 6.60 | 7.35 | | | | | | |
| 04-15-93 | 50 | 900 | 25.3 | 6.58 | 7.36 | | | | | | |
| 04-16-93 | 51 | 1000 | 25.5 | 6.58 | 8.16 | | | | | | |
| 04-17-93 | 52 | 1000 | 25.3 | 6.58 | 7.33 | | | | | | |
| 04-18-93 | 53 | 1000 | 24.9 | 6.65 | 7.41 | | | | | | |
| 04-19-93 | 54 | 900 | 25.2 | 6.56 | 7.53 | | | | | | |
| 04-20-93 | 55 | 900 | 25.4 | 6.56 | 7.56 | | | | | | |
| 04-21-93 | 56 | 1000 | 25.4 | 6.61 | 7.31 | | | | | | |
| 04-22-93 | 57 | 900 | 24.9 | 6.61 | | | | | | | |
| 04-23-93 | 58 | 900 | 24.8 | 6.63 | 7.08 | | | | | | |
| 04-24-93 | 59 | 1000 | 25.1 | 6.60 | 7.59 | | | | | | |
| 04-25-93 | 60 | 800 | 25.2 | 6.59 | 7.79 | | | | | | |
| 04-26-93 | 61 | 900 | 25.3 | 6.60 | 7.08 | | | | | | |
| 04-27-93 | 62 | 1000 | 24.4 | 6.70 | 7.70 | | | | | | |
| 04-28-93 | 63 | 900 | 25.0 | 6.60 | 7.32 | | | | | | |
| 04-29-93 | 64 | 1000 | 25.0 | 6.65 | 7.17 | | | | | | |
| 04-30-93 | 65 | 1000 | 25.2 | 6.64 | 7.17 | | | | | | |
| 05-01-93 | 66 | 1000 | 25.4 | 6.62 | 7.90 | | | | | | |
| 05-02-93 | 67 | 1000 | 25.5 | 6.62 | 7.71 | | | | | | |
| 05-03-93 | 68 | 1000 | 25.4 | 6.61 | 7.82 | | | | | | |
| 05-04-93 | 69 | 1000 | 25.6 | 6.65 | 7.22 | | | | | | |
| 05-05-93 | 70 | 900 | 25.5 | 6.66 | 7.16 | | | | | | |
| 05-06-93 | 71 | 1000 | 25.1 | 6.65 | 7.09 | | | | | | |
| 05-07-93 | 72 | 900 | 24.9 | 6.68 | 7.23 | | | | | | |
| 05-08-93 | 73 | 1640 | 25.2 | 6.56 | 5.38 | | | | | | |
| 05-09-93 | 74 | 1440 | 25.1 | 6.61 | 5.29 | | | | | | |
| 05-10-93 | 75 | 1000 | 24.9 | 6.62 | 6.94 | | | | | | |
| 05-11-93 | 76 | 900 | 25.0 | 6.64 | 7.36 | | | | | | |
| 05-12-93 | 77 | 1000 | 25.1 | 6.66 | 6.90 | | | | | | |
| 05-13-93 | 78 | 1000 | 25.0 | 6.69 | 7.15 | | | | | | |
| 05-14-93 | 79 | 800 | 24.8 | 6.72 | 7.01 | | | | | | |
| 05-15-93 | 80 | 800 | 24.8 | 6.73 | 7.41 | | | | | | |
| 05-16-93 | 81 | 1100 | 25.1 | 6.71 | 7.61 | | | | | | |
| 05-17-93 | 82 | 900 | 24.8 | 6.62 | 7.55 | | | | | | |
| 05-18-93 | 83 | 1000 | 24.8 | 6.64 | 7.62 | | | | | | |
| 05-19-93 | 84 | 1000 | 25.1 | 6.74 | 7.20 | | | | | | |
| 05-20-93 | 85 | 900 | 25.3 | 6.69 | 6.96 | | | | | | |
| 05-21-93 | 86 | 900 | 24.9 | 6.68 | 7.14 | | | | | | |
| 05-22-93 | 87 | 800 | 24.6 | 6.81 | 7.33 | | | | | | |
| 05-23-93 | 88 | 1000 | 24.5 | 6.74 | 7.17 | | | | | | |
| 05-24-93 | 89 | 800 | 24.7 | 6.67 | 7.12 | | | | | | |
| 05-25-93 | 90 | 1000 | 24.8 | 6.82 | 6.76 | | | | | | |
| 05-26-93 | 91 | 900 | 24.9 | 6.78 | 7.13 | | | | | | |
| 05-27-93 | 92 | 900 | 24.9 | 6.66 | 6.97 | | | | | | |
| 05-28-93 | 93 | 1000 | 25.0 | 6.66 | 6.95 | | | | | | |
| 05-29-93 | 94 | 1200 | 25.1 | 6.67 | 7.22 | | | | | | |
| 05-30-93 | 95 | 900 | 25.0 | 6.73 | 6.60 | | | | | | |

| | | | | | | | | | | | |
|----------|-----|------|------|------|------|------|-------|-----|------|------|---------|
| 09-16-93 | 204 | 930 | 25.2 | 6.70 | 6.44 | 34.0 | 102.6 | 402 | 0.00 | 0.00 | 0.0000 |
| 09-17-93 | 205 | 930 | 24.6 | 8.66 | 6.50 | | | | | | |
| 09-18-93 | 206 | 800 | 24.8 | 6.77 | 6.78 | | | | | | |
| 09-19-93 | 207 | 1300 | 24.5 | 6.72 | 7.10 | | | | | | |
| 09-20-93 | 208 | 930 | 24.4 | 6.68 | 6.87 | | | | | | |
| 09-21-93 | 209 | 930 | 25.9 | 8.80 | 6.25 | | | | | | |
| 09-22-93 | 210 | 900 | 25.2 | 8.61 | 6.48 | | | | | | |
| 09-23-93 | 211 | 930 | 25.1 | 8.69 | 6.81 | 34.0 | 102.6 | 412 | 0.00 | 0.00 | 0.00000 |
| 09-24-93 | 212 | 1000 | 25.2 | 6.70 | 6.61 | | | | | | |
| 09-25-93 | 213 | 900 | 25.0 | 6.72 | 6.92 | | | | | | |
| 09-26-93 | 214 | 2100 | 25.6 | 6.78 | 7.01 | | | | | | |
| 09-27-93 | 215 | 1000 | 25.3 | 6.77 | 6.73 | | | | | | |
| 09-28-93 | 216 | 900 | 24.8 | 6.75 | 6.91 | | | | | | |
| 09-29-93 | 217 | 930 | 24.8 | 6.72 | 6.91 | 34.0 | 102.6 | 407 | 0.00 | 0.00 | 0.00003 |
| 09-30-93 | 218 | 1300 | 24.6 | 6.56 | 6.91 | | | | | | |
| 10-01-93 | 219 | 1100 | 24.2 | 8.66 | 8.09 | | | | | | |
| 10-02-93 | 220 | 830 | 24.7 | 6.70 | 6.91 | | | | | | |
| 10-03-93 | 221 | 830 | 24.9 | 6.72 | 6.71 | | | | | | |
| 10-04-93 | 222 | 1000 | 24.5 | 8.69 | 6.92 | | | | | | |
| 10-05-93 | 223 | 1000 | 24.9 | 8.65 | 6.68 | | | | | | |
| 10-06-93 | 224 | 930 | 24.5 | 6.77 | 7.08 | | | | | | |
| 10-07-93 | 225 | 930 | 24.9 | 8.82 | 6.97 | 40.8 | 102.6 | 407 | 0.00 | 0.00 | 0.00000 |
| 10-08-93 | 226 | 1000 | 24.9 | 8.77 | 7.01 | | | | | | |
| 10-09-93 | 227 | 1130 | 24.9 | 8.77 | 7.04 | | | | | | |
| 10-10-93 | 228 | 1130 | 24.6 | 6.77 | 7.14 | | | | | | |
| 10-11-93 | 229 | 1600 | 24.6 | 6.79 | 7.28 | | | | | | |
| 10-12-93 | 230 | 1000 | 24.5 | 6.74 | 7.12 | | | | | | |
| 10-13-93 | 231 | 930 | 24.7 | 8.76 | 7.03 | | | | | | |
| 10-14-93 | 232 | 1000 | 24.6 | 6.74 | 7.04 | 34.0 | 102.6 | 403 | 0.00 | 0.00 | 0.00000 |
| 10-15-93 | 233 | 1000 | 24.7 | 6.67 | 6.99 | | | | | | |
| 10-16-93 | 234 | 1230 | 24.6 | 6.67 | 7.00 | | | | | | |
| 10-17-93 | 235 | 1030 | 24.8 | 6.73 | 6.91 | | | | | | |
| 10-18-93 | 236 | 1000 | 24.7 | 8.69 | 6.89 | | | | | | |
| 10-19-93 | 237 | 900 | 24.7 | 8.67 | 7.03 | | | | | | |
| 10-20-93 | 238 | 1000 | 24.7 | 8.75 | 7.21 | | | | | | |
| 10-21-93 | 239 | 1000 | 24.7 | 6.76 | 7.04 | 34.0 | 102.6 | 405 | 0.00 | 0.00 | 0.00000 |
| 10-22-93 | 240 | 800 | 24.6 | 8.79 | 6.84 | | | | | | |
| 10-23-93 | 241 | 700 | 24.3 | 6.80 | 7.27 | | | | | | |
| 10-24-93 | 242 | 800 | 24.3 | 8.75 | 7.18 | | | | | | |
| 10-25-93 | 243 | 1000 | 24.3 | 6.65 | 6.54 | | | | | | |
| 10-26-93 | 244 | 1000 | 24.6 | 6.67 | 7.09 | | | | | | |
| 10-27-93 | 245 | 1000 | 24.3 | 6.76 | 7.08 | | | | | | |
| 10-28-93 | 246 | 1000 | 24.8 | 6.78 | 7.02 | 34.0 | 102.6 | 386 | 0.00 | 0.00 | 0.00016 |
| 10-29-93 | 247 | 1230 | 24.6 | 6.78 | 6.91 | | | | | | |
| 10-30-93 | 248 | 700 | 24.9 | 6.82 | 7.00 | | | | | | |
| 10-31-93 | 249 | 800 | 24.6 | 6.87 | 6.97 | | | | | | |
| 11-01-93 | 250 | 930 | 24.3 | 6.76 | 7.00 | | | | | | |
| 11-02-93 | 251 | 930 | 24.7 | 8.79 | 6.97 | | | | | | |
| 11-03-93 | 252 | 1000 | 24.9 | 6.74 | 6.90 | | | | | | |
| 11-04-93 | 253 | 1000 | 24.9 | 6.70 | 6.91 | 34.0 | 102.6 | 401 | 0.00 | 0.00 | 0.00000 |
| 11-05-93 | 254 | 1000 | 25.1 | 6.78 | 6.81 | | | | | | |
| 11-06-93 | 255 | 600 | 25.2 | 8.70 | 6.69 | | | | | | |
| 11-07-93 | 256 | 1000 | 24.9 | 8.74 | 6.75 | | | | | | |
| 11-08-93 | 257 | 1000 | 24.8 | 8.75 | 6.90 | | | | | | |

| | | | | | | | | | | | |
|----------|-----|------|------|-------|-------|------|-------|------|-------|-------|----------|
| 11-09-93 | 258 | 1000 | 24.7 | 6.74 | 6.94 | 34.0 | 102.6 | 387 | 0.00 | 0.00 | 0.00000 |
| 11-10-93 | 259 | 1000 | 24.7 | 6.70 | 6.82 | | | | | | |
| 11-11-93 | 260 | 1100 | 24.6 | 6.74 | 6.84 | | | | | | |
| 11-12-93 | 261 | 1000 | 24.8 | 6.72 | 6.72 | | | | | | |
| 11-13-93 | 262 | 1230 | 24.9 | 6.75 | 6.68 | | | | | | |
| 11-14-93 | 263 | 900 | 25.0 | 6.74 | 6.66 | | | | | | |
| 11-15-93 | 264 | 1000 | 25.1 | 6.71 | 6.56 | | | | | | |
| 11-16-93 | 265 | 1000 | 24.9 | 6.66 | 6.68 | | | | | | |
| 11-17-93 | 266 | 1200 | 25.1 | 6.66 | 6.01 | | | | | | |
| 11-18-93 | 267 | 1000 | 24.9 | 6.73 | 6.76 | 34.0 | 102.6 | 381 | 0.00 | 0.015 | 0.00004 |
| 11-19-93 | 268 | 1000 | 24.9 | 6.75 | 6.72 | | | | | | |
| 11-20-93 | 269 | 830 | 24.9 | 6.72 | 6.58 | | | | | | |
| 11-21-93 | 270 | 1300 | 24.5 | 6.71 | 6.99 | | | | | | |
| 11-22-93 | 271 | 930 | 24.5 | 6.71 | 6.79 | | | | | | |
| 11-23-93 | 272 | 1000 | 24.6 | 6.69 | 6.99 | 34.0 | 102.6 | 401 | 0.00 | 0.041 | 0.00014 |
| 11-24-93 | 273 | 1300 | 24.6 | 6.80 | 6.41 | | | | | | |
| 11-25-93 | 274 | 900 | 24.6 | 6.77 | 6.91 | | | | | | |
| 11-26-93 | 275 | 1200 | 24.4 | 6.70 | 7.07 | | | | | | |
| 11-27-93 | 276 | 1100 | 24.7 | 6.74 | 6.84 | | | | | | |
| 11-28-93 | 277 | 1100 | 25.2 | 6.72 | 6.59 | | | | | | |
| 11-29-93 | 278 | 600 | 24.1 | 7.38 | 7.42 | | | | | | |
| MEAN | | | 25.0 | 6.69 | 6.93 | 34.9 | 100.9 | 380 | 0.00 | 0.00 | 0.00004 |
| MINIMUM | | | 23.9 | 6.25 | 5.29 | 27.2 | 85.5 | 344 | 0.00 | 0.00 | 0.00000 |
| MAXIMUM | | | 26.9 | 7.38 | 8.16 | 47.6 | 102.6 | 568 | 0.00 | 0.00 | 0.00021 |
| Std Dev | | | 0.33 | 0.116 | 0.514 | 3.78 | 5.13 | 39.8 | 0.000 | 0.000 | 0.000061 |
| N | | | 278 | 278 | 277 | 40 | 40 | 39 | 38 | 37 | 37 |

APPENDIX 50

ROUTINE WATER QUALITY CHEMICAL METHODS FOR ALL SAMPLES ANALYZED IN THE BIOMONITORING TRAILER AND AT THE UMD/WREC LABORATORY

| Parameter | Method ^a |
|------------------|--|
| Alkalinity | Method 2320 B. Titration Method |
| Ammonia-nitrogen | Method 4500-NH ₃ . Ammonia Selective Electrode Method |
| Chlorine | Method 4500-Cl G. DPD Colorimetric Method |
| Conductivity | Method 2510 B. Laboratory Method |
| Dissolved Oxygen | Method 4500-O G. Membrane Electrode Method |
| Hardness | Method 2340 C. EDTA Titrimetric Method |
| pH | Method 4500-H ⁺ B. Electrometric Method |
| Temperature | Method 2550 B. Laboratory and Field Methods |

^a All methods taken from Standard Methods (APHA et al., 1989).

APPENDIX 51

SCREENING-LEVEL ASSESSMENT OF THE NEAR- AND FAR-FIELD
DILUTION OF THE SURFICIAL GROUNDWATER DISCHARGE
FROM BEACH POINT INTO THE BUSH RIVER

FINAL REPORT - JOB 1437

SCREENING-LEVEL ASSESSMENT OF THE NEAR-FIELD DILUTION OF GROUNDWATER DISCHARGES FROM THE BEACH POINT SITE

Prepared for:
U.S. ARMY BIOMEDICAL RESEARCH AND DEVELOPMENT LABORATORY
FORT DETRICK, BUILDING 568
FREDERICK MD 21702-5010

Prepared by:
NAJARIAN ASSOCIATES, INC.
ONE INDUSTRIAL WAY WEST
EATONTOWN, NEW JERSEY 07724

JULY, 1994

EXECUTIVE SUMMARY

This study was initiated to assess the potential impact to the Bush River due to contaminated groundwater emanating from the Army Proving Ground site in Harford County, Maryland. Towards this objective, an EPA-approved near-field model was adapted to the receiving-waters adjacent to the APG site. The model approach and input was developed to provide a conservative assessment of receiving water impacts. This analysis suggests a near-field dilution level of approximately 42:1 for the application of acute criteria and a near-field dilution level of approximately 168:1 for the application of chronic criteria. Thus, contaminants introduced into Bush River receiving-waters at a concentration of 42 ppb will be diluted locally to a concentration of approximately 1 ppb or less. By applying these dilution factors to the groundwater quality data from the APG site, near-field concentrations within the Bush River can be projected. These results indicate that Tables 8 and 9, none of the projected receiving water concentrations would exceed Maryland's current acute or chronic criteria.

TABLE OF CONTENTS

| | PAGE |
|---|-----------|
| 1. Introduction | |
| 1.1 Background..... | 1 |
| 1.2 Objectives..... | 2 |
| 1.3 Study Area..... | 2 |
| 2. Part I: Near-Field Model Study | |
| 2.1 Approach..... | 8 |
| 2.2 Near-Field Model Selection | 9 |
| 2.3 Estimation of Groundwater Input Parameters | 10 |
| 2.4 Ambient Receiving Water Input..... | 11 |
| 2.5 Conceptualized Discharge Schematization..... | 13 |
| 2.6 Near-field Model Results..... | 13 |
| 2.7 Model Sensitivity Analysis | 14 |
| 3. Part II: Far-field Dilution Estimates | |
| 3.1 Definitions..... | 21 |
| 3.2 Tidal Prism Method | 21 |
| 3.3 Fraction of Freshwater Method..... | 24 |
| 3.4 Previous Dye-Tracer Study Results | 24 |
| 3.5 Calculation of Total Near-Field Dilution..... | 26 |
| 3.6 Application of Dilution Factors to Toxic Discharge Data | 27 |
| 4. Part III: Mixing Zone Considerations | |
| 4.1 Mixing Zone Regulations | 30 |
| 4.2 Site Applicability | 33 |
| 4.3 Conformance with Criteria | 35 |
| 5. Summary and Conclusions | 38 |

LIST OF TABLES

| | PAGE |
|---|------|
| 1. Sediment Metal Concentrations in the Bush River | 7 |
| 2. Sediment PAH and TOC Data for the Bush River | 7 |
| 3. Calculation Table for Tidal Prism Method | 23 |
| 4. Calculation Table for Fraction of Freshwater Method | 25 |
| 5. Computed Dilutions of Volatile Organics | 28 |
| 6. Computed Dilutions of Heavy Metals | 29 |
| 7. Toxic Substances Criteria for Ambient Surface Waters (Fresh Water) | 32 |
| 8. Comparison of Near-Field Concentrations with Acute Criteria | 36 |
| 9. Comparison of Near-Field Concentrations with Chronic Criteria | 37 |

LIST OF FIGURES

PAGE

1. Bush River Navigation Chart3

2. Simulated Dilution vs. Discharge Rate15

3. Simulated Dilution vs. Discharge Length16

4 Simulated Dilution vs. Discharge Depth.....18

5. Simulated Dilution vs. Ambient Current19

6. Simulated Dilution vs. Ambient Salinity.....20

1. INTRODUCTION

1.1 Background

The U.S Army Aberdeen Proving Ground (APG) is located between the Bush River and the Upper Chesapeake Bay in Harford County, Maryland. Currently, the U.S. Army Biomedical Research and Development Laboratory (USABRDL) is conducting biomonitoring evaluation studies of groundwater contamination at the APG site. To complement such studies, the USABRDL requires information regarding potential impacts of such groundwater contamination on Bush River water quality. Such impacts depend upon the ability of these receiving waters to mix and dilute contaminants introduced by groundwater sources. Accordingly, Najarian Associates, Inc. was contracted by the USABRDL to conduct preliminary model studies of groundwater-plume dilution in the Bush River.

The initial dilution of groundwater-discharge plumes depends, in part, upon the physical characteristics of both the source and the ambient receiving waters. These characteristics include the quantity/quality of the discharge, the geometrical configuration of the source, the ambient current speed, the ambient density stratification, and the water depth. These latter three factors control the near-field mixing of discharged constituents with surrounding receiving-waters, primarily through the mechanisms of plume buoyancy and initial discharge momentum. Typically, all such factors are incorporated into plume-dilution models, including simplified "screening-level" approaches.

1.2 Objectives


The basic goal of this study is to estimate, using EPA-approved screening-level

techniques, the initial ("near-field") dilution and flushing rate ("far-field" dilution) of groundwater discharge plumes within the adjacent surface waters. Such an assessment, when combined with relevant discharge concentration data, would provide first-order estimates of resulting near-field concentrations — receiving-water concentrations that would be expected over relatively short distances from the discharge site (e.g., ones, tens and hundreds of meters) and over short time periods (e.g., seconds and minutes). Our second objective is to compute such concentrations for selected toxic constituents and to compare the results to available baseline data. As a result, potential water quality impacts may be quantified. A third study goal is to assess the applicability of a "regulatory mixing zone" — a localized discharge zone in which local water quality standards may be exceeded. Such zones are routinely allowed for wastewater treatment plant discharges. If applicable to the subject groundwater-discharge plume, a mixing zone may allow for local exceedences of water quality standards.

1.3 Study Area

The goals and procedures outlined above are based on relevant characteristics of the study area. Surface elevations at the APG Beach Point site lie approximately 14.5 feet above sea level. An unsaturated zone occurs from elevation 14.5 ft down and at depths as shallow as 1 ft. The site is underlain by an impermeable clay lens located at a depth ranging from approximately 70 feet to 120 ft. A lower confined aquifer known as the Canal Creek Aquifer occurs below the upper confining unit.

The Bush River is a relatively small and shallow tributary estuary located on the western shore of the Upper Chesapeake Bay (Figure 1). The Bush River has a length of



approximately 14 km. Bush River MLW depths generally increase from an approximate 0.9-1.5 m range (near the head) to a 2.7-4.0 m range (near the mouth). The average Mean-Low-Water (MLW) depth is approximately 2 m. The average annual freshwater

discharge to the Bush River from its 316-km² drainage area is only about 4.7 m³/sec (Pritchard, 1976).

Bush River tides are predominantly semi-diurnal (NOAA, 1978), with a mean range at Pond Point of 0.4 m and a spring range of approximately 0.5 m (NOS, 1993a). Inside the entrance to the Bush River, about 2 miles south of Beach Point, average maximum tidal current speeds are approximately 0.3 m/sec (1.0 ft/sec; NOS, 1993).

The MLW volume of the Bush River is approximately 6.0×10^7 m³; the inter-tidal volume (i.e., the "tidal prism") is about 1.2×10^7 m³ (Pritchard, 1976). Thus, the inter-tidal volume comprises a relatively large fraction (approximately 20%) of the River's MLW volume. In contrast, the mean runoff volume per day (4.1×10^5 m³/day) corresponds to a relatively small fraction of the River's MLW volume. Pritchard (1976) estimates that the MLW volume corresponds to approximately 150 days of mean freshwater inflow.

Due to its relatively small areal extent, depth, and freshwater inflow, the Bush River typically exhibits relatively small horizontal and vertical salinity gradients (Carter, 1976). For example, observations of system-wide salinity variations of only 3 ppt are reported in the Bush River (Martin Marietta, 1986; Carter, 1976). The longitudinal salinity difference is controlled primarily by the adjacent Chesapeake Bay salinities and secondarily by freshwater inflows to the Bush River from Otter Point Creek (Carter, 1976). Consequently, Bush River salinities do not always decrease monotonically in a

landward direction. Intratidal variations of Bush River salinities range from approximately 0.5 ppt near the mouth to near zero at the head (Carter, 1976).

Susquehanna River inflows to the Upper Chesapeake Bay influence the salinity distribution and circulation of the Bush River and other small tributary estuaries (Pritchard, 1976). As noted by Pritchard (1969), salinities of these embayments are controlled largely by the relatively low salinities of the Upper Bay and by Upper Bay variability. For example, Martin Marietta (1986) reports observations of Bush River salinities varying from 0.7 ppt (at the River's head-of-tide) to 3.7 ppt (at River mouth), consistent with prevailing local Bay salinities. However, the salinity of the Upper Bay and its tributaries also varies in response to seasonal changes in Susquehanna River flows, as described by Pritchard (1976).

During the autumn, when Susquehanna River flows are relatively low, Upper Chesapeake Bay salinities are constant or slowly increasing. At this time, the local freshwater inflow to the Bush River may be sufficient to set up an estuarine type gravitational circulation, with a net outflow of fresher surface waters and net inflow of more saline bottom waters. During the spring freshet, when salinities of the Upper Bay decrease over time, corresponding salinities of the tributary embayments (e.g., the Bush River) are relatively high. Consequently, a reverse estuarine circulation pattern may develop, with a net bottom outflow of more saline water from the River to the Bay proper, and a net surface inflow from the Bay proper into the River. Such seasonal gravitational circulation patterns enhance the flushing of the Bush River and other tributary embayments. Thus, the available far-field dilution would be greater at these times.

The Bush River at the Beach Point site is classified as a *Designated Use II: Shellfish Harvesting Waters* (COMAR 26.08.02.02B) — an actual or potential area for shellfish

propagation. Approximately one mile upstream from Beach Point, the Bush River classification changes to *Designated Use I: Water Contact Recreation, and Protection of Aquatic Life* (i.e., the basic swimmable/fishable classification).

Toxic water quality conditions in the Bush River are reported in a recent study of contaminants in Chesapeake Bay/Tributary sediments conducted by the Maryland Department of the Environment (Eskin et al., 1994). The MDE's toxics monitoring program includes a Bush River sampling station near Gum Point (Figure 1, 39°26'06"N, 76°14'36"W), approximately 5 km (3 miles) north of the Beach Point site. Annual (1988-91) sediment metals concentrations for this stations are reproduced below in Table 1; individual observations of sediment PAH and TOC for 1991 are reproduced in Table 2. In the tables, both the No Observable Effect Level (NOEL) and the Probable Effects Level (PEL) are listed for each parameter, where available. Such levels provide guidance regarding the potential toxicity of these parameters (MacDonald, 1993).

As indicated in the MDE report, most sediment concentrations observed in the Bush River fall above the NOEL but below the PEL. Thus, toxic effects due to each trace metal are unlikely (individually). However, compared to other tributary stations, the Bush River Nickel levels are somewhat high.

Where sediment quality guidelines are available, the data also suggest that toxicity due to individual PAH and TOC compounds is unlikely. In fact, most Bush River PAH concentrations observed in the MDE's 1991 survey fall below the corresponding NOELs.

TABLE 1: SEDIMENT METAL CONCENTRATIONS IN THE BUSH RIVER

| Parameter | Guideline Limits (ppm) | | Observed Concentrations (ppm) | | | |
|-----------|------------------------|-----|-------------------------------|-------|-------|-------|
| | NOEL | PEL | 1988 | 1989 | 1990 | 1991 |
| As | 8 | 64 | 34.37 | 45.80 | 38.30 | 19.70 |
| Cd | 1 | 7.5 | 0.01 | 0.15 | 0.40 | 0.50 |
| Cr | 33 | 240 | 68.92 | 69.00 | 82.00 | 59.00 |
| Cu | 28 | 270 | 43.32 | 45.00 | 47.50 | 40.00 |
| Pb | 21 | 160 | 42.25 | 66.00 | 42.60 | 54.00 |
| Hg | 0.1 | 1.4 | 0.099 | 0.244 | 0.150 | 0.190 |
| Ni | — | — | 53.79 | 57.00 | 47.10 | 52.00 |
| Zn | 68 | 300 | 205.0 | 216.0 | 222.3 | 189.0 |

TABLE 2: SEDIMENT PAH AND TOC DATA FOR THE BUSH RIVER

| Parameter | Guideline Limits (ppb) | | Observed Concentrations (ppb) |
|------------------------|------------------------|-------|-------------------------------|
| | NOEL | PEL | |
| anthracene | 85 | 740 | 75.0 |
| fluoranthene | 380 | 3,200 | 24.4 |
| pyrene | — | — | 224.9 |
| benzo(A)anthracene | 160 | 1,300 | 75.0 |
| benzo(B)fluoranthene | — | — | 187.4 |
| benzo(K)fluoranthene | — | — | 56.2 |
| benzo(A)pyrene | 230 | 2,500 | 112.5 |
| benzo(G, H, 1)perylene | — | — | 150.0 |

2. NEAR-FIELD MODEL STUDY

2.1 Approach

In this preliminary assessment, the impact of groundwater emanating from the APG site was evaluated based on the potential dilution available within the Bush River. Within the near-field, it is postulated that a groundwater plume intersects the channel of the Bush River and seeps into this River as a submerged buoyant plume. Based on the available geologic information about the APG site, it is also postulated that the discharge from the surface aquifer is limited to a relatively narrow band that parallels the interface between the Bush River and the subject APG site. As this groundwater plume is less dense (less saline) than the adjacent estuarine water, it would rise and subsequently mix with the ambient receiving-waters. Thus, it is diluted through mechanisms of buoyancy-based entrainment and mixing associated with ambient currents.

These same physical mechanisms are known to dilute effluent plumes discharged from the outfall diffusers of wastewater treatment plants (WWTP) and electrical generation facilities. Thus, the dilution of groundwater emanating from the APG site can be conceptualized as the dilution of the discharge from a line diffuser. This approach allows the near-field impact of this groundwater discharge to be simulated using established plume-dilution models that have been developed primarily for such outfall diffusers.

The limitations of this approach are twofold. First, most plume-dilution models also simulate momentum-based entrainment, a phenomenon that has a negligible effect upon the groundwater plume. Therefore, the port configuration on the

conceptualized line diffuser must be set to minimize initial momentum. Secondly, the conceptualization of the groundwater discharge surface as a line diffuser will concentrate the plume's impact within the near-field condition. As this effect reduces the dimensions of the near-field plume and the calculated levels of dilution, the model result will provide a conservative estimate of field conditions.

2.2 Near-Field Model Selection

A series of EPA-approved, plume-dilution models for surface waters are documented in "Initial Mixing Characteristics of Municipal Ocean Discharges" by Muellenhoff et al. (EPA/600/3-85/073a, 1985) and by reviewed by Alam et al. (1982). These models provide a convenient and accepted choice of screening-level techniques. As noted above, such models are usually applied to discharges from line sources such as outfall diffuser manifold legs. In the present case, however, the subject discharges are emanating primarily from the adjacent stream bed.

While all EPA-approved plume models can be run, ULINE is preferred for dilution calculations since it is based directly on laboratory experiments and has proved its usefulness in recent outfall studies conducted throughout the U.S.(e.g., Najarian Assoc., 1990). The ULINE model also neglects initial discharge momentum, consistent with the problem at hand.

Typically, outfall dilution studies are conducted under "critical" (minimum-dilution; Tetrattech, 1982) conditions. These conditions are defined as lowest 10th-percentile ambient currents, and highest 10th-percentile ambient density stratification conditions. However, Maryland regulations require that the acute toxicity criteria be applied under conditions of mean low water and "minimum daily averaged

1-hour tidal velocity" (COMAR 26.08.02.05C); chronic criteria are applied under conditions of mean water level and average tidal velocity (COMAR 26.08.02.05D).

Thus, the input data for the selected near-field model (ULINE) should reflect these required conditions.

To adapt ULINE to the study area, an input of site-specific data is required. Such model input data fall into three categories: (1) groundwater discharge parameters; (2) receiving-water parameters; and (3) outfall diffuser parameters. The selection of such parameters is described below:

2.3 Estimation of Groundwater Input Parameters

Information relating the site's local geology and hydrogeologic conditions was obtained from previous studies by Jacobs Engineering Group and K-V Associates, Inc. (1994). These studies indicate that subsurface flows within the surficial aquifer are bounded by a sub-surface clay layer and that the portion of the aquifer underlying the Beach Point peninsula is both homogeneous and isotropic. On-site data indicates that the dominate, long-term groundwater flow direction would be down-gradient towards the Bush River.

Based on K-V Associates measurements and Jacobs Engineering calculations (Jacobs Engineering Group, 1994), the average saturated thickness of the surficial aquifer at the Beach Point peninsula is approximately 18.8 m (61.8 ft). Also, the Bush River frontage along the subject peninsula is approximately 448 m (1,470 ft). Approximately 366 m (1,200 ft) of frontage is directed in a northeast direction; approximately 82 m (270 ft) of the northern peninsula terminus is directed in a

northwest direction. Thus, the cross-section area of the saturated aquifer that parallels the Bush River is approximately 8,423 m² (90,846 ft²).

As conditions within the surficial aquifer are homogeneous and isotropic, horizontal velocity measurements at a single location may be used to evaluate the aquifer discharge. K-V Associates measured a 25-hour average horizontal velocity of 0.27 m/day (0.89 ft/day) at a 15-m deep well site located near the middle of the Beach Point Peninsula. This flow was directed 61.8° northeast of magnetic north, approximately 24° riverward from the main peninsula axis. Note that the above velocity measurement is a 24-hour average value (discounting short-term fluctuations due to tidal influences).

Jacobs Engineering estimates the groundwater seepage rate to be 0.0136 m³/sec (41,509 ft³/day). This estimate utilizes the directional component of the horizontal velocity that is *orthogonal* to the Bush River frontage. That is, the measured horizontal velocity (i.e., 0.27 m/day) is multiplied by a sin(24°) factor along the 366 m (1,200-ft) frontage:

$$\begin{aligned}\text{Discharge} &= (0.27 \text{ m/day velocity})(\sin 24^\circ)(18.8 \text{ m depth})(366 \text{ m frontage}) \\ &\quad + (0.27 \text{ m/day velocity})(18.8 \text{ m depth})(82 \text{ m frontage}) \\ &= 0.0136 \text{ m}^3/\text{sec}\end{aligned}$$

2.4 Ambient Receiving Water Input

Information relating the hydrographic conditions within the Bush River were obtained from a review of the NOAA Tidal Current Table data, NOAA navigation charts, and previous studies of the Bush River by Martin Marietta (1986).

Ambient Current Velocity in Bush River

A review of the NOAA Tidal Current Table data at a location of 0.6 km southwest of Bush Point indicates that the maximum flood velocity is 0.305 m/sec (1.0 ft/sec or 0.6 knot). Assuming a pure sinusoidal tidal velocity distribution, having a 0.305 m/sec amplitude and 12.4 hour period, the "minimum daily average 1-hour tidal velocity was computed. That is, the assumed tidal current magnitude profile was averaged over a 1-hour window centered at slack tide (approximately hour 6.2). The resulting speed was estimated to be 0.04 m/sec (see Appendix A) . Thus, as a conservative estimate, an ambient current speed of 0.03 m/sec was input to the near-field model as a representative acute condition tidal current speed. Likewise, for the same sinusoidal profile, a tidal average speed was computed as 0.19 m/sec. This latter average is representative of the chronic condition speed.

Ambient salinity and temperature in Bush River

A representative receiving-water salinity of 3 ppt is assumed in this study, consistent with the limited available data (e.g., Martin Marietta, 1986). Similarly, a temperature of 21° C is assumed for the Bush River. This salinity and temperature correspond to a receiving-water density of only 1.00031 g/cm³. Thus the selected values are conservative, inasmuch as only limited plume buoyancy will be available to dilute the discharges. Note that the assumed density of the groundwater discharges is 1.00024 g/cm³ (i.e., only 0.007% lighter than the ambient receiving waters).

Depth of discharge

Based on the most recent NOAA navigation chart for the Bush River, an average water depth of 1.5 m (5 ft) is assumed at the Bush River discharge site. However, a more conservative estimate of half this depth, 0.76 m (2.5 ft), is chosen as a representative model discharge depth. This selection reflects the fact that some of the discharge occurs along the shallow frontage of the peninsula.

2.5 Conceptualized Discharge Schematization

As noted above, the 448-m-wide groundwater discharge plume at the Beach Point site seeps into the adjacent Bush River. Thus, this discharge is assumed to comprise a 448-m-long line source of buoyancy. By analogy, the corresponding plume in the Bush River is assumed to be discharged from a 295-port diffuser having a total length of 448 m and a "port" spacing of 1.5 m (5 ft). As noted by Roberts et al. (1989), the dilution level of a buoyancy-dominated plume is nearly independent of the number of ports, and depends primarily on *plume length* and buoyancy. The schematized diameter of each port is 1.2 m (4 ft). Such large diameter ports allowed momentum-related effects to be minimized. In addition, further testing showed that Model results are insensitive to port diameter (see section 2.7).

2.6 Near-Field Model Results

Using information described above, an input file was constructed for the ULINE model and a series of simulation were conducted. *The results indicate that a near-field dilution ratio of 159:1 is appropriate for application of acute toxicity criteria. Similarly, model results indicate that a near-field dilution ratio of 1,006:1 is appropriate for application of chronic toxicity criteria.* Having limited buoyancy and initial

momentum, this near-field dilution is effected largely by the large discharge length of 427 m (1,400 ft) and ambient current field.


2.7 Model Sensitivity Analysis

The accuracy of any model results depends, in part, upon the accuracy of selected input parameter values. Accordingly, a model sensitivity analysis was performed to evaluate effects of individual parameter variations on model output. Results of this analysis are plotted in Figures 2-6 and described below.

Figure 2 displays model sensitivity to variations in the groundwater discharge to the Bush River. As noted above, the selected discharge rate of $0.0136 \text{ m}^3/\text{sec}$ (0.48 cfs) corresponds to a dilution ratio of 159. However, higher discharge rates are likely to decrease the near-field dilution (Figure 2). For example, a near doubling of the discharge rate to $0.025 \text{ m}^3/\text{sec}$ nearly halves the dilution to a ratio of only 88.

Likewise, Figure 3 represents the model sensitivity to the selected discharge length (i.e., the assumed length of the "line source"). As noted above, the discharge length is estimated to be 448 m (1,470 ft). This value assumes the extent of the lateral discharge to be the entire length of the Beach Point Peninsula fronting Bush River. However, a more conservative estimate of approximately half this length (i.e., 224 m, 735 ft) corresponds to a simulated near-field dilution ratio of 81. Also, a discharge of length of (152 m) 500 ft corresponds to a dilution ratio of 54. Overall, the dilution is somewhat sensitive to, and approximately linearly related to, this assumed discharge length.


SIMULATED DILUTION vs. DISCHARGE



A51-21

FIGURE 2

DILUTION vs. DISCHARGE LENGTH




The model results are also somewhat dependent upon the assumed discharge depth. Figure 4 shows a nearly linear relation between the simulated dilution ratio and discharge depth. In this figure, note that the average water depth in the near field is approximately 1.5 m (5 ft). However, a conservative value of half this depth (i.e., 0.75 m or 2.5 ft) is used in all other simulations. This conservative assumption results in a dilution of 159 — nearly half of the corresponding dilution for a 1.5-ft-deep discharge (i.e., 319).

Typically, near-field models are sensitive to the ambient current speed. Figure 5 suggests that the adapted model is no exception. For example, the estimated tenth-percentile speed (i.e., 0.03 m/sec) corresponds to a dilution ratio of 159, while a speed of 0.02 m/sec corresponds to a dilution of 106. Thus, it is unlikely that uncertainties in the speed estimate will greatly enhance the computed dilution levels.


Finally, the adapted model assumes a Bush River salinity of 3 ppt. Figure 6 suggests that the model is relatively insensitive to other representative values for this parameter.

Based on the overall results of the model sensitivity analysis, a more conservative estimate of the near-field dilution factor would lie within the range of 54 to 159 for application of acute toxicity criteria. Similarly, a conservative estimate of the near-field dilution factor would lie within the range of 499 to 1,006 for application of chronic toxicity criteria.

DILUTION vs. DISCHARGE DEPTH




SIMULATED DILUTION vs. AMBIENT CURRENT



A51-25

FIGURE 5

DILUTION vs. AMBIENT SALINITY



3. FAR-FIELD DILUTION ESTIMATES

3.1 Definitions

The Bush River receiving waters can be divided into two zones based on the physical mechanisms of mixing. In the near-field, rapid mixing is caused primarily by the discharge plume's buoyancy and initial discharge momentum, in combination with ambient currents. As noted above, near-field length scales are typically ones to tens of meters; near-field time scales are typically seconds to tens of minutes.

In the far-field, the plume is mixed passively by ambient processes such as turbulence and tidal dispersion. Far-field length scales are typically kilometer distances from the discharge site; far-field time scales are hours, days or more.

The following sections provide preliminary assessments of the far-field dilution (i.e., the flushing rate) in the Bush River Estuary. Methodologies for such techniques are documented in "Water Quality Assessment: a Screening Procedure for Toxic and Conventional Pollutants" (EPA/600/6-85/002b). Input required for these analyses includes the local tidal range, mean freshwater inflow and salinity data.

3.2 Tidal Prism Method

The tidal prism method was first introduced by Ketchum (1951) and later elaborated (EPA, 1985). This method divides an estuary into a number of hypothetical segments having lengths of approximately one tidal excursion. Within each segment, complete mixing is assumed. Thus, the flushing flow for the i^{th} segment, Q_{fi} , is given by the tidal prism volume for that segment, P_i , divided by the dominant tidal period, T (i.e., 1 tidal cycle = 12.42 hours or 44,712 sec). The

corresponding residence time for that segment is computed as the total segment volume (i.e., $P_i + V_i$, where V_i is the MLW volume) divided by the segment flushing

rate. The total segment residence time is the sum of all individual segment residence times.

Application of this method to the Bush River Estuary is illustrated in Table 3. In this application, the Bush River is divided into 6 hypothetical segments as delineated in the previous one-dimensional modeling study by Carter (1976). Typical segments are approximately 2-3 km long, comparable to one tidal excursion. Table 3 lists corresponding intertidal volumes and MLW (i.e., "subtidal") volumes for each segment. Table 3 also lists the computed segment "exchange ratio," r_i , defined as the tidal prism, P_i , divided by the total segment volume, $P_i + V_i$. Thus, the computed residence time for the i^{th} segment (in units of tidal cycles) is simply the inverse of r_i . The tidal prism method calculations indicate that the total residence time for the Bush River is approximately 31 days. Moreover, the flushing time below the Beach Point site (i.e., for segments 2 and 1) is approximately 13 days (24.3 cycles). This suggests that the Bush River has limited "memory" of past discharges from the site. It should be noted, however, that the tidal prism method may underestimate the flushing time of an estuary since the selected segments are not, in reality, completely mixed.

Based on Table 3, the theoretical flushing rate at the Beach Point site, Q_{f2} , is given by the ratio P_2/T or approximately $1.0 \times 10^6 \text{ m}^3$ per tidal cycle i.e., per 44,712 seconds). Thus, the corresponding far-field dilution is obtained by dividing this flushing rate by the assumed groundwater discharge flow, $0.0136 \text{ m}^3/\text{sec}$ (see Eq. 6).

TABLE 3: CALCULATION TABLE FOR TIDAL PRISM METHOD

| Segment Number | Segment Dimensions | | | | Subtidal Water Volume, V_i (m^3) | Intertidal Water Volume $P_{i,2}$ (m^3) | Segment Exchange Ratio r_i | Segment Flushing Time, T_i (Tidal Cycles) |
|----------------|-----------------------------------|---------------------------------|------------------------------------|--------------------|--|---|------------------------------|---|
| | Starting Distance Above Mouth (m) | Ending Distance Above Mouth (m) | Distance of Center Above Mouth (m) | Segment Length (m) | | | | |
| 6 | 18,810 | 16,110 | 17,460 | 2,700 | 2.0×10^6 | 4.3×10^5 | 0.18 | 5.6 |
| 5 | 15,310 | 11,910 | 13,610 | 3,400 | 6.1×10^6 | 8.7×10^5 | 0.12 | 8.3 |
| 4 | 11,910 | 9,260 | 10,585 | 2,650 | 8.0×10^6 | 8.5×10^5 | 0.10 | 10.0 |
| 3 | 9,260 | 6,560 | 7,910 | 2,700 | 1.1×10^7 | 9.5×10^5 | 0.08 | 12.5 |
| 2 | 6,560 | 3,700 | 5,130 | 2,860 | 1.3×10^7 | 1.0×10^6 | 0.07 | 14.3 |
| 1 | 3,700 | 0 | 1,850 | 3,700 | 1.6×10^7 | 1.7×10^6 | 0.10 | 10.0 |
| | | | | | | | | |
| | | | | | | | | |
| | | | | | | | | |
| | | | | | | | | |
| | | | | | | | | |
| | | | | | | | | |
| | | | | | | | | |
| | | | | | | | | |
| | | | | | | | | |
| | | | | | | | | |
| | | | | | | | | |
| | | | | | | | $\sum_{i=1}^n T_i =$ | 60.7 cycles (31.4 days) |

This yields an estimated far-field dilution of 1,618. Again, note that this method likely overestimates this dilution since it assumes complete mixing within segments.

3.3 Fraction of Freshwater Method

This alternate method for estimating estuarine flushing times uses the freshwater content of estuarine segments as a tracer. That is, the "fraction of freshwater" for the i^{th} segment, f_i , is defined as the difference between the Bush River entrance salinity and the i^{th} segment salinity, $S_s - S_i$, divided by the entrance salinity, S_s (see Table 4, sixth column). The corresponding "freshwater volume," W_i , for that segment is approximated by the fraction of freshwater, f_i , multiplied by the mean segment volume, V_i . Thus, in the steady state, the segment flushing time is simply the segment freshwater volume divided by the freshwater inflow, R . Again, the total residence time is obtained by summing the individual segment flushing times.

Application of this method to the Bush River Estuary is illustrated in Table 4. Note that the segmentation used for this method is the same as that employed in the previous section. The fraction-of-freshwater-method calculations estimate the total residence time for the Bush River as 57 days — somewhat longer than the tidal prism method estimate. However, the computed flushing time below the Beach Point site (i.e., for segments 2 and 1) is nearly the same — approximately 14 days (26.7 cycles). This also suggests that the Bush River has limited "memory" of past discharges from the site.

3.4 Previous Dye-Tracer Study Results

Far-field dilution also may be computed empirically based on field experiments. Accordingly, the dye-tracer data collected by Carter (1976) in the Bush River allows

→ Up Estuary

A51-31

for such computations. Under spring (high-flow) conditions, Carter reports a steady dye dilution ratio of 267 (i.e., 61.4 ppb effluent concentration to 0.23 ppb measured).

Likewise, under fall (low-flow) conditions, Carter reports a dilution ratio of 315 (i.e., 50.5 ppb initial concentration to 0.16 ppb measured). Carter's data provides a more conservative estimate of the far-field dilution than the segmented (complete mixing) approaches described in sections 3.2 and 3.3. For this reason, and because it is based on direct measurements, the above-mentioned dilution ratios of 267 and 315 are adopted for use in the calculation of total near-field dilution.

3.5 Calculation of Total Near-Field Dilution

For a conservative constituent, the far-field, near-field and total near-field dilutions (S_f , S_n and S_t , respectively) are defined (e.g., Fischer, 1979; Adams, 1990) as:

$$S_f = \frac{C_o - C_a}{C_f - C_a} \quad (1)$$

$$S_n = \frac{C_o - C_f}{C_n - C_f} \quad (2)$$

$$S_t = \frac{C_o - C_a}{C_n - C_a} \quad (3)$$

where C_o , C_n and C_f are the constituent concentrations at the discharge, at the end of the near-field, and in the far-field, respectively. Note that in these definitions, the far-field concentration serves as a background concentration for the near-field in the same way that the ambient concentration serves as a background for the total near-field. These definitions may be combined to yield (Adams, 1990):

$$S_t = \frac{1}{\frac{1}{S_n} + \frac{1}{S_f} - \frac{1}{S_n S_f}} \quad (4)$$

or

$$\frac{1}{S_t} \sim \frac{1}{S_n} + \frac{1}{S_f} \quad (5)$$

This inverse sum indicates that the total dilution in the near field is always less than both the near-field and far-field dilutions. Moreover, the effectiveness of near-field mixing is always limited by the magnitude of the far-field mixing.

In the absence of other pollutant sources (i.e., $C_a=0$ in Eq. 1), the far-field dilution reduces to a ratio of either concentrations or flows (Adams, 1990):

$$S_f = \frac{C_o}{C_f} = \frac{Q_f}{Q_o} \quad (6)$$

where: C_o is the discharge concentration; C_f is the far-field concentration; Q_o is the discharge flow; and Q_f is the far-field "flushing flow" or downestuary transport rate (Officer, 1976). Note that the flushing flow is not a measurable flow rate, but is the equivalent rate at which mass is transported seaward.

Based on an application of Eq. 5 with a value of $S_n=50$, the total near-field dilutions for acute toxicity conditions is calculated to be approximately 42 and 43 for under spring and fall conditions, respectively. Similarly, an application of Eq. 5 with a value of $S_n=450$, yield total near-field dilutions for chronic toxicity conditions of approximately 168 and 185 for under spring and fall conditions, respectively.

3.6 Application of Dilution Factors to Toxics Discharge Data

Using the above calculated dilution factors, near-field concentrations can be estimated for the parameters of interest. This estimate would be equal to the mean groundwater

concentration divided by the dilution factor. Thus, using the above dilution factor, a pollutant concentration of 42 ppb in the groundwater would be diluted to 1 ppb within

the near-field limits of the Bush River. In this manner, near-field concentrations for all detected groundwater constituents are calculated. These results concentrations are shown in Tables 5 and 6. An examination of these Table reveals that the expected near-field impacts will be near or below the method detection limit for most parameters. For such constituents, these analyses suggest that on-site contamination will not produce any discernable effect on the water quality of the Bush River.

Table 5: Computed Dilutions of Volatile Organics (ug/l)

| Constituent | Groundwater Concentration | <u>Receiving-Water Concentration</u> | |
|---------------------------|------------------------------|--------------------------------------|-----------|
| | | Acute* | Chronic** |
| Chlorobenzene | 2 | 0.05 | 0.01 |
| Chloroform | 12 | 0.29 | 0.07 |
| 1,1-Dichloroethene | 5 | 0.12 | 0.03 |
| cis-1,2-Dichloroethene | 640 | 15.24 | 3.81 |
| 1,1,1,2-Tetrachloroethane | 3 | 0.07 | 0.02 |
| 1,1,2,2-Tetrachloroethane | 17,000 | 404.76 | 101.19 |
| Tetrachloroethene | 90 | 2.14 | 0.54 |
| 1,1,2-Trichloroethane | 110 | 2.62 | 0.65 |
| Trichloroethene | 1,600 | 38.10 | 9.52 |
| Vinyl Chloride | 2 | 0.05 | 0.01 |

* Based on observed average concentration and dilution ratio of 42:1

** Based on observed average concentration and dilution ratio of 168:1

Table 6: Computed Dilutions of Heavy Metals (ug/l)

| Constituent | Groundwater Concentration | <u>Receiving-Water Concentration</u> | |
|-------------|------------------------------|--------------------------------------|-----------|
| | | Acute* | Chronic** |
| Aluminum | 1,500 | 35.71 | 8.93 |
| Antimony | 5 | 0.12 | 0.03 |
| Arsenic | 7 | 0.17 | 0.04 |
| Beryllium | 5 | 0.12 | 0.03 |
| Boron | 70 | 1.67 | 0.42 |
| Cadmium | 1.4 | 0.03 | 0.01 |
| Chromium | 5 | 0.12 | 0.03 |
| Cobalt | 110 | 2.62 | 0.65 |
| Copper | 8 | 0.19 | 0.05 |
| Lead | 5 | 0.12 | 0.03 |
| Mercury | 0.5 | 0.01 | 0.003 |
| Molybdenum | 40 | 0.95 | 0.24 |
| Nickel | 180 | 4.29 | 1.07 |
| Selenium | 5 | 0.12 | 0.03 |
| Silver | 1 | 0.02 | 0.006 |
| Thallium | 5 | 0.12 | 0.03 |
| Tin | 100 | 2.38 | 0.60 |
| Zinc | 310 | 7.38 | 1.85 |

* Based on observed average concentration and dilution ratio of 42:1

** Based on observed average concentration and dilution ratio of 168:1

4. MIXING ZONE CONSIDERATIONS

The concept of mixing zones is normally applied in the case of wastewater facilities and power plants that discharge to tidal/coastal waterways via an outfall diffuser manifold. The following sections summarize Maryland's mixing zone regulations and explore their applicability to the subject groundwater discharges.

4.1 Mixing Zone Regulations

A mixing zone is defined (COMAR 26.08.01.01B) as "an area contiguous to a discharge where surface water quality or groundwater quality does not have to meet: (a) all water quality criteria; or (b) all requirements otherwise applicable to the natural water." That is, pollutant concentrations within the mixing zone may exceed the specified water quality standards within a localized vicinity of the outfall. The dimensions of this zone are normally comparable to those associated with the initial dilution processes (e.g., 10-100 meters). Thus, marine organisms would only be exposed to concentrations exceeding the specified criteria for a brief period during a transit of the mixing zone (Muellenhoff et al., 1985). Of course, water quality criteria must still be met outside this zone. Moreover, as stated in COMAR 26.08.02.05A, mixing zones may be allowed "provided that the following requirements are met outside the mixing zone: (a) there shall be no interference with biological communities or populations of indigenous species to a degree which is damaging to the aquatic life or ecosystem; (b) there shall be no diminishing of other legitimate beneficial uses; (c) mixing zones may not form barriers to the migratory routes of aquatic life; (d) mixing zones shall be designated and located to protect surface waters and shallow water shoreline areas; (e) the general water

quality criteria set out in regulation .03B(1)-(3) of this chapter apply within this mixing zone."

Four types of numerical toxic substance criteria are promulgated by the MDE: (a) human health criteria through ingestion of public water supplies; (b) the wholesomeness of fish for human consumption; (c) fresh, estuarine, and salt water aquatic life criteria from *acute* toxicity impacts; and (d) fresh, estuarine, and salt water aquatic life criteria from *chronic* toxicity impacts. For the purpose of applying numerical toxic substance criteria, the Bush River Area (Sub-Basin 02-13-07) is classified by the MDE as "fresh water" (Comar 26.08.02.03-1B) and as "Use II" (shellfish harvesting waters). Corresponding toxic substances criteria for both ambient surface waters ("fresh water") and human health ("fish consumption") are listed in Table 7. Note that toxic criteria related to drinking water do not apply within shellfish harvesting waters.

The following regulations (COMAR 26.08.02.05D) pertain to the application of toxic substance *Chronic Criteria* for the protection of Aquatic Life. First, in terms of size, the Regulatory Mixing Zone (i.e., the mixing zone in which the Chronic Criteria are applied) may not exceed 10 percent of the cross-sectional area of the receiving waters. Also, *the Chronic Criteria are applied under flow conditions determined from site-specific data for the mean tide level, average tidal velocity, and, when appropriate, the design streamflow.* Based on the MDE's discretion, a plume-dilution study also may be required for the application of these criteria. In accordance with COMAR 26.08.02.06(a), conditions pertinent to the application of toxic substance *Chronic Criteria* are also relevant to the application of toxic substance *Human Health Criteria*.

Table 7: Applicable Toxic Substances Criteria for the Bush River (ppb)

| Constituent | <u>Aquatic Life</u> | | <u>Human Health</u> |
|-----------------------|---------------------|---------|---------------------|
| | Acute | Chronic | Fish Consumption |
| Arsenic (tri) | 360 | 190 | |
| Cadmium | 3.9 | 1.1 | |
| Chromium (hex) | 16 | 11 | |
| Chromium (tri) | 1700 | 210 | 3,433,000 |
| Copper | 18 | 12 | |
| Cyanide | 22 | 5.2 | |
| Lead | 82 | 3.2 | |
| Mercury | 2.4 | 0.012 | 0.146 |
| Nickel | 1,400 | 160 | 100 |
| Selenium | 20 | 5 | |
| Silver | 4.1 | 0.12 | |
| Zinc | 120 | 110 | |
| Aldrin | 3 | | 0.00079 |
| Benzidine | | | 0.0053 |
| 1,1DCE | | | 18.5 |
| DDT | 1.1 | 0.001 | 0.00024 |
| Dieldrin | 2.5 | 0.0019 | 0.00076 |
| Endrin | 0.18 | 0.0023 | |
| Lindane | 2 | 0.08 | 0.625 |
| PCP | 20 | 13 | |
| PCB | 2 | 0.014 | 0.00079 |
| 1,1,1TCA | | | 1,030,000 |
| Toxaphene | 0.73 | 0.0002 | 0.0073 |
| TCE | | | 807 |
| Tributyltin (TBT) | | 0.026 | |
| 2,3,7,8 TCDD (Dioxin) | | | 0.0000012 |

The following regulations (COMAR 26.08.02.05C) pertain to the application of toxic substance *Acute Criteria* for the protection of Aquatic Life. Such acute criteria (for low-velocity dischargers) must be met within a short distance from the outfall using the most restrictive of the following conditions:

- (a) Within 10 percent of the distance (in any spatial direction) from the discharge to the edge of the Regulatory Mixing Zone (RMZ) used for the application of the chronic criteria.
- (b) Within a distance of 50 times the "discharge length scale" (i.e., 50 times the square root of the cross-sectional area of the discharge outlet).
- (c) Within a distance of 5 times the local water depth in any horizontal direction from the discharge outlet.

Thus, the region where the acute criteria are applied, sometimes called the "Toxic Dilution Zone" (TDZ), extends outward from the discharge to a distance no larger than any of the three lengths cited above. Moreover, the TDZ must not occupy more than 5 percent of the cross-sectional area of the receiving waters.

Also, the acute criteria are applied under flow conditions determined from site-specific data for the mean low water elevation, the minimum daily averaged 1-hour tidal velocity and, when appropriate, design streamflow conditions. Such tidal stages are deemed "critical" (minimum-dilution) conditions.

4.2 Site Applicability

The regulatory conditions relevant to mixing zones were developed for outfall diffusers. Their application to an area-wide groundwater discharge is somewhat uncertain due to the unknown extent of this area. It seems likely that the discharge area is confined within the the RMZ, which for this case extends approximately 500 feet from the river

bank. However, the extent of the TDZ is more poorly defined as it partially depends on the configuration of the discharge. Thus, the applicability of the mixing zone concept must be further examined.

The concept of mixing zones provides an intermediate approach to the assessment of toxic constituent impacts. The two other approaches generally in use are "end of pipe" and "fully mixed". The "end of pipe" approach allows for zero dilution -- criteria are directly applied to the discharge water quality. This simplified approach yields the most stringent results which, in turn, most ensures the protection of aquatic biota. However, it may lack a physical or biological basis in terms of receiving water impacts. Thus, its results are not considered to be defensible -- it is effectively a non-technically based policy decision. Conversely, the "fully mixed" (or far-field) approach considers the total volume of the receiving water segment to be instantly available for the dilution of a discharge. This approach is generally far less stringent than the "end of pipe" approach. However, it may not adequately address biological impacts in the receiving water due to near-field conditions. The concept of mixing zones was evolved to address these limitations. Mixing zones allow consideration of the initial dilution process. This approach also protects aquatic life by limiting the potential for exposure to elevated concentrations to a minor portion of the waterway. As only a portion of the available dilution is considered in this approach, mixing zones provide a much more stringent basis for regulation than the "fully mixed" approach.

In the application of the mixing zone concept to the Bush River, it was necessary to conceptualize the area-wide groundwater discharge from the APG as a line source (diffuser). Due to this approach, potential impacts could be assessed in a technically defensible manner. However, this approach had a side effect of concentrating the

groundwater impact into a smaller area. *That is, this approach minimized the available near-field dilution and projected higher receiving water concentrations than would actually occur. Thus, the mixing zone analysis produced an overly conservative result. In the near-field, actual receiving water concentrations would be less than those projected due to more dispersed nature of the groundwater plume.* In the far-field, projected and actual receiving water concentrations should be comparable. However far-field concentrations would always be less than the near-field projections.

4.3 Conformance with Criteria

In Tables 8 and 9, the anticipated near-field receiving water concentrations are compared with the applicable water quality criteria. As shown in these Tables, no exceedences of these criteria are projected.

There will be a detectable impact on the receiving water concentrations due to volatile organic constituents (such as cis-1,2 dichloroethene; 1,1,2,2 tetrachloroethane; and trichloroethene). However, with the exception of trichloroethene (TCE), Maryland has not implemented criteria for these constituents.

**Table 8: Comparison of Projected Near-Field Concentration with
Applicable Toxic Substances Acute Criteria (ppb)**

| Constituent | Near-Field Concentration | Aquatic Life Acute Criteria |
|----------------|-----------------------------|--------------------------------|
| Arsenic (tri) | 0.17 (total) | 360 |
| Cadmium | 0.03 | 3.9 |
| Chromium (hex) | 0.12 (total) | 16 |
| Chromium (tri) | | 1700 |
| Copper | 0.19 | 18 |
| Cyanide | | 22 |
| Lead | 0.12 | 82 |
| Mercury | 0.01 | 2.4 |
| Nickel | 4.29 | 1,400 |
| Selenium | 0.12 | 20 |
| Silver | 0.02 | 4.1 |
| Zinc | 7.38 | 120 |
| DDT | | 1.1 |
| Dieldrin | | 2.5 |
| Endrin | | 0.18 |
| Lindane | | 2 |
| PCP | | 20 |
| PCB | | 2 |
| Toxaphene | | 0.73 |

**Table 8: Comparison of Projected Near-Field Concentration with
Applicable Toxic Substances Chronic and Human-Health Criteria (ppb)**

| Constituent | Near-Field Concentration | Aquatic Life Chronic Criteria | Human-Health Fish Consump. |
|-----------------------|-----------------------------|----------------------------------|-------------------------------|
| Arsenic (tri) | 0.04 (total) | 190 | |
| Cadmium | 0.01 | 1.1 | |
| Chromium (hex) | 0.03 (total) | 11 | |
| Chromium (tri) | | 210 | 3,433,000 |
| Copper | 0.05 | 12 | |
| Cyanide | | 5.2 | |
| Lead | 0.03 | 3.2 | |
| Mercury | 0.003 | 0.012 | 0.146 |
| Nickel | 1.07 | 160 | 100 |
| Selenium | 0.03 | 5 | |
| Silver | 0.006 | 0.12 | |
| Zinc | 1.85 | 110 | |
| | | | |
| Aldrin | | | 0.00079 |
| Benzidine | | | 0.0053 |
| 1,1DCE | | | 18.5 |
| DDT | | 0.001 | 0.00024 |
| Dieldrin | | 0.0019 | 0.00076 |
| Endrin | | 0.0023 | |
| Lindane | | 0.08 | 0.625 |
| PCP | | 13 | |
| PCB | | 0.014 | 0.00079 |
| 1,1,1TCA | | | 1,030,000 |
| Toxaphene | | 0.0002 | 0.0073 |
| TCE | 9.52 | | 807 |
| Tributyltin (TBT) | | 0.026 | |
| 2,3,7,8 TCDD (Dioxin) | | | 0.0000012 |

5. SUMMARY AND CONCLUSIONS

This study was initiated to assess the potential impact to the Bush River due to

contaminated groundwater emanating from the APG site in Harford County, Maryland. Towards this objective, an EPA-approved near-field model was adapted to the receiving-waters adjacent to the APG site. The model approach and input was developed to provide a conservative assessment of receiving water impacts. This analysis suggests a near-field dilution level of approximately 42:1 for the application of acute criteria and a near-field dilution level of approximately 168:1 for the application of chronic criteria. Thus, contaminants introduced into Bush River receiving-waters at a concentration of 42 ppb will be diluted locally to a concentration of approximately 1 ppb or less. By applying these dilution factors to the groundwater quality data from the APG site, near-field concentrations within the Bush River can be projected. As shown in Tables 8 and 9, none of the projected receiving water concentrations exceed Maryland's current acute or chronic criteria.

There will be a detectable impact on the receiving water concentrations of certain volatile organic constituents (such as cis-1,2 dichloroethene; 1,1,2,2 tetrachloroethane; and trichloroethene). Should Maryland enact more stringent criteria for these parameters, these results and conclusions must be re-evaluated. In such a case, a field study would be recommended. Such a study would include in-situ current measurement, dye release and tracking, CTD casts, and/or water quality data collection.

REFERENCES

- Adams, E.E., 1990. "Analysis of initial mixing in the Hackensack River near the outfall for the Bergen County Utility Authority wastewater Treatment Facility." Report prepared for Clinton Bogert Associates, Englewood Cliffs, New Jersey, 88 pp.
- Adams, E.E., Stalzenbach, K.D., and Harleman, D.R.F., 1975, "Near and Far Field Analysis of Thermal Discharges into Large Bodies Of Water," Technical Report No. 205, R.M. Parsons Laboratory for Water Resources and Hydrodynamics, MIT.
- Alam, A.M., Harleman, D.R.F., and Colonell, J.M., 1982. "Evaluation of selected initial dilution models," *ASCE, Jour. Envir. Engnr. Div.*, 108:159-186.
- Carter, H.H., 1976. "Simple one dimensional kinematic model results for the Bush River and Romney Creek." Chesapeake Bay Institute, Special Report 49, Reference 76-2, 37 pp.
- Eskin, R.A., Rowland, K.H., Alegre, D.Y., and Magnien, R.E., 1994. "Contaminants in Chesapeake Bay sediments (1984-1991)." Chapter 4 of Draft Report, Maryland Department of the Environment, June 1994, 164 pp.
- Fischer, H.B., List, E.J., Koh, R.C.Y., Imberger, N.H., and Brooks, N.H., 1979, Mixing in Inland and Coastal Waters, Academic Press.
- K-V Associates, Inc., 1994. "Hydrogeologic study of Beach Point, Edgewood Area Aberdeen Proving Grounds, Maryland." Report prepared for Jacobs Engineering Group, Bel Air Maryland, 15 pp.
- K-V Associates, Inc., 1994. "Estimated discharge rates in the surficial aquifer, Beach Point (Peninsula) study area, Aberdeen Proving Ground - Edgewood Area, Maryland." Letter report calculations, 1 p.
- MacDonald, 1993. "Development of an approach to the assessment of sediment quality in Florida Coastal Waters." Report, Florida Department of Environmental Regulations.
- Martin Marietta Environmental Systems, 1986. "Prediction of the effects of nutrient loadings from a power plant at Perryman, MD on the water quality


of the Bush River Estuary." Report prepared for the Maryland Department of Natural Resources, Power Plant Research Program, Annapolis Maryland.

- Muellenhoff, W.P., et al, 1985, "Initial Mixing Characteristics of Municipal Ocean Discharges," Pacific Division, Environmental Research Laboratory, U.S. EPA, Newport, Oregon.
- Najarian & Associates, Inc., 1990. "A Field and Model Study of the Dilution Capacity of the Passaic Valley Sewerage Commissioner's Wastewater Outfall System." Technical Report prepared for Passaic Valley Sewerage Commissioners, Newark, New Jersey.
- Officer, C.B., 1976. Physical Oceanography of Estuaries and Associated Coastal Waters, John Wiley and Sons.
- Pritchard, D.W., 1976. "Distribution of contaminants and excess heat in the Bush River and Romney Creek from the proposed Perryman Electric Power Plant." Chesapeake Bay Institute, Special Report 58, Reference 76-13, 30 pp.
- Roberts, P.J.W., Snyder, W.H., and Baumgartner, D.J., 1989. "Ocean outfalls. I: Submerged wastefield formation." *ASCE, Jour. Envir. Engnr. Div.*, 115:1-25.
- Roberts, P.J.W., Snyder, W.H., and Baumgartner, D.J., 1989. "Ocean outfalls. II: Spatial Evolution of Submerged Wastefield." *ASCE, Jour. Envir. Engnr. Div.*, 115:26-48.
- Roberts, P.J.W., Snyder, W.H., and Baumgartner, D.J., 1989. "Ocean outfalls. III: Effect of diffuser design on submerged wastefield." *ASCE, Jour. Envir. Engnr. Div.*, 115:49-71.
- Tetra Tech, Inc, 1982, "Design of 301(h) Monitoring programs for Municipal Wastewater Discharges to Marine Waters," EPA 430/9-82-010.
- Tetra Tech, Inc, 1982, "Revised Section 301(h) Technical Support Document", EPA 430/9-82-011.
- USEPA, 1985, "Technical Support Document for Water Quality-based Toxics Control," EPA-440/4-85-032.

USEPA, 1990, "Technical Support Document for Water Quality-based Toxics Control," EPA-/505/2-90-001.


APPENDIX A

| TIME (hours) | AMPL. (m) | |
|-----------------|--------------|----------|
| | 0 | 0 |
| 0.125 | 0.019274205 | 0.019274 |
| 0.25 | 0.038471361 | 0.038471 |
| 0.375 | 0.057514728 | 0.057515 |
| 0.5 | 0.07632818 | 0.076328 |
| 0.625 | 0.094836512 | 0.094837 |
| 0.75 | 0.112965735 | 0.112966 |
| 0.875 | 0.130643379 | 0.130643 |
| 1 | 0.147798777 | 0.147799 |
| 1.125 | 0.164363351 | 0.164363 |
| 1.25 | 0.180270883 | 0.180271 |
| 1.375 | 0.195457785 | 0.195458 |
| 1.5 | 0.209863346 | 0.209863 |
| 1.625 | 0.22342998 | 0.22343 |
| 1.75 | 0.236103455 | 0.236103 |
| 1.875 | 0.247833108 | 0.247833 |
| 2 | 0.25857205 | 0.258572 |
| 2.125 | 0.268277354 | 0.268277 |
| 2.25 | 0.276910221 | 0.27691 |
| 2.375 | 0.284436141 | 0.284436 |
| 2.5 | 0.290825031 | 0.290825 |
| 2.625 | 0.296051351 | 0.296051 |
| 2.75 | 0.300094208 | 0.300094 |
| 2.875 | 0.302937441 | 0.302937 |
| 3 | 0.304569684 | 0.30457 |
| 3.125 | 0.304984413 | 0.304984 |
| 3.25 | 0.304179969 | 0.30418 |
| 3.375 | 0.302159569 | 0.30216 |
| 3.5 | 0.298931289 | 0.298931 |
| 3.625 | 0.294508033 | 0.294508 |
| 3.75 | 0.288907484 | 0.288907 |
| 3.875 | 0.282152031 | 0.282152 |
| 4 | 0.274268677 | 0.274269 |
| 4.125 | 0.265288936 | 0.265289 |
| 4.25 | 0.255248705 | 0.255249 |
| 4.375 | 0.24418812 | 0.244188 |
| 4.5 | 0.232151395 | 0.232151 |
| 4.625 | 0.219186648 | 0.219187 |
| 4.75 | 0.205345703 | 0.205346 |
| 4.875 | 0.190683891 | 0.190684 |
| 5 | 0.175259821 | 0.17526 |
| 5.125 | 0.159135153 | 0.159135 |
| 5.25 | 0.142374342 | 0.142374 |
| 5.375 | 0.125044392 | 0.125044 |
| 5.5 | 0.107214578 | 0.107215 |
| 5.625 | 0.088956175 | 0.088956 |
| 5.75 | 0.07034217 | 0.070342 |
| 5.875 | 0.051446972 | 0.051447 |
| 6 | 0.032346116 | 0.032346 |



| | | | | | | |
|----------------|--------------|----------|----------|--|--|--|
| 6.125 | 0.013115957 | 0.013116 | | | | |
| 6.25 | -0.006166633 | 0.006167 | 0.043252 | | | |
| 6.375 | -0.025424572 | 0.025425 | | | | |
| 6.5 | -0.044580877 | 0.044581 | | | | |
| 6.625 | -0.06355897 | 0.063559 | | | | |
| 6.75 | -0.082282986 | 0.082283 | | | | |
| 6.875 | -0.100678077 | 0.100678 | | | | |
| 7 | -0.118678700 | 0.118671 | | | | |
| 7.125 | -0.136188955 | 0.136189 | | | | |
| 7.25 | -0.153162787 | 0.153163 | | | | |
| 7.375 | -0.169524352 | 0.169524 | | | | |
| 7.5 | -0.185208246 | 0.185208 | | | | |
| 7.625 | -0.200151771 | 0.200152 | | | | |
| 7.75 | -0.214295191 | 0.214295 | | | | |
| 7.875 | -0.227581968 | 0.227582 | | | | |
| 8 | -0.239958988 | 0.239959 | | | | |
| 8.125 | -0.251376774 | 0.251377 | | | | |
| 8.25 | -0.261789684 | 0.26179 | | | | |
| 8.375 | -0.271156092 | 0.271156 | | | | |
| 8.5 | -0.279438556 | 0.279439 | | | | |
| 8.625 | -0.286603967 | 0.286604 | | | | |
| 8.75 | -0.292623681 | 0.292624 | | | | |
| 8.875 | -0.297473635 | 0.297474 | | | | |
| 9 | -0.30113444 | 0.301134 | | | | |
| 9.125 | -0.303591463 | 0.303591 | | | | |
| 9.25 | -0.304834882 | 0.304835 | | | | |
| 9.375 | -0.304859727 | 0.30486 | | | | |
| 9.5 | -0.303665897 | 0.303666 | | | | |
| 9.625 | -0.301258166 | 0.301258 | | | | |
| 9.75 | -0.297646158 | 0.297646 | | | | |
| 9.875 | -0.292844312 | 0.292844 | | | | |
| 10 | -0.286871824 | 0.286872 | | | | |
| 10.125 | -0.279752568 | 0.279753 | | | | |
| 10.25 | -0.271515004 | 0.271515 | | | | |
| 10.375 | -0.262192061 | 0.262192 | | | | |
| 10.5 | -0.251821007 | 0.251821 | | | | |
| 10.625 | -0.240443302 | 0.240443 | | | | |
| 10.75 | -0.228104426 | 0.228104 | | | | |
| 10.875 | -0.214853706 | 0.214854 | | | | |
| 11 | -0.200744109 | 0.200744 | | | | |
| 11.125 | -0.18583204 | 0.185832 | | | | |
| 11.25 | -0.170177108 | 0.170177 | | | | |
| 11.375 | -0.153841896 | 0.153842 | | | | |
| 11.5 | -0.136891701 | 0.136892 | | | | |
| 11.625 | -0.119394284 | 0.119394 | | | | |
| 11.75 | -0.101419589 | 0.10142 | | | | |
| 11.875 | -0.083039469 | 0.083039 | | | | |
| 12 | -0.064327401 | 0.064327 | | | | |
| 12.125 | -0.045358184 | 0.045358 | | | | |
| 12.25 | -0.026207648 | 0.026208 | | | | |
| 12.375 | -0.006952348 | 0.006952 | | | | |
| AVERAGE SPEED= | | 0.192927 | | | | |

TIDAL VELOCITIES



Estimated Discharge Rates in the Surficial Aquifer Beach Point (Peninsula) Study Area

Aberdeen Proving Ground-Edgewood Area, Maryland

Assumptions: V_{hor} measured in MW-33B.1 is continuous throughout the Surficial Aquifer at the Beach Point peninsula; V_{hor} orientation (flow direction) is constant with actual aquifer conditions over one complete tidal cycle (discounting short-term fluctuations due to tidal changes); the Surficial Aquifer at Beach Point is homogeneous, characterized by isotropic flow conditions.

MW-33B.1 (screened 41'-46'): $V_{hor} = 0.89$ ft/day (61.8 NE)
Average saturated thickness in Surficial Aquifer = 61.8'
Length of Beach Point peninsula fronting Bush River = 1,470'
Area of saturated Surficial Aquifer fronting Bush River = 90,882.7 ft²

1. If we assume V_{hor} is oriented perpendicular (orthogonal) to the Bush River frontage in the Surficial Aquifer, then the estimated groundwater discharge from the Surficial Aquifer to the Bush River side of Beach Point can be calculated as shown below:

$$\begin{aligned}\text{Estimated Discharge Rate} &= (V_{hor})(\text{Area of Surficial Aquifer X-Section fronting Bush River}) \\ &= (0.89 \text{ ft/day})(61.8')(1,470') \\ &= 80,852.9 \text{ ft}^3/\text{day} = 604,780.0 \text{ gallons/day}\end{aligned}$$

2. However, a more realistic estimation of discharge through the Surficial Aquifer along the Bush River involves using the directional component of V_{hor} orthogonal to the Bush River frontage. If this is performed, then the estimated groundwater discharge from the Surficial Aquifer to the Bush River side of Beach Point can be calculated as shown below:

$$\begin{aligned}\text{Estimated Discharge Rate} &= (V_{hor})(\text{Area of Surficial Aquifer X-Section fronting Bush River})(\text{component of } V_{hor} \text{ orthogonal to Bush River frontage}) \\ &= (0.89 \text{ ft/day})(61.8')(1,200.3')(\sin 23.8) + (0.89 \text{ ft/day})(61.8')(270.3') \\ &= 41,508.7 \text{ ft}^3/\text{day} = 310,484.8 \text{ gallons/day}\end{aligned}$$